

Frobenius splitting of Schubert varieties of semi-infinite flag manifolds*

Syu KATO[†]

July 3, 2021

Abstract

We exhibit basic algebro-geometric results on the formal model of semi-infinite flag varieties and its Schubert varieties over an algebraically closed field \mathbb{K} of characteristic $\neq 2$ from scratch. We show that the formal model of a semi-infinite flag variety admits a unique nice (ind)scheme structure, its projective coordinate ring has a \mathbb{Z} -model, and it admits a Frobenius splitting compatible with the boundaries and opposite cells in positive characteristic. This establishes the normality of the Schubert varieties of the quasi-map space with a fixed degree (instead of their limits proved in [K, Math. Ann. **371** no.2 (2018)]) when $\text{char } \mathbb{K} = 0$ or $\gg 0$, and the higher cohomology vanishing of their nef line bundles in arbitrary characteristic $\neq 2$. Some particular cases of these results play crucial roles in our proof [K, arXiv:1805.01718] of a conjecture by Lam-Li-Mihalcea-Shimozono [J. Algebra **513** (2018)] that describes an isomorphism between affine and quantum K -groups of a flag manifold.

Introduction

The semi-infinite flag varieties are variants of affine flag varieties that encode the modular representation theory of a semi-simple Lie algebra, representation theory of a quantum group at roots of unity, and representation theory of an affine Lie algebra at the critical level. They originate from the ideas of Lusztig [64] and Drinfeld, put forward by Feigin-Frenkel [24], and subsequently polished by the works of Braverman, Finkelberg, and their collaborators [29, 23, 2, 9, 10, 11, 12, 13]. They (mainly) employed the ind-model of semi-infinite flag varieties and achieved spectacular success on the geometric Langlands correspondence [2, 7], on the quantum K -groups of flag manifolds [11], and on their (conjectural) relation to the finite \mathcal{W} -algebras [10].

In [51], we have initiated the study of the formal model of a semi-infinite flag variety (over \mathbb{C}) that follows the classical description of flag varieties [57, 69, 63, 58] more closely than the above. We refer this formal model of a semi-infinite flag variety as a “semi-infinite flag manifold” since we hope to justify that it is “smooth” in a sense. However, the analysis in [51] has two defects: the relation

*MSC2010: 20G44

[†]Department of Mathematics, Kyoto University, Oiwake Kita-Shirakawa Sakyo Kyoto 606-8502 JAPAN E-mail: syuchan@math.kyoto-u.ac.jp

with the ind-models of semi-infinite flag varieties is unclear, and the treatment there is rather ad hoc (it is just an indscheme whose set of \mathbb{C} -valued points have the desired property, and lacks a characterization as a functor cf. [3, 22]). The former defect produces difficulty in the discussion of deep properties on the identification between the equivariant K -group of a semi-infinite flag manifold and the equivariant quantum K -group of a flag manifold [47], that is in turn inspired by the works of Givental and Lee [32, 33]. The goal of this paper is to study semi-infinite flag manifolds in characteristic $\neq 2$ from scratch, and resolve the above defects. In particular, we verify that the scheme in [51] is the universal one among all the indschemes with similar set-theoretic properties, and provide new proofs of the normality of Zastava spaces [11] and of the semi-infinite flag manifolds [51].

It is possible to regard our works ([46, 49, 51, 47, 48]) as a part of catch-up of Peterson's original construction [74] of his isomorphism [61] between the quantum cohomology of a flag manifold and the cohomology of an affine Grassmannian in the K -theoretic setting. From this view-point, this paper provides some varieties considered in [74, Lecture 11] with their appropriate compactifications. Hence, though there are still some missing pieces to complete the original program along the lines in [74], this paper provides a step to fully examine his ideas.

To explain our results more precisely, we introduce more notation: Let \mathfrak{g} denote a simple Lie algebra (given in terms of root data and the Chevalley generators) over an algebraically closed field \mathbb{K} of characteristic $\neq 2$. Let G denote the connected simply-connected algebraic group over \mathbb{K} such that $\mathfrak{g} = \text{Lie } G$. Let $H \subset G$ be a Cartan subgroup and let N be a unipotent radical of G that is normalized by H . We set $B := HN$ and $\mathcal{B} := G/B$ (the flag manifold of G). Let $\mathbf{I}^+ \subset G(\mathbb{K}[[z]])$ denote the Iwahori subgroup that contains B , and let $\mathbf{I}^- \subset G(\mathbb{K}[[z^{-1}]])$ be its opposite Iwahori subgroup. Let $\tilde{\mathfrak{g}}$ denote the untwisted affine Kac-Moody algebra associated to \mathfrak{g} . Let W and W_{af} be the finite Weyl group and the affine Weyl group of \mathfrak{g} , respectively. The coroot lattice Q^\vee of \mathfrak{g} yields a natural subgroup $\{t_\beta\}_{\beta \in Q^\vee} \subset W_{\text{af}}$. Let $w_0 \in W$ be the longest element.

Our first main result is as follows:

Theorem A (\doteq Theorem 3.18 and Proposition 3.26). *There is an indscheme $\mathbf{Q}_G^{\text{rat}}$ with the following properties:*

1. *The indscheme $\mathbf{Q}_G^{\text{rat}}$ is expressed as the union of infinite type integral schemes flat over \mathbb{Z} ;*
2. *If we set $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}} := \mathbf{Q}_G^{\text{rat}} \otimes_{\mathbb{Z}} \mathbb{K}$, then we have*

$$(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}(\mathbb{K}) \cong G(\mathbb{K}((z)))/(H(\mathbb{K})N(\mathbb{K}((z))))$$

that intertwines the natural $G(\mathbb{K}((z))) \times \mathbb{G}_m(\mathbb{K})$ -actions on the both sides, where \mathbb{G}_m is the loop rotation;

3. *The functor*

$$\text{Aff}_{\mathbb{K}}^{\text{op}} \ni R \mapsto G(R((z)))/(H(R)N(R((z)))) \in \text{Sets.}$$

is coarsely ind-representable by $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}$ (see §3.3 for the convention).

One can equip $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}$ with an indscheme structure using the arc scheme of the basic affine space $\overline{G/N}$. Such an indscheme *cannot* coincide with ours (in general) by the appearance of the nontrivial nilradicals [71, 27, 26]. In fact, such an indscheme defines a radicial thickening of $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}$.

The set of $H \times \mathbb{G}_m$ -fixed points of $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}$ is in bijection with W_{af} . Let $p_w \in (\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}$ be the point corresponding to $w \in W_{\text{af}}$. We set $\mathbb{O}(w) := \mathbf{I}^+ p_w$ and $\mathbb{O}^-(w) := \mathbf{I}^- p_w$ for each $w \in W_{\text{af}}$.

Theorem A has some applications to the theory of quasi-map spaces from \mathbb{P}^1 to \mathcal{B} presented in [29, 23, 11, 12, 13] as follows:

Theorem B. *In the above settings, it holds:*

1. (Corollary 2.38 and Theorem 3.30) *If $\text{char } \mathbb{K} > 0$, then the scheme $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}$ admits an \mathbf{I}^{\pm} -canonical Frobenius splitting that is compatible with $\mathbb{O}(w)$'s and $\mathbb{O}^-(v)$'s ($w, v \in W_{\text{af}}$);*
2. (Lemma 3.10 and Theorem 3.13) *For each $w, v \in W_{\text{af}}$, the intersection $\mathcal{Q}(v, w) := \overline{\mathbb{O}(w)} \cap \overline{\mathbb{O}^-(v)}$ is reduced. It is irreducible when $v = w_0 t_{\beta}$ for some $\beta \in Q^{\vee}$;*
3. (Lemma 3.7 and Corollary 4.24) *For each $w, v \in W_{\text{af}}$, the scheme $\mathcal{Q}(v, w)$ is weakly normal. It is (irreducible and) normal when $\text{char } \mathbb{K} = 0$ or $\text{char } \mathbb{K} \gg 0$;*
4. (Lemma 3.28) *For each $\beta \in Q_+^{\vee}$, the set of \mathbb{K} -valued points of the scheme $\mathcal{Q}(w_0 t_{\beta}, e)$ is in bijection with the set of (\mathbb{K} -valued) Drinfeld-Plücker data. In particular, $\mathcal{Q}(w_0 t_{\beta}, e)$ is isomorphic to the quasi-map space in [29] when $\mathbb{K} = \mathbb{C}$.*

Theorem B is a key result at the deepest part (correspondence between natural bases) in our proof ([47]) of a conjecture of Lam-Li-Mihalcea-Shimozono [60] about the comparison between the equivariant K -group of the affine Grassmannian of G and the equivariant small quantum K -group of \mathcal{B} . In [47], we also prove that $\mathcal{Q}(w_0 t_{\beta}, w)$ admits only rational singularities (and hence it is Cohen-Macaulay) when $\mathbb{K} = \mathbb{C}$ on the basis of Theorem B. We remark that Theorem B 3) is proved in [11, 12] when $v = w_0 t_{\beta}$, $w = e$, and $\mathbb{K} = \mathbb{C}$.

Our proof of Theorem B 1) is not at all standard, and in fact it forms the core of the technical contributions in this paper. To appreciate its contents, let us recall that there are two standard ways to construct a Frobenius splitting of \mathcal{B} ([15]): One is to consider the Bott-Samelson-Demazure-Hansen resolution of \mathcal{B} , that reduces the assertion to the case of a point, which is a Schubert variety with trivial Frobenius splitting. Another is to analyze the space of global sections of (some power of) the canonical bundle of \mathcal{B} .

However, the both of the above proof strategies do not work for $\mathbf{Q}_G^{\text{rat}}$. The first one fails since any Schubert variety of $\mathbf{Q}_G^{\text{rat}}$ is infinite-dimensional, and carries rich internal structure by itself. The second one fails since the canonical bundle of $\mathbf{Q}_G^{\text{rat}}$ simply does not make sense, at least naively. These require us some new ideas in order to prove Theorem B 1). Our idea here is as follows: An interpretation of the filtrations in [50] reduces the existence of a Frobenius splitting of $\mathbf{Q}_G^{\text{rat}}$ into a property of the Frobenius splitting of the corresponding thick affine flag manifold [49, Corollary B]. This property can be seen as a special case of some homological property in the representation theory of affine

Lie algebras ([18, 20]), but it is proved only for characteristic zero. Thus, we use Kashiwara's theory of global basis ([43, 44]) to transfer such a homological property into the positive characteristic setting (Proposition 2.19).

In the rest of introduction, we assume $\mathbb{K} = \mathbb{C}$ for the sake of simplicity. Let P be the weight lattice of H , and let $P_+ \subset P$ denote its subset corresponding to dominant weights. For each $\lambda \in P$, we have an equivariant line bundle $\mathcal{O}_{\mathbf{Q}_G^{\text{rat}}}(\lambda)$ on $\mathbf{Q}_G^{\text{rat}}$, whose restriction to $\mathcal{Q}(v, w)$ is denoted by $\mathcal{O}_{\mathcal{Q}(v, w)}(\lambda)$. Associated to $\lambda \in P_+$, we have a level zero extremal weight module $\mathbb{X}(\lambda)$ of $U(\widehat{\mathfrak{g}})$ in the sense of Kashiwara [44]. We know that $\mathbb{X}(\lambda)$ is equipped with two kinds of Demazure modules, and a distinguished basis (the global basis).

Corollary C (\doteq Theorem 3.33). *Let $w, v \in W_{\text{af}}$. For each $\lambda \in P_+$, we have*

$$H^{>0}(\mathcal{Q}(v, w), \mathcal{O}_{\mathcal{Q}(v, w)}(\lambda)) = \{0\}.$$

The space $H^0(\mathcal{Q}(v, w), \mathcal{O}_{\mathcal{Q}(v, w)}(\lambda))^\vee$ is the intersection of two Demazure modules of $\mathbb{X}(\lambda)$ spanned by a subset of the global basis of $\mathbb{X}(\lambda)$ if λ is strictly dominant. If we have $w', v' \in W_{\text{af}}$ such that $\mathcal{Q}(v', w') \subset \mathcal{Q}(v, w)$, then the restriction map

$$H^0(\mathcal{Q}(v, w), \mathcal{O}_{\mathcal{Q}(v, w)}(\lambda)) \twoheadrightarrow H^0(\mathcal{Q}(v', w'), \mathcal{O}_{\mathcal{Q}(v', w')}(\lambda))$$

is surjective.

Note that Corollary C adds new vanishing region to [12, Theorem 3.1 1)]. We also provide parabolic versions of Theorem A, Theorem B, and Corollary C. We have a description of $H^0(\mathcal{Q}(v, w), \mathcal{O}_{\mathcal{Q}(v, w)}(\lambda))$ for general $\lambda \in P_+$ that is more complicated (Theorem B.6).

Let $\mathcal{B}_{2, \beta}$ be the space of genus zero stable maps with two marked points to \mathcal{B} with the class of its image $\beta \in Q_+^\vee \subset Q^\vee \cong H_2(\mathcal{B}, \mathbb{Z})$. We have evaluation maps $e_j : \mathcal{B}_{2, \beta} \rightarrow \mathcal{B}$ for $j = 1, 2$. The following purely geometric result is a byproduct of our proof, that maybe of independent interest.

Corollary D (\doteq Corollary 4.19). *Let $\beta \in Q_+^\vee$, and let $x, y \in \mathcal{B}$. The space $(e_1^{-1}(x) \cap e_2^{-1}(\overline{By}))$ is connected if it is nonempty.*

Note that Corollary D is contained in [16] whenever $\{x\}, \overline{By} \subset \mathcal{B}$ are in general position.

The plan of this paper is as follows: In section one, we collect basic material needed in the sequel. In section two, after recalling generalities on Frobenius splitting and representation theory of quantum loop algebras, we construct the indscheme $\mathbf{Q}_G^{\text{rat}}$ and equip it with a Frobenius splitting (Corollary 2.38). In section three, we first interpret $\mathbf{Q}_G^{\text{rat}}$ as an indscheme (coarsely) representing the coset $G(\mathbb{K}((z)))/(H(\mathbb{K})N(\mathbb{K}((z))))$ (Theorem A). Using this, we identify some Richardson varieties of $\mathbf{Q}_G^{\text{rat}}$ with quasi-map spaces (Theorem 3.30) and present their cohomological properties (Theorem 3.33), and hence prove (large parts of) Theorem B and Corollary C. Since our construction equips quasi-map spaces with Frobenius splittings (Lemma 3.7), they are automatically weakly normal. Moreover, we explain how to connect characteristic zero and positive characteristic (§3.5). In section four, we analyze the fibers of the graph space resolutions of quasi-map spaces and deduce that Richardson varieties of semi-infinite flag manifolds (over \mathbb{C}) are normal based on the weak normality proved in the previous section. This proves the remaining part of Theorem B. Our analysis here

contains an inductive proof that the fibers of the evaluation maps of the space of genus zero stable maps to flag varieties are connected (Corollary D). In Appendix A, we give a proof of the normality of (the ind-pieces of) $\mathbf{Q}_G^{\text{rat}}$ (that works also in the positive characteristic setting) and present an analogue of the Kempf vanishing theorem [52] for $\mathbf{Q}_G^{\text{rat}}$. Appendix B exhibits the structure of global sections of nef line bundles of Richardson varieties of $\mathbf{Q}_G^{\text{rat}}$.

Note that Theorem B equips a quasi-map space from \mathbb{P}^1 to \mathcal{B} with a Frobenius splitting compatible with the boundaries. However, the notion of boundary in quasi-map spaces depend on a configuration of points in \mathbb{P}^1 (we implicitly set them to $\{0, \infty\} \subset \mathbb{P}^1$ throughout this paper). This makes our analogues of open Richardson varieties not necessarily smooth contrary to the original case [76] (see also [29, §8.4.1]). The author hope to give more account of this, as well as the factorization structure ([29, §6.3]) from the view point presented in this paper, in his future works.

1 Preliminaries

We work over an algebraically closed field \mathbb{K} unless stated otherwise. A vector space is a \mathbb{K} -vector space, and a graded vector space refers to a \mathbb{Z} -graded vector space whose graded pieces are finite-dimensional and its grading is bounded from the above or from the below. Tensor products are taken over \mathbb{K} unless specified otherwise.

Let A be a PID. For a graded free A -module $M = \bigoplus_{m \in \mathbb{Z}} M_m$, we set $M^\vee := \bigoplus_{m \in \mathbb{Z}} \text{Hom}_A(M_m, A)$, where $\text{Hom}_A(M_m, A)$ is understood to be degree $-m$.

As a rule, we suppress \emptyset and associated parenthesis from notation. This particularly applies to $\emptyset = \mathbf{J} \subset \mathbf{I}$ frequently used to specify parabolic subgroups.

1.1 Groups, root systems, and Weyl groups

We refer to [21, 57] for precise expositions of general material presented in this subsection.

Let G be a connected, simply connected simple algebraic group of rank r over an algebraically closed field \mathbb{K} , and let B and H be a Borel subgroup and a maximal torus of G such that $H \subset B$. We set $N (= [B, B])$ to be the unipotent radical of B and let N^- be the opposite unipotent subgroup of N with respect to H . We denote the Lie algebra of an algebraic group by the corresponding German small letter. We have a (finite) Weyl group $W := N_G(H)/H$. For an algebraic group E , we denote its set of $\mathbb{K}[z]$ -valued points by $E[z]$, its set of $\mathbb{K}[[z]]$ -valued points by $E[[z]]$, and its set of $\mathbb{K}((z))$ -valued points by $E((z))$ etc... Let $\mathbf{I} \subset G[[z]]$ be the preimage of $B \subset G$ via the evaluation at $z = 0$ (the Iwahori subgroup of $G[[z]]$). We set $\mathbf{I}^- \subset G[[z^{-1}]]$ be the opposite Iwahori subgroup of \mathbf{I} in $G((z))$ with respect to H . By abuse of notation, we might consider \mathbf{I} and $G[[z]]$ as pro-algebraic groups over \mathbb{K} whose \mathbb{K} -valued points are given as these.

Let $P := \text{Hom}_{gr}(H, \mathbb{G}_m)$ be the weight lattice of H , let $\Delta \subset P$ be the set of roots, let $\Delta_+ \subset \Delta$ be the set of roots that yield root subspaces in \mathfrak{b} , and let $\Pi \subset \Delta_+$ be the set of simple roots. We set $\Delta_- := -\Delta_+$. Let Q^\vee be the dual lattice of P with a natural pairing $\langle \bullet, \bullet \rangle : Q^\vee \times P \rightarrow \mathbb{Z}$. We define $\Pi^\vee \subset Q^\vee$ to be the set of positive simple coroots, and let $Q_+^\vee \subset Q^\vee$ be the set of non-negative

integer span of Π^\vee . For $\beta, \gamma \in Q^\vee$, we define $\beta \geq \gamma$ if and only if $\beta - \gamma \in Q_+^\vee$. We set $P_+ := \{\lambda \in P \mid \langle \alpha^\vee, \lambda \rangle \geq 0, \forall \alpha^\vee \in \Pi^\vee\}$ and $P_{++} := \{\lambda \in P \mid \langle \alpha^\vee, \lambda \rangle > 0, \forall \alpha^\vee \in \Pi^\vee\}$. Let $\mathbf{I} := \{1, 2, \dots, r\}$. We fix bijections $\mathbf{I} \cong \Pi \cong \Pi^\vee$ such that $i \in \mathbf{I}$ corresponds to $\alpha_i \in \Pi$, its coroot $\alpha_i^\vee \in \Pi^\vee$, and a simple reflection $s_i \in W$ corresponding to α_i . Let $\{\varpi_i\}_{i \in \mathbf{I}} \subset P_+$ be the set of fundamental weights (i.e. $\langle \alpha_i^\vee, \varpi_j \rangle = \delta_{ij}$) and $\rho := \sum_{i \in \mathbf{I}} \varpi_i = \frac{1}{2} \sum_{\alpha \in \Delta^+} \alpha \in P_+$.

For a subset $\mathbf{J} \subset \mathbf{I}$, we define $P(\mathbf{J})$ as the standard parabolic subgroup of G corresponding to \mathbf{J} . I.e. we have $\mathfrak{b} \subset \mathfrak{p}(\mathbf{J}) \subset \mathfrak{g}$ and $\mathfrak{p}(\mathbf{J})$ contains the root subspace corresponding to $-\alpha_i$ ($i \in \mathbf{I}$) if and only if $i \in \mathbf{J}$. Let $H \subset L(\mathbf{J}) \subset P(\mathbf{J})$ be the standard Levi subgroup (that is isomorphic to the quotient of $P(\mathbf{J})$ by its unipotent radical). Then, the set of characters of $P(\mathbf{J})$ is identified with $P_{\mathbf{J}} := \sum_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{Z} \varpi_i$. We also set $P_{\mathbf{J},+} := \sum_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{Z}_{\geq 0} \varpi_i = P_+ \cap P_{\mathbf{J}}$ and $P_{\mathbf{J},++} := \sum_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{Z}_{\geq 1} \varpi_i = P_{++} \cap P_{\mathbf{J}}$. We define $W_{\mathbf{J}} \subset W$ to be the reflection subgroup generated by $\{s_i\}_{i \in \mathbf{J}}$. It is the Weyl group of $[L(\mathbf{J}), L(\mathbf{J})]$ and $L(\mathbf{J})$. We define $\rho_{\mathbf{J}}$ to be the half-sum of positive roots whose root spaces are contained in the unipotent radical of $\mathfrak{p}(\mathbf{J})$.

Let $\Delta_{\text{af}} := \Delta \times \mathbb{Z}\delta \cup \{m\delta\}_{m \neq 0}$ be the untwisted affine root system of Δ with its positive part $\Delta_+ \subset \Delta_{\text{af},+}$. We set $\alpha_0 := -\vartheta + \delta$, $\Pi_{\text{af}} := \Pi \cup \{\alpha_0\}$, and $\mathbf{I}_{\text{af}} := \mathbf{I} \cup \{0\}$, where ϑ is the highest root of Δ_+ . We set $W_{\text{af}} := W \rtimes Q^\vee$ and call it the affine Weyl group. It is a reflection group generated by $\{s_i \mid i \in \mathbf{I}_{\text{af}}\}$, where s_0 is the reflection with respect to α_0 . We also have a reflection $s_\alpha \in W_{\text{af}}$ corresponding to $\alpha \in \Delta \times \mathbb{Z}\delta \subsetneq \Delta_{\text{af}}$. Let $\ell : W_{\text{af}} \rightarrow \mathbb{Z}_{\geq 0}$ be the length function and let $w_0 \in W$ be the longest element in $W \subset \bar{W}_{\text{af}}$. Together with the normalization $t_{-\vartheta^\vee} := s_\vartheta s_0$ (for the coroot ϑ^\vee of ϑ), we introduce the translation element $t_\beta \in W_{\text{af}}$ for each $\beta \in Q^\vee$.

For each $i \in \mathbf{I}_{\text{af}}$, we have a connected algebraic group $SL(2, i)$ that is isomorphic to $SL(2)$ equipped with an inclusion $SL(2, i)(\mathbb{K}) \subset G((z))$ as groups corresponding to $\pm\alpha_i \in \mathbf{I}_{\text{af}}$. Let $\rho_{\pm\alpha_i} : \mathbb{G}_m \rightarrow SL(2, i)$ denote the unipotent one-parameter subgroup corresponding to $\pm\alpha_i \in \Delta_{\text{af}}$. We set $B_i := SL(2, i) \cap \mathbf{I}$, that is a Borel subgroup of $SL(2, i)$. For each $i \in \mathbf{I}$, we set $P_i := P(\{i\})$. For each $i \in \mathbf{I}_{\text{af}}$, we set $\mathbf{I}(i) := SL(2, i)\mathbf{I}$. Each $\mathbf{I}(i)$ can be regarded as a pro-algebraic group.

As a variation of [57, Chapter VI], we say an indscheme \mathfrak{X} over \mathbb{K} admits a $G((z))$ -action if it admits an action of \mathbf{I} and $SL(2, i)$ ($i \in \mathbf{I}_{\text{af}}$) as (ind)schemes over \mathbb{K} that coincides on $B_i = (\mathbf{I} \cap SL(2, i))$, and they generate a $G((z))$ -action on the set of closed points of \mathfrak{X} (the latter is a group action on a set). We consider the notion of $G((z))$ -equivariant morphisms accordingly.

We set

$$Q_{<}^\vee := \{\beta \in Q^\vee \mid \langle \beta, \alpha_i \rangle < 0, \forall i \in \bar{\mathbf{I}}\}.$$

Let \leq be the Bruhat order of W_{af} . In other words, $w \leq v$ holds if and only if a subexpression of a reduced decomposition of v yields a reduced decomposition of w . We define the generic (semi-infinite) Bruhat order $\leq_{\frac{\infty}{2}}$ as:

$$w \leq_{\frac{\infty}{2}} v \Leftrightarrow wt_\beta \leq vt_\beta \quad \text{for every } \beta \in Q_{<}^\vee \text{ such that } \langle \beta, \alpha_i \rangle \ll 0 \text{ for } i \in \mathbf{I}. \quad (1.1)$$

By [64], this defines a preorder on W_{af} . Here we remark that $w \leq v$ if and only if $w \geq_{\frac{\infty}{2}} v$ for $w, v \in W$. We also have

$$w \leq_{\frac{\infty}{2}} v \Leftrightarrow ww_0 \geq_{\frac{\infty}{2}} vw_0 \quad w, v \in W_{\text{af}}. \quad (1.2)$$

For proofs and related results, we refer to [51, §2.2] and [74, Lecture 13].

For each $u \in W$ and $\beta \in Q^\vee$, we set

$$\ell^{\frac{\infty}{2}}(ut_\beta) := \ell(u) + \sum_{\alpha \in \Delta_+} \langle \beta, \alpha \rangle = \ell(u) + 2 \langle \beta, \rho \rangle.$$

Theorem 1.1 (Lusztig [64] cf. [61] Proposition 4.4). *For each $w, v \in W_{\text{af}}$ such that $w \leq_{\frac{\infty}{2}} v$, there exists $\alpha \in \Delta_+^{\text{af}}$ such that $w \leq_{\frac{\infty}{2}} s_\alpha v \leq_{\frac{\infty}{2}} v$ and $\ell^{\frac{\infty}{2}}(s_\alpha v) = \ell^{\frac{\infty}{2}}(v) + 1$. \square*

For each $\lambda \in P_+$, we denote the corresponding Weyl module by $V(\lambda)$ (see e.g. [1, Proposition 1.22] and [41, Theorem 5]). By convention, $V(\lambda)$ is a finite-dimensional indecomposable G -module with a cyclic B -eigenvector \mathbf{v}_λ^0 (highest weight vector) with its H -weight λ whose character obeys the Weyl character formula. For a semi-simple H -module V , we set

$$\text{ch } V := \sum_{\lambda \in P} e^\lambda \cdot \dim_{\mathbb{K}} \text{Hom}_H(\mathbb{K}_\lambda, V).$$

If V is a \mathbb{Z} -graded H -module in addition, then we set

$$\text{gch } V := \sum_{\lambda \in P, n \in \mathbb{Z}} q^n e^\lambda \cdot \dim_{\mathbb{K}} \text{Hom}_H(\mathbb{K}_\lambda, V_n). \quad (1.3)$$

Let $\mathcal{B} := G/B$ and call it the flag manifold of G . We have the Bruhat decomposition

$$\mathcal{B} = \bigsqcup_{w \in W} \mathbb{O}_{\mathcal{B}}(w) \quad (1.4)$$

into B -orbits such that $\dim \mathbb{O}_{\mathcal{B}}(w) = \ell(w_0) - \ell(w)$ for each $w \in W \subset W_{\text{af}}$. We set $\mathcal{B}(w) := \overline{\mathbb{O}_{\mathcal{B}}(w)} \subset \mathcal{B}$.

For each $\lambda \in P$, we have a line bundle $\mathcal{O}_{\mathcal{B}}(\lambda)$ such that

$$H^0(\mathcal{B}, \mathcal{O}_{\mathcal{B}}(\lambda)) \cong V(\lambda)^*, \quad \mathcal{O}_{\mathcal{B}}(\lambda) \otimes_{\mathcal{O}_{\mathcal{B}}} \mathcal{O}_{\mathcal{B}}(-\mu) \cong \mathcal{O}_{\mathcal{B}}(\lambda - \mu) \quad \lambda, \mu \in P_+.$$

For each $w \in W$, let $p_w \in \mathbb{O}_{\mathcal{B}}(w)$ be the unique H -fixed point. We normalize p_w (and hence $\mathbb{O}_{\mathcal{B}}(w)$) so that the restriction of $\mathcal{O}_{\mathcal{B}}(\lambda)$ to p_w is isomorphic to $\mathbb{K}_{-ww_0\lambda}$ for every $\lambda \in P_+$. (Here we warn that the convention differs from [47].)

1.2 Representations of affine and current algebras

In the rest of this section, we work over $\mathbb{K} = \mathbb{C}$, the field of complex numbers. Material in this subsection without a reference can be found in [40, 41]. Every result in this subsection is transferred to an arbitrary field in §2.2.

Let $\tilde{\mathfrak{g}}$ denote the untwisted affine Kac-Moody algebra associated to \mathfrak{g} . I.e. we have

$$\tilde{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[z, z^{-1}] \oplus \mathbb{C}K \oplus \mathbb{C}d,$$

where K is central, $[d, X \otimes z^m] = mX \otimes z^m$ for each $X \in \mathfrak{g}$ and $m \in \mathbb{Z}$, and for each $X, Y \in \mathfrak{g}$ and $f, g \in \mathbb{C}[z^{\pm 1}]$ it holds:

$$[X \otimes f, Y \otimes g] = [X, Y] \otimes fg + (X, Y)_{\mathfrak{g}} \cdot K \cdot \text{Res}_{z=0} f \frac{\partial g}{\partial z},$$

where $(\bullet, \bullet)_{\mathfrak{g}}$ denotes the G -invariant bilinear form such that $(\alpha^\vee, \alpha^\vee)_{\mathfrak{g}} = 2$ for a long simple root α . Let E_i, F_i ($i \in \mathbf{I}_{\text{af}}$) denote the Kac-Moody generators of $\tilde{\mathfrak{g}}$ corresponding to α_i . We set $\tilde{\mathfrak{h}} := \mathfrak{h} \oplus \mathbb{C}K \oplus \mathbb{C}d$. Let \mathfrak{J} be the Lie subalgebra of $\tilde{\mathfrak{g}}$ generated by E_i ($i \in \mathbf{I}_{\text{af}}$) and $\tilde{\mathfrak{h}}$, and \mathfrak{J}^- be the Lie subalgebra of $\tilde{\mathfrak{g}}$ generated by F_i ($i \in \mathbf{I}_{\text{af}}$) and $\tilde{\mathfrak{h}}$. For each $i \in \mathbf{I}_{\text{af}}$ and $n \geq 0$, we set $E_i^{(n)} := \frac{1}{n!} E_i^n$ and $F_i^{(n)} := \frac{1}{n!} F_i^n$.

We define

$$Q^{\text{af}, \vee} := \mathbb{Z}d \oplus \bigoplus_{i \in \mathbf{I}_{\text{af}}} \mathbb{Z}\alpha_i^\vee \subset \tilde{\mathfrak{h}}, \quad P^{\text{af}} := \mathbb{Z}\delta \oplus \bigoplus_{i \in \mathbf{I}_{\text{af}}} \mathbb{Z}\Lambda_i \subset \tilde{\mathfrak{h}}^*,$$

and a pairing $Q^{\text{af}, \vee} \times P^{\text{af}} \rightarrow \mathbb{Z}$ such that

$$\langle \alpha_i^\vee, \Lambda_j \rangle = \delta_{ij} \quad (i, j \in \mathbf{I}_{\text{af}}), \quad \langle \alpha_i^\vee, \delta \rangle \equiv 0, \quad \langle d, \Lambda_i \rangle = \delta_{i0} \quad (i \in \mathbf{I}_{\text{af}}), \quad \langle d, \delta \rangle = 1.$$

We have a projection map

$$P^{\text{af}} \ni \Lambda = k\delta + \sum_{i \in \mathbf{I}_{\text{af}}} a_i \Lambda_i \mapsto \bar{\Lambda} = \sum_{i \in \mathbf{I}} a_i \varpi_i \in P,$$

that has a unique splitting $P \subset P^{\text{af}}$ whose image is orthogonal to $d, K \in \tilde{\mathfrak{h}}$. We set $P_+^{\text{af}} := \sum_{i \in \mathbf{I}_{\text{af}}} \mathbb{Z}_{\geq 0} \Lambda_i$. Each $\Lambda \in P_+^{\text{af}}$ defines an irreducible integrable highest weight module $L(\Lambda)$ of $\tilde{\mathfrak{g}}$ with its highest weight vector \mathbf{v}_Λ . In addition, each $\lambda \in P_+$ defines a level zero extremal weight module $\mathbb{X}(\lambda)$ of $\tilde{\mathfrak{g}}$ by means of the specialization of the quantum parameter $q = 1$ in [43, Proposition 8.2.2] and [44, §5.1], that is integrable and K acts by 0. The module $\mathbb{X}(\lambda)$ carries a cyclic \mathfrak{h} -weight vector \mathbf{v}_λ such that:

$$H\mathbf{v}_\lambda = \lambda(H)\mathbf{v}_\lambda \quad (H \in \mathfrak{h}), \quad K\mathbf{v}_\lambda = 0 = d\mathbf{v}_\lambda, \quad E_i\mathbf{v}_\lambda = 0 \quad (i \in \mathbf{I}), \quad \text{and} \quad F_0\mathbf{v}_\lambda = 0.$$

(We can deduce that $\mathbb{X}(\lambda)$ is the maximal integrable $\tilde{\mathfrak{g}}$ -module that possesses a cyclic vector with the above properties [43, §8.1].) Moreover, each $w = ut_\beta \in W_{\text{af}}$ ($u \in W, \beta \in Q^\vee$) defines an element $\mathbf{v}_{w\lambda} \in \mathbb{X}(\lambda)$ such that

$$H\mathbf{v}_{w\lambda} = (w\lambda)(H)\mathbf{v}_{w\lambda} \quad (H \in \mathfrak{h}), \quad K\mathbf{v}_{w\lambda} = 0, \quad d\mathbf{v}_{w\lambda} = -\langle \beta, \lambda \rangle \mathbf{v}_{w\lambda}$$

up to sign (see [43, §8.1]). We call a vector in $\{\mathbf{v}_{w\lambda}\}_{w \in W_{\text{af}}}$ an extremal weight vector of $\mathbb{X}(\lambda)$.

We set $\mathfrak{g}[z] := \mathfrak{g} \otimes_{\mathbb{C}} \mathbb{C}[z]$ and regard it as a Lie subalgebra of $\tilde{\mathfrak{g}}$. We have $\mathfrak{J} \subset \mathfrak{g}[z] + \mathbb{C}K + \mathbb{C}d$. The Lie algebra $\mathfrak{g}[z]$ is graded, and its grading is the internal grading of $\tilde{\mathfrak{g}}$ given by d .

For each $\lambda \in P_+$, we set

$$\mathbb{W}_w(\lambda) := U(\mathfrak{J})\mathbf{v}_{w\lambda} \subset \mathbb{X}(\lambda).$$

These are the $q = 1$ cases of the Demazure modules of $\mathbb{X}(\lambda)$, as well as the generalized global Weyl modules in the sense of [28]. We set $\mathbb{W}(\lambda) := \mathbb{W}_{w_0}(\lambda)$. By construction, the both of $\mathbb{X}(\lambda)$ and $\mathbb{W}_w(\lambda)$ are semi-simple as $(H \times \mathbb{G}_m)$ -module, where \mathbb{G}_m acts on z by $a : z^m \mapsto a^m z^m$ ($m \in \mathbb{Z}$).

Theorem 1.2 (LNSSS [62], Chari-Ion [18], cf. [46] Theorem 1.6). *For each $\lambda \in P_+$, the \mathfrak{J} -action on $\mathbb{W}(\lambda)$ prolongs to $\mathfrak{g}[z]$, and it is isomorphic to the global Weyl module of $\mathfrak{g}[z]$ in the sense of Chari-Pressley [19]. Moreover, $\mathbb{W}(\lambda)$ is a projective module in the category of $\mathfrak{g}[z]$ -modules whose restriction to \mathfrak{g} is a direct sum of modules in $\{V(\mu)\}_{\mu \leq \lambda}$. \square*

Theorem 1.3 ([46] Theorem 3.3 and Corollary 3.4). *Let $\lambda, \mu \in P_+$ and $w \in W$. We have a unique (up to scalar) injective degree zero \mathfrak{J} -module map*

$$\mathbb{W}_w(\lambda + \mu) \hookrightarrow \mathbb{W}_w(\lambda) \otimes \mathbb{W}_w(\mu).$$

Sketch of proof. For each $\lambda, \mu \in P_+$, the projectivity of $\mathbb{W}(\lambda + \mu)$ in the sense of Theorem 1.2 yields a unique graded $\mathfrak{g}[z]$ -module map

$$\mathbb{W}(\lambda + \mu) \longrightarrow \mathbb{W}(\lambda) \otimes \mathbb{W}(\mu)$$

of degree 0. This map is injective by examining the specializations to local Weyl modules (for their definitions, see [46, Theorem 1.4], or Lemma 2.20 and Remark 2.21). By examining the \mathfrak{J} -cyclic vectors, it uniquely restricts to a map

$$\mathbb{W}_w(\lambda + \mu) \longrightarrow \mathbb{W}_w(\lambda) \otimes \mathbb{W}_w(\mu)$$

up to scalar. This map must be also injective as the ambient map is so. \square

1.3 Semi-infinite flag manifolds

We work over \mathbb{C} as in the previous subsection. Material in this section is reproved in the setting of characteristic $\neq 2$ in §2.4 and §3.2 (cf. §3.3). We define the semi-infinite flag manifold as the reduced indscheme such that:

- We have a closed embedding

$$\mathbf{Q}_G^{\text{rat}} \subset \prod_{i \in \mathbf{I}} \mathbb{P}(V(\varpi_i) \otimes \mathbb{C}((z))); \quad \text{and}$$

- We have an equality $\mathbf{Q}_G^{\text{rat}}(\mathbb{C}) = G((z))/ (H(\mathbb{C}) \cdot N((z)))$.

This is a pure indscheme of ind-infinite type [51]. Note that the group $Q^\vee \subset H((z))/H(\mathbb{C})$ acts on $\mathbf{Q}_G^{\text{rat}}$ from the right. The indscheme $\mathbf{Q}_G^{\text{rat}}$ is equipped with a $G((z))$ -equivariant line bundle $\mathcal{O}_{\mathbf{Q}_G^{\text{rat}}}(\lambda)$ for each $\lambda \in P$. Here we normalized so that $\Gamma(\mathbf{Q}_G^{\text{rat}}, \mathcal{O}_{\mathbf{Q}_G^{\text{rat}}}(\lambda))$ is $B^-((z))$ -cocyclic to a H -weight vector with its H -weight $-\lambda$. We warn that this convention is twisted by $-w_0$ from that of [47], and complies with [51].

Theorem 1.4 ([29, 23, 51, 64]). *We have an \mathbf{I} -orbit decomposition*

$$\mathbf{Q}_G^{\text{rat}} = \bigsqcup_{w \in W_{\text{af}}} \mathbb{O}(w)$$

with the following properties:

1. Each $\mathbb{O}(w)$ is isomorphic to \mathbb{A}^∞ and have a unique $(H \times \mathbb{G}_m)$ -fixed point;
2. The right action of $\gamma \in Q^\vee$ on $\mathbf{Q}_G^{\text{rat}}$ yields the translation $\mathbb{O}(w) \mapsto \mathbb{O}(wt_\gamma)$;
3. We have $\mathbb{O}(w) \subset \overline{\mathbb{O}(v)}$ if and only if $w \leq_{\frac{\infty}{2}} v$;
4. The relative dimension of $\mathbb{O}(ut_\beta)$ ($u \in W, \beta \in Q^\vee$) and $\mathbb{O}(e)$, counted as the difference of the cardinality of the maximal chain of intermediate \mathbf{I} -orbits to a common smaller \mathbf{I} -orbits, is $\ell_{\frac{\infty}{2}}(ut_\beta)$. \square

For each $w \in W_{\text{af}}$, let $\mathbf{Q}_G(w)$ denote the closure of $\mathbb{O}(w)$. We refer $\mathbf{Q}_G(w)$ as a Schubert variety of $\mathbf{Q}_G^{\text{rat}}$ (corresponding to $w \in W_{\text{af}}$).

Let $S = \bigoplus_{\lambda \in P_{j,+}} S(\lambda)$ be a $P_{j,+}$ -graded commutative ring such that $S(0) = A$ is a PID, S is torsion-free over A , and S is generated by $\bigoplus_{i \in \mathbb{I} \setminus \mathbb{J}} S(\varpi_i)$. We define

$$\text{Proj } S = (\text{Spec } S \setminus E)/H \subset \prod_{i \in \mathbb{I} \setminus \mathbb{J}} \mathbb{P}_A(S(\varpi_i)^\vee) \quad (1.5)$$

as the $P_{j,+}$ -graded proj over $\text{Spec } A$, where E is the locus that whole of $S(\varpi_i)$ vanishes for some $i \in \mathbb{I} \setminus \mathbb{J}$ (the irrelevant locus).

Theorem 1.5 ([51] Theorem 4.26 and Corollary 4.27). *For each $w \in W_{\text{af}}$, it holds:*

$$\mathbf{Q}_G(w) \cong \text{Proj} \bigoplus_{\lambda \in P_+} \mathbb{W}_{ww_0}(\lambda)^\vee,$$

where the multiplication of the ring $\bigoplus_{\lambda} \mathbb{W}_{ww_0}(\lambda)^\vee$ is given by Theorem 1.3.

1.4 Quasi-map spaces and Zastava spaces

We work over \mathbb{C} as in the previous subsection. Here we recall basics of quasi-map spaces from [29, 23].

We have W -equivariant isomorphisms $H^2(\mathcal{B}, \mathbb{Z}) \cong P$ and $H_2(\mathcal{B}, \mathbb{Z}) \cong Q^\vee$. This identifies the (integral points of the) nef cone of \mathcal{B} with $P_+ \subset P$ and the effective cone of \mathcal{B} with Q_+^\vee . A quasi-map (f, D) is a map $f : \mathbb{P}^1 \rightarrow \mathcal{B}$ together with a Π^\vee -colored effective divisor

$$D = \sum_{\alpha \in \Pi^\vee, x \in \mathbb{P}^1(\mathbb{C})} m_x(\alpha^\vee) \alpha^\vee \otimes [x] \in Q^\vee \otimes_{\mathbb{Z}} \text{Div } \mathbb{P}^1 \quad \text{with } m_x(\alpha^\vee) \in \mathbb{Z}_{\geq 0}.$$

For $i \in \mathbb{I}$, we set $D_i := \langle D, \varpi_i \rangle \in \text{Div } \mathbb{P}^1$. We call D the defect of the quasi-map (f, D) . Here we define the (total) degree of the defect by

$$|D| := \sum_{\alpha \in \Pi^\vee, x \in \mathbb{P}^1(\mathbb{C})} m_x(\alpha^\vee) \alpha^\vee \in Q_+^\vee.$$

For each $\beta \in Q_+^\vee$, we set

$$\mathcal{Q}(\mathcal{B}, \beta) := \{f : \mathbb{P}^1 \rightarrow X \mid \text{quasi-map s.t. } f_*[\mathbb{P}^1] + |D| = \beta\},$$

where $f_*[\mathbb{P}^1]$ is the class of the image of \mathbb{P}^1 multiplied by the degree of $\mathbb{P}^1 \rightarrow \text{Im } f$. We denote $\mathcal{Q}(\mathcal{B}, \beta)$ by $\mathcal{Q}(\beta)$ in case there is no danger of confusion.

Definition 1.6 (Drinfeld-Plücker data). Consider a collection $\mathcal{L} = \{(\psi_\lambda, \mathcal{L}^\lambda)\}_{\lambda \in P_+}$ of inclusions $\psi_\lambda : \mathcal{L}^\lambda \hookrightarrow V(\lambda) \otimes_{\mathbb{C}} \mathcal{O}_{\mathbb{P}^1}$ of line bundles \mathcal{L}^λ over \mathbb{P}^1 . The data \mathcal{L} is called a Drinfeld-Plücker data (DP-data) if the canonical inclusion of G -modules

$$\eta_{\lambda, \mu} : V(\lambda + \mu) \hookrightarrow V(\lambda) \otimes V(\mu)$$

induces an isomorphism

$$\eta_{\lambda, \mu} \otimes \text{id} : \psi_{\lambda+\mu}(\mathcal{L}^{\lambda+\mu}) \xrightarrow{\cong} \psi_\lambda(\mathcal{L}^\lambda) \otimes_{\mathcal{O}_{\mathbb{P}^1}} \psi_\mu(\mathcal{L}^\mu)$$

for every $\lambda, \mu \in P_+$.

Theorem 1.7 (Drinfeld, see Finkelberg-Mirković [29]). *The variety $\mathcal{Q}(\beta)$ is isomorphic to the variety formed by isomorphism classes of the DP-data $\mathcal{L} = \{(\psi_\lambda, \mathcal{L}^\lambda)\}_{\lambda \in P_+}$ such that $\deg \mathcal{L}^\lambda = \langle w_0\beta, \lambda \rangle$. In addition, $\mathcal{Q}(\beta)$ is an irreducible variety of dimension $2\langle \rho, \beta \rangle + \ell(w_0)$.*

For each $w \in W$, let $\mathcal{Z}(\beta, w) \subset \mathcal{Q}(\beta)$ be the locally closed subset consisting of quasi-maps that are defined at $z = 0$, and their values at $z = 0$ are contained in $\mathcal{B}(w) \subset \mathcal{B}$. We set $\mathcal{Q}(\beta, w) := \overline{\mathcal{Z}(\beta, w)}$. (Hence, we have $\mathcal{Q}(\beta) = \mathcal{Q}(\beta, e)$.)

Theorem 1.8 (Finkelberg-Mirković [29]). *Let \mathbb{K} be an algebraically closed field (that is not necessarily characteristic zero), and let $\mathcal{Q}(\beta)_{\mathbb{K}}$ and $\mathcal{Z}(\beta, w_0)_{\mathbb{K}}$ be the spaces obtained by replacing the base field \mathbb{C} with \mathbb{K} in Definition 1.6. For each $\beta \in Q_+^\vee$, the space $\mathcal{Z}(\beta, w_0)_{\mathbb{K}}$ is an irreducible affine scheme equipped with an action of $(B \times \mathbb{G}_m)$ over \mathbb{K} . In addition, this action has a unique fixed point.*

Remarks on proof. Theorem 1.8 is proved in [29] for $\mathbb{K} = \mathbb{C}$ using [70], and is proved in the current setting in [7] using [8]. One can also replace the usage of [70] with [81, Corollary 5.3.8] along the lines of [29]. \square

For each $\lambda \in P$ and $w \in W$, we have a G -equivariant line bundle $\mathcal{O}_{\mathcal{Q}(\beta, w)}(\lambda)$ (and its pro-object $\mathcal{O}_{\mathcal{Q}}(\lambda)$) obtained by the (tensor products of the) pull-backs $\mathcal{O}_{\mathcal{Q}(\beta, w)}(\varpi_i)$ of the i -th $\mathcal{O}(1)$ via the embedding

$$\mathcal{Q}(\beta, w) \hookrightarrow \prod_{i \in I} \mathbb{P}(V(\varpi_i) \otimes_{\mathbb{C}} \mathbb{C}[z]_{\leq -\langle w_0\beta, \varpi_i \rangle}) \quad (1.6)$$

for each $\beta \in Q_+^\vee$.

We have embeddings $\mathcal{B} \subset \mathcal{Q}(\beta) \subset \mathbf{Q}_G(e)$ such that the line bundles $\mathcal{O}(\lambda)$ ($\lambda \in P$) correspond to each other by restrictions ([12, 46, 51]).

2 Semi-infinite flag manifolds over $\mathbb{Z}[\frac{1}{2}]$

We keep the settings of the previous section. In this section, we sometimes work over a (commutative) ring or a non-algebraically closed field. For a ring S or a scheme \mathfrak{X} , we may write S_A and \mathfrak{X}_A if it is defined over A . In addition, we may consider their scalar extensions $S_B := S_A \otimes_A B$ and \mathfrak{X}_B for a ring map $A \rightarrow B$.

2.1 Frobenius splittings

Let \mathbb{k} be a field, and let p be a prime. We assume $\text{char } \mathbb{k} = p$, $\mathbb{k} \subset \mathbb{K}$, and the p -th power map is invertible on \mathbb{k} (e.g. $\mathbb{k} = \mathbb{F}_p$ or $\overline{\mathbb{F}_p}$) throughout this subsection.

We follow the generality on Frobenius splittings in [15], that considers separated schemes of finite type. We sometimes use the assertions from [15] without finite type assumption when the assertion is independent of that, whose typical disguises are properness, finite generation, and the Serre vanishing theorem.

Definition 2.1 (Frobenius splitting of a ring). Let R be a commutative ring over \mathbb{k} , and let $R^{(1)}$ denote the set R equipped with the map

$$R \times R^{(1)} \ni (r, m) \mapsto r^p m \in R^{(1)}.$$

This equips $R^{(1)}$ an R -module structure over \mathbb{k} (the \mathbb{k} -vector space structure on $R^{(1)}$ is also twisted by the p -th power operation), together with a map

$\iota : R.1 \rightarrow R^{(1)}$. An R -module map $\phi : R^{(1)} \rightarrow R$ is said to be a Frobenius splitting if $\phi \circ \iota$ is an identity.

Note that ι in Definition 2.1 must be an inclusion if we have a Frobenius splitting ϕ . Since the p -th power map in \mathbb{k} is invertible, we can twist the (scalar multiplication part of the) \mathbb{k} -vector space structure of R ($\cong R^{(1)}$ as sets) to make ι into a \mathbb{k} -linear map, without making it into an R -linear map (when $R \neq \mathbb{k}$).

Definition 2.2 (Frobenius splitting of a scheme). Let \mathfrak{X} be a separated scheme defined over \mathbb{k} . Let Fr be the (relative) Frobenius endomorphism of \mathfrak{X} (that induces a \mathbb{k} -linear endomorphism). We have a natural inclusion $\iota : \mathcal{O}_{\mathfrak{X}} \rightarrow \text{Fr}_* \mathcal{O}_{\mathfrak{X}}$. A Frobenius splitting of \mathfrak{X} is a $\mathcal{O}_{\mathfrak{X}}$ -linear morphism $\phi : \text{Fr}_* \mathcal{O}_{\mathfrak{X}} \rightarrow \mathcal{O}_{\mathfrak{X}}$ such that the composition $\phi \circ \iota$ is the identity.

Definition 2.3 (Compatible splitting). Let $\mathfrak{Y} \subset \mathfrak{X}$ be a closed immersion of separated schemes defined over \mathbb{k} . A Frobenius splitting ϕ of \mathfrak{X} is said to be compatible with \mathfrak{Y} if $\phi(\text{Fr}_* \mathcal{I}_{\mathfrak{Y}}) \subset \mathcal{I}_{\mathfrak{Y}}$, where $\mathcal{I}_{\mathfrak{Y}} := \ker(\mathcal{O}_{\mathfrak{X}} \rightarrow \mathcal{O}_{\mathfrak{Y}})$. Compatible Frobenius splitting of a pair of a commutative ring and its quotient ring is defined through their spectrums.

Remark 2.4. A Frobenius splitting of \mathfrak{X} compatible with \mathfrak{Y} induces a Frobenius splitting of \mathfrak{Y} (see e.g. [15, Remark 1.1.4 (ii)]).

Theorem 2.5 ([15] Lemma 1.1.14 and Exercise 1.1.E). *Let \mathfrak{X} be a separated scheme of finite type over \mathbb{k} with semiample line bundles $\mathcal{L}_1, \dots, \mathcal{L}_r$. If \mathfrak{X} admits a Frobenius splitting, then the multi-section ring*

$$\bigoplus_{n_1, \dots, n_r \geq 0} \Gamma(\mathfrak{X}, \mathcal{L}_1^{\otimes n_1} \otimes \dots \otimes \mathcal{L}_r^{\otimes n_r})$$

admits a Frobenius splitting ϕ . Moreover, a closed subscheme $\mathfrak{Y} \subset \mathfrak{X} = \text{Proj } R$ admits a compatible Frobenius splitting if and only if the homogeneous ideal $I_{\mathfrak{Y}} \subset R$ that defines \mathfrak{Y} satisfies $\phi(I_{\mathfrak{Y}}) \subset I_{\mathfrak{Y}}$, i.e. the pair $(R, R/I_{\mathfrak{Y}})$ admits a compatible Frobenius splitting ϕ . \square

Definition 2.6 (Canonical splitting). Let \mathfrak{X} be a separated scheme over \mathbb{k} equipped with a B -action. A Frobenius splitting ϕ is said to be B -canonical if it is H -fixed, and each $i \in \mathbf{I}$ yields

$$\rho_{\alpha_i}(z^p)\phi(\rho_{\alpha_i}(-z)f) = \sum_{j=0}^{p-1} \frac{z^j}{j!} \phi_{i,j}(f) \quad z \in \mathbb{k}, \quad (2.1)$$

where $\phi_{i,j} \in \text{Hom}_{\mathcal{O}_{\mathfrak{X}}}(\text{Fr}_* \mathcal{O}_{\mathfrak{X}}, \mathcal{O}_{\mathfrak{X}})$. We similarly define the notion of B^- -canonical splitting (resp. \mathbf{I} -canonical splitting and \mathbf{I}^- -canonical splitting) by using $\{\rho_{-\alpha_i}\}_{i \in \mathbf{I}}$ (resp. $\{\rho_{\alpha_i}\}_{i \in \mathbf{I}_{\text{af}}}$ and $\{\rho_{-\alpha_i}\}_{i \in \mathbf{I}_{\text{af}}}$) instead. Canonical splittings of a commutative ring S over \mathbb{k} is defined through its spectrum.

Proposition 2.7 (cf. [15] Proposition 4.1.8). *Let $S = \bigoplus_{m \geq 0} S_m$ be a graded ring with $S_0 = \mathbb{k}$ such that*

- S is equipped with a degree preserving \mathbf{I} -action;
- Each S_m is a graded \mathbb{k} -vector space compatible with the multiplication;

- We have an \mathbf{I} -canonical Frobenius splitting $\phi : S^{(1)} \rightarrow S$.

Then, the induced map

$$\phi^\vee : S_m^\vee \longrightarrow S_{pm}^\vee \quad m \in \mathbb{Z}_{\geq 0}$$

satisfies

$$\phi^\vee(E_i^{(n)} \mathbf{v}) = E_i^{(pn)} \phi^\vee(\mathbf{v}) \quad \forall i \in \mathbf{I}_{\text{af}}, n \in \mathbb{Z}_{\geq 0}, \mathbf{v} \in S_m^\vee.$$

Similar results hold for the \mathbf{I}^- - and B^\pm -actions.

Remark 2.8. In the opinion of the author, a merit of Proposition 2.7 over [15, Proposition 4.1.8] is that it becomes obvious that a projective variety \mathfrak{X} with a B -action has at most one B -canonical splitting whenever the space of global sections of all ample line bundles are (or can be made) B -cocyclic compatible with multiplications (cf. [15, Theorem 4.1.15] and Corollary 2.35).

Proof of Proposition 2.7. The condition that S_m is a graded vector space implies $S_m \xrightarrow{\cong} (S_m^\vee)^\vee$ for each $m \in \mathbb{Z}_{\geq 0}$. By [15, Proposition 4.1.8], each $\mathbf{w} \in S_{pm} \subset S^{(1)}$ satisfies $\phi(E_i^{(pn)} \mathbf{w}) = E_i^{(n)} \phi(\mathbf{w})$ for $i \in \mathbf{I}_{\text{af}}$ and $n \geq 0$. Using the natural non-degenerate invariant pairing $\langle \bullet, \bullet \rangle$ between S_m^\vee and S_m , we compute the most LHS of

$$\langle \mathbf{v}, \phi(E_i^{(p)} \mathbf{w}) \rangle = \langle \mathbf{v}, E_i \phi(\mathbf{w}) \rangle = - \langle \phi^\vee(E_i \mathbf{v}), \mathbf{w} \rangle$$

by using the invariance under the corresponding unipotent action as

$$\begin{aligned} \langle \mathbf{v}, \phi(E_i^{(p)} \mathbf{w}) \rangle &= - \sum_{k_1=1}^p \langle E^{(k_1)} \phi^\vee(\mathbf{v}), E_i^{(p-k_1)} \mathbf{w} \rangle \\ &\dots = \sum_{m=1}^p \sum_{k_\bullet > 0, k_1+k_2+\dots+k_m=p} (-1)^m \langle E_i^{(k_1)} E_i^{(k_2)} \dots \phi^\vee(\mathbf{v}), \mathbf{w} \rangle \\ &= - \langle E^{(p)} \phi^\vee(\mathbf{v}), \mathbf{w} \rangle \end{aligned}$$

since we have $E_i^{(k_1)} E_i^{(k_2)} \dots E_i^{(k_m)} \in p\mathbb{Z}E_i^{(p)}$ except for $k_1 = p, 0 = k_2 = \dots$. This implies the case $n = 1$.

Similarly, we have

$$\langle \mathbf{v}, \phi(E_i^{(pn)} \mathbf{w}) \rangle = \sum_{m=1}^n \sum_{k_\bullet > 0, k_1+k_2+\dots+k_m=n} (-1)^m \langle E_i^{(pk_1)} E_i^{(pk_2)} \dots \phi^\vee(\mathbf{v}), \mathbf{w} \rangle.$$

Compared with

$$\langle \mathbf{v}, E_i^{(n)} \phi(\mathbf{w}) \rangle = \sum_{m=1}^n \sum_{k_\bullet > 0, k_1+k_2+\dots+k_m=n} (-1)^m \langle \phi^\vee(E_i^{(k_1)} E_i^{(k_2)} \dots \mathbf{v}), \mathbf{w} \rangle$$

using induction on n , we conclude the result. \square

2.2 Representations of affine Lie algebras over \mathbb{Z}

In this section, we systematically use the global basis theory [41, 43, 44, 45, 65, 35] by specializing the quantum parameter q to 1. Therefore, we might refer these references without an explicit declaration that we specialize q .

We consider the Kostant-Lusztig \mathbb{Z} -form $U_{\mathbb{Z}}^+$ (resp. $U_{\mathbb{Z}}^-$) of $U([\mathfrak{J}, \mathfrak{J}])$ (resp. $U([\mathfrak{J}^-, \mathfrak{J}^-])$) obtained as the specialization $q = 1$ of the $\mathbb{Z}[q, q^{-1}]$ -integral form of the quantized enveloping algebras [66, §23.2].

Remark 2.9. We remark that $U_{\mathbb{Z}}^{\pm}$ are the same integral forms dealt in [31], and also coincide with the integral forms obtained through the Drinfeld presentation ([4, §2] and [73, Lemma 2.5]).

Note that $U_{\mathbb{Z}}^{\pm}$ are equipped with the \mathbb{Z} -bases $\mathbf{B}(\mp\infty)$ obtained by the specialization $q = 1$ of the lower global basis [41] (see also [66, §25]). In view of [65, 43], we have an idempotent \mathbb{Z} -integral form

$$\begin{aligned} \dot{U}_{\mathbb{Z}} &= \bigoplus_{\Lambda \in P^{\text{af}}} U_{\mathbb{Z}}^- U_{\mathbb{Z}}^+ a_{\Lambda} && \text{such that} \\ a_{\Lambda} a_{\Gamma} &= \delta_{\Lambda, \Gamma} a_{\Lambda} && \Lambda, \Gamma \in P^{\text{af}} \text{ and,} \\ E_i^{(m)} a_{\Lambda} &= a_{\Lambda + m\alpha_i} E_i^{(m)}, \quad F_i^{(m)} a_{\Lambda} = a_{\Lambda - m\alpha_i} F_i^{(m)} && i \in \mathbf{I}_{\text{af}}, m \in \mathbb{Z}_{\geq 0}. \end{aligned}$$

We set $\dot{U}_{\mathbb{Z}}^{\geq 0} \subset \dot{U}_{\mathbb{Z}}$ to be the subalgebra generated by $\{F_i^{(m)}\}_{i \in \mathbf{I}, m \in \mathbb{Z}_{\geq 0}}$, $\{a_{\Lambda}\}_{\Lambda \in P^{\text{af}}}$, and $U_{\mathbb{Z}}^+$.

If a $\dot{U}_{\mathbb{Z}}$ -module M admits a decomposition

$$M = \bigoplus_{\Lambda \in P^{\text{af}}} a_{\Lambda} M,$$

then we call this the P^{af} -weight decomposition. If $\Lambda \in P^{\text{af}}$ satisfies $a_{\Lambda} M \neq 0$, then we call Λ a P^{af} -weight of M . In case M is defined over a field \mathbb{k} , we define the P^{af} -character of M as

$$\text{gch } M := \sum_{\Lambda \in P^{\text{af}}} e^{\Lambda} \dim_{\mathbb{k}} a_{\Lambda} M$$

whenever the RHS makes sense. For each $n \in \mathbb{Z}$, we set

$$M_n := \sum_{\Lambda \in P^{\text{af}}, \langle d, \Lambda \rangle = n} a_{\Lambda} M \subset M$$

and call it the d -degree n -part of M . Note that these are consistent with (1.3) through the identification $q = e^{\delta}$.

For each $\lambda \in P$, we set

$$a_{\lambda}^0 M := \sum_{\Lambda \in P^{\text{af}}, \lambda = \bar{\Lambda}} a_{\Lambda} M. \quad (2.2)$$

We call the decomposition

$$M = \bigoplus_{\lambda \in P} a_{\lambda}^0 M,$$

the P -weight decomposition. We call a non-zero element of $a_\Lambda M$ (resp. $a_\lambda^0 M$) a P^{af} -weight vector of M (resp. a P -weight vector of M). We also call $\lambda \in P$ with $a_\lambda^0 M \neq \{0\}$ a P -weight of M .

Similarly, we have the Kostant-Lusztig \mathbb{Z} -form $U_{\mathbb{Z}}^{0,+}$ (resp. $U_{\mathbb{Z}}^{0,-}$) of $U(\mathfrak{n})$ (resp. $U(\mathfrak{n}^-)$). We have $U_{\mathbb{Z}}^{0,+} \subset U_{\mathbb{Z}}^+$ and $U_{\mathbb{Z}}^{0,-} \subset U_{\mathbb{Z}}^-$. In view of the characterization of global bases ([41]), we find that $\mathbf{B}^0(\mp\infty) := \mathbf{B}(\mp\infty) \cap U_{\mathbb{Z}}^{0,\pm}$ define \mathbb{Z} -bases of $U^{0,\pm}$.

We set $\dot{U}_{\mathbb{Z}}^0 \subset \dot{U}_{\mathbb{Z}}$ to be the subalgebra of $\dot{U}_{\mathbb{Z}}$ generated by $\{E_i^{(m)}, F_i^{(m)}\}_{i \in \mathbf{I}, m \in \mathbb{Z}_{\geq 0}}$, $\{a_\Lambda\}_{\Lambda \in P^{\text{af}}}$. For a field \mathbb{k} , a $\dot{U}_{\mathbb{k}}^{\geq 0}$ -module M with a P^{af} -weight decomposition is said to be $\dot{U}_{\mathbb{k}}^0$ -integrable if its $\{E_i^{(m)}, F_i^{(m)}\}_{m \geq 0}$ -action induces a $SL(2, i)_{\mathbb{k}}$ -action whose $(SL(2, i) \cap H)_{\mathbb{k}}$ -eigenvalues are given by the P -weights for each $i \in \mathbf{I}$.

Note that if a $U(\tilde{\mathfrak{g}}_{\mathbb{C}})$ -module V over \mathbb{C} carries a cyclic $\tilde{\mathfrak{h}}_{\mathbb{C}}$ -weight vector whose weight belongs to P^{af} and each of its $\tilde{\mathfrak{h}}_{\mathbb{C}}$ -weight space is finite-dimensional, then we have a $\dot{U}_{\mathbb{Z}}$ -lattice $V_{\mathbb{Z}}$ inside V . In such a case, the module $V_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{k}$ admits P^{af} - or P -weight decompositions.

We have the Chevalley involution of $\dot{U}_{\mathbb{Z}}$ defined as:

$$\theta(E_i^{(m)}) = F_i^{(m)}, \theta(F_i^{(m)}) = E_i^{(m)}, \quad \text{and} \quad \theta(a_\Lambda) = a_{-\Lambda} \quad i \in \mathbf{I}_{\text{af}}, m \in \mathbb{Z}_{\geq 0}, \Lambda \in P^{\text{af}}.$$

Definition 2.10 ([45] Definition 2.4 and §2.8). A $U(\tilde{\mathfrak{g}}_{\mathbb{C}})$ -module V over \mathbb{C} with a cyclic $\tilde{\mathfrak{h}}_{\mathbb{C}}$ -weight vector \mathbf{v} is said to be compatible with the negative global basis if we have

$$U_{\mathbb{Z}}^- \mathbf{v} = \bigoplus_{b \in \mathbf{B}(\infty)} \mathbb{Z} b \mathbf{v} \subset V.$$

If (V, \mathbf{v}) is compatible with the negative global basis, then we set

$$\mathbf{B}^-(V) = \mathbf{B}^-(V, \mathbf{v}) := \{b \mathbf{v} \mid b \in \mathbf{B}(\infty) \text{ s.t. } b \mathbf{v} \neq 0\} \subset V$$

and refer them as the negative global basis of V .

Compatibility with the positive global basis of V and the positive global basis $\mathbf{B}^+(V) = \mathbf{B}^+(V, \mathbf{v})$ of V is defined similarly.

Theorem 2.11 (Kashiwara [41] Theorem 5). *We have:*

1. For each $\Lambda \in P_+^{\text{af}}$, the $\tilde{\mathfrak{g}}_{\mathbb{C}}$ -module $L(\Lambda)_{\mathbb{C}}$ is compatible with the negative global basis;
2. For each $\lambda \in P_+$, we have

$$V(\lambda)_{\mathbb{C}} = \bigoplus_{b \in \mathbf{B}^0(\infty)} \mathbb{C} b \mathbf{v}_\lambda^0.$$

We set $\mathbf{B}(\Lambda) := \mathbf{B}^-(L(\Lambda)_{\mathbb{C}}, \mathbf{v}_\Lambda)$ for each $\Lambda \in P_+^{\text{af}}$.

For each $\Lambda \in P_+^{\text{af}}$ and $\lambda \in P_+$, we set

$$L(\Lambda)_{\mathbb{Z}} := U_{\mathbb{Z}}^- \mathbf{v}_\Lambda \subset L(\Lambda)_{\mathbb{C}} \quad \text{and} \quad V(\lambda)_{\mathbb{Z}} := (U_{\mathbb{Z}}^{0,-}) \mathbf{v}_\lambda^0 \subset V(\lambda)_{\mathbb{C}}.$$

Here $V(\lambda)_{\mathbb{Z}}$ acquires the action of $\dot{U}_{\mathbb{Z}}^0$ thanks to the splitting $P \hookrightarrow P^{\text{af}}$.

Corollary 2.12. *We have:*

1. For each $\Lambda, \Gamma \in P_+^{\text{af}}$, we have a natural inclusion $L(\Lambda + \Gamma)_{\mathbb{Z}} \hookrightarrow L(\Lambda)_{\mathbb{Z}} \otimes_{\mathbb{Z}} L(\Gamma)_{\mathbb{Z}}$ of $\dot{U}_{\mathbb{Z}}$ -modules, that is a direct summand as \mathbb{Z} -modules;
2. For each $\lambda, \mu \in P_+$, we have a natural inclusion $V(\lambda + \mu)_{\mathbb{Z}} \hookrightarrow V(\lambda)_{\mathbb{Z}} \otimes_{\mathbb{Z}} V(\mu)_{\mathbb{Z}}$ of $\dot{U}_{\mathbb{Z}}^0$ -modules, that is a direct summand as \mathbb{Z} -modules.

Proof. Since the two cases are completely parallel, we only prove the first case. The $\tilde{\mathfrak{g}}$ -module $L(\Lambda)_{\mathbb{C}} \otimes_{\mathbb{C}} L(\Gamma)_{\mathbb{C}}$ decomposes into the direct sum of integrable highest weight modules ([40, Proposition 9.10]), with a direct summand $L(\Lambda + \Gamma)_{\mathbb{C}}$. In view of [41, Theorem 3], it gives rise to the $\mathbb{Z}[\mathfrak{q}]$ -lattice of the quantized version of $L(\Lambda) \otimes L(\Gamma)$ compatible with those of $L(\Lambda + \Gamma)$ via the natural embedding. By setting $\mathfrak{q} = 1$, we obtain a direct sum decomposition of $L(\Lambda)_{\mathbb{Z}} \otimes_{\mathbb{Z}} L(\Gamma)_{\mathbb{Z}}$ as \mathbb{Z} -modules with its direct summand $L(\Lambda + \Gamma)_{\mathbb{Z}}$. \square

Theorem 2.13 (Kashiwara [43] Proposition 8.2.2). *For each $\lambda \in P_+$, the $\tilde{\mathfrak{g}}_{\mathbb{C}}$ -module $\mathbb{X}(\lambda)_{\mathbb{C}}$ is compatible with the negative/positive global basis (for every extremal weight vector).* \square

For each $\lambda \in P_+$, we set

$$\mathbb{X}(\lambda)_{\mathbb{Z}} := \dot{U}_{\mathbb{Z}} \mathbf{v}_{\lambda} \subset \mathbb{X}(\lambda)_{\mathbb{C}}.$$

Theorem 2.14 (Kashiwara [45]). *Let $\lambda \in P_+$. There exists a \mathbb{Z} -basis $\mathbf{B}(\mathbb{X}(\lambda))$ of $\mathbb{X}(\lambda)_{\mathbb{Z}}$ that contains the negative/positive global basis of $\mathbb{X}(\lambda)_{\mathbb{Z}}$ constructed from every extremal weight vector of $\mathbb{X}(\lambda)_{\mathbb{C}}$.*

Proof. We set $\mathbf{B}(\mathbb{X}(\lambda))$ to be the specialization of the global basis of a quantum loop algebra module [43, Proposition 8.2.2]. Then, it is compatible with the global basis generated from an extremal weight vectors by [45, Theorem 3.3]. \square

For each $\lambda \in P_+$ and $w \in W_{\text{af}}$, we define

$$\mathbb{W}_w(\lambda)_{\mathbb{Z}} := U_{\mathbb{Z}}^+ \mathbf{v}_{w\lambda} \subset \mathbb{X}(\lambda)_{\mathbb{C}} \quad \text{and} \quad \mathbb{W}_w^-(\lambda)_{\mathbb{Z}} := U_{\mathbb{Z}}^- \mathbf{v}_{w\lambda} \subset \mathbb{X}(\lambda)_{\mathbb{C}}.$$

We set $\mathbb{W}(\lambda)_{\mathbb{Z}} := \mathbb{W}_{w_0}(\lambda)_{\mathbb{Z}}$ and $\mathbb{W}^-(\lambda)_{\mathbb{Z}} := \mathbb{W}_e^-(\lambda)_{\mathbb{Z}}$.

Lemma 2.15 (Naito-Sagaki). *For each $\lambda \in P_+$ and $w, v \in W_{\text{af}}$, we have $\mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}} \subset \mathbb{W}_{vw_0}(\lambda)_{\mathbb{Z}}$ if $w \leq_{\infty} v$. If we have $\lambda \in P_{++}$ in addition, then we have $\mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}} \subset \mathbb{W}_{vw_0}(\lambda)_{\mathbb{Z}}$ if and only if $w \leq_{\infty} v$.*

Proof. Apply the inclusion relation of the (labels of the) global basis in [72, Corollary 5.2.5] (see also [45, §2.8]). \square

Corollary 2.16. *For each $\lambda \in P_+$, $w \in W_{\text{af}}$, and $i \in \mathbf{I}_{\text{af}}$, we have $\mathbb{W}_{s_i w w_0}(\lambda)_{\mathbb{K}} \subset \mathbb{W}_{w w_0}(\lambda)_{\mathbb{K}}$ if $s_i w \leq_{\infty} w$. In this case, $\mathbb{W}_{w w_0}(\lambda)_{\mathbb{K}}$ inherits an action of $SL(2, i)_{\mathbb{K}}$ from $\mathbb{X}(\lambda)_{\mathbb{K}}$.*

Proof. The first part of the assertion is the special case of Lemma 2.15. Given this, it remains to notice that a lift of $s_i \in W_{\text{af}}$ sends $\mathbf{v}_{w w_0 \lambda}$ to $\pm \mathbf{v}_{s_i w w_0 \lambda}$, and hence the Bruhat decomposition of $\mathbf{I}(i)_{\mathbb{K}}$ (into two $\mathbf{I}_{\mathbb{K}}$ -double cosets) implies that $\mathbb{W}_{w w_0}(\lambda)_{\mathbb{K}}$ is stable under $\mathbf{I}(i)_{\mathbb{K}}$. \square

Lemma 2.17. *For each $\lambda \in P_+$ and $w \in W_{\text{af}}$, it holds:*

1. Each $\beta \in Q^\vee$ defines a $\dot{U}_{\mathbb{Z}}$ -module automorphism τ_β on $\mathbb{X}(\lambda)_{\mathbb{Z}}$ determined by $\tau_\beta(\mathbf{v}_\lambda) := \mathbf{v}_{t_\beta \lambda}$. Moreover, we have $\tau_\beta \mathbf{B}(\mathbb{X}(\lambda)) = \mathbf{B}(\mathbb{X}(\lambda))$;
2. We have $\theta^*(\mathbb{X}(\lambda)_{\mathbb{Z}}) \cong \mathbb{X}(-w_0 \lambda)_{\mathbb{Z}}$ as $\dot{U}_{\mathbb{Z}}$ -modules. Moreover, we have $\theta^* \mathbf{B}(\mathbb{X}(\lambda)) = \mathbf{B}(\mathbb{X}(-w_0 \lambda))$;
3. We have $\mathbb{W}_w(\lambda)_{\mathbb{Z}} = \mathbb{W}_w(\lambda)_{\mathbb{C}} \cap \mathbb{X}(\lambda)_{\mathbb{Z}}$;
4. We have a $U_{\mathbb{Z}}^-$ -cyclic vector of $\theta^*(\mathbb{W}_w(\lambda)_{\mathbb{Z}})$ with weight $-w\lambda = ww_0(-w_0\lambda)$. In particular, we have

$$\theta^*(\mathbb{W}_w(\lambda)_{\mathbb{Z}}) \cong \mathbb{W}_{ww_0}^-(-w_0\lambda)_{\mathbb{Z}} \quad \text{and} \quad \theta^*(\mathbb{W}_w^-(\lambda)_{\mathbb{Z}}) \cong \mathbb{W}_{ww_0}(-w_0\lambda)_{\mathbb{Z}}.$$

Proof. We borrow the setting of [43, §8.1 and §8.2].

We prove the first assertion. Since \mathbf{v}_λ and $\mathbf{v}_{t_\beta \lambda}$ obeys the same relation, τ_β defines an automorphism as $\tilde{\mathfrak{g}}_{\mathbb{C}}$ -modules. The latter assertion follows from Theorem 2.14.

We prove the second assertion. The defining equation of $\theta^*(\mathbf{v}_\lambda)$ is the same as the cyclic vector $\mathbf{v}_{-w_0\lambda} \in \mathbb{X}(-w_0\lambda)_{\mathbb{C}}$ as $\tilde{\mathfrak{g}}_{\mathbb{C}}$ -modules. This yields a $\tilde{\mathfrak{g}}_{\mathbb{C}}$ -module isomorphism $\eta : \theta^*(\mathbb{X}(\lambda)_{\mathbb{C}}) \rightarrow \mathbb{X}(-w_0\lambda)_{\mathbb{C}}$. Since θ exchanges $U_{\mathbb{Z}}^\pm$ and \mathbf{v}_λ is cyclic, we deduce that $\eta(\theta^*(\mathbb{X}(\lambda)_{\mathbb{Z}})) = \dot{U}_{\mathbb{Z}} \mathbf{v}_{-\lambda} \subset \mathbb{X}(-w_0\lambda)_{\mathbb{C}}$. By Theorem 2.14, we conclude $\theta^* \mathbf{B}(\mathbb{X}(\lambda)) = \mathbf{B}(\mathbb{X}(-w_0\lambda))$ as required.

We prove the third assertion. By Theorem 2.14, the \mathbb{Z} -basis of $\mathbb{W}(\lambda)_{\mathbb{Z}}$ is formed by the non-zero elements of $\mathbf{B}(-\infty)\mathbf{v}_{w_0\lambda}$ and forms a direct summand of $\mathbb{X}(\lambda)_{\mathbb{Z}}$ as \mathbb{Z} -modules. Hence, the case $w = w_0$ follows. For $w \in W$, we apply [43, Lemma 8.2.1] repeatedly to deduce the assertion from the $w = w_0$ case by using $\mathbf{B}(-\infty)\mathbf{v}_{w\lambda} \subset \mathbf{B}(-\infty)\mathbf{v}_{w_0\lambda}$. For $w = ut_\beta \in W_{\text{af}}$ with $u \in W, \beta \in Q^\vee$, we additionally apply $\tau_{w_0\beta}$ to conclude the assertion.

We prove the fourth assertion. The vector $\theta^*(\mathbf{v}_{w\lambda})$ is a $U_{\mathbb{Z}}^-$ -cyclic vector of $\theta^*(\mathbb{W}_w(\lambda)_{\mathbb{Z}})$, and its weight is

$$-w\lambda = ww_0(-w_0\lambda).$$

Hence, we conclude the assertion (using the fact that θ is an involution). \square

Theorem 2.18 ([44] Proposition 8.6 and [5] Corollary 4.15). *Let $\lambda \in P_+$. The unique (d -degree zero) $\dot{U}_{\mathbb{Z}}^{\geq 0}$ -module map*

$$\Psi_\lambda : \mathbb{W}(\lambda)_{\mathbb{Z}} \hookrightarrow \bigotimes_{i \in \mathbf{I}} \mathbb{W}(\varpi_i)_{\mathbb{Z}}^{\otimes \langle \alpha_i^\vee, \lambda \rangle}, \quad (2.3)$$

that sends \mathbf{v}_λ to the tensor product of $\{\mathbf{v}_{\varpi_i}\}_{i \in \mathbf{I}}$'s, is injective and defines a direct summand as \mathbb{Z} -modules.

Proof. We set $\mathbb{X}^\otimes := \bigotimes_{i \in \mathbf{I}} \mathbb{X}(\varpi_i)_{\mathbb{Z}}^{\otimes \langle \alpha_i^\vee, \lambda \rangle}$. The map Ψ_λ exists as $\bigotimes_{i \in \mathbf{I}} \mathbf{v}_{\varpi_i}^{\otimes \langle \alpha_i^\vee, \lambda \rangle}$ obeys the same defining condition as the extremal weight vector $\mathbf{v}_\lambda \in \mathbb{X}(\lambda)_{\mathbb{Z}}$, and the comultiplication of $\tilde{\mathfrak{g}}$ induces an algebra map $\Delta : \dot{U}_{\mathbb{Z}} \subset \dot{U}_{\mathbb{Z}} \otimes \dot{U}_{\mathbb{Z}}$ ([66, §23.1.5]).

The map Ψ_λ is injective by [5, Corollary 4.15 and Remark 4.17]. By [44, Theorem 8.5], the $\mathfrak{q} = 1$ specializations of the global bases yield the $\dot{U}_{\mathbb{Z}}$ -spans of the extremal weight vectors in $\mathbb{X}(\lambda)_{\mathbb{Z}}$ and \mathbb{X}^\otimes up to the action of some rings of (partially symmetric) polynomials with integer coefficients, respectively. This

yields a splitting of Ψ_λ as \mathbb{Z} -modules, where the \mathbb{Z} -module structure of the RHS is coming from the crystal lattice of (the \mathfrak{q} -version of) \mathbb{X}^\otimes (as in [44, (8.8)]). We call this crystal lattice \mathbf{L}_1 .

We have another $\mathbb{Z}[\mathfrak{q}]$ -lattice inside (the \mathfrak{q} -version of) \mathbb{X}^\otimes obtained by the tensor product of the crystal lattices of (the \mathfrak{q} -versions of) $\mathbb{X}(\varpi_i)$'s. We call this $\mathbb{Z}[\mathfrak{q}]$ -lattice \mathbf{L}_2 .

By [44, Proposition 8.6], the global basis of \mathbf{L}_1 is written in terms of the (tensor product) global bases of \mathbf{L}_2 by an upper-unitriangular matrix C (valued in $\mathbb{Q}[\mathfrak{q}]$, with finitely many non-zero entry in each row). Hence, the corresponding bases in \mathbb{X}^\otimes are related by an upper-unitriangular matrix C obtained as the $\mathfrak{q} = 1$ specialization of C (valued in \mathbb{Q}). In view of the fact that Δ is an algebra morphism (and [44, Theorem 8.5]), we deduce $\mathbf{L}_1 \subset \mathbf{L}_2$. Hence, we have $\mathbb{Z} \otimes_{\mathbb{Z}[\mathfrak{q}]} \mathbf{L}_1 \subset \mathbb{Z} \otimes_{\mathbb{Z}[\mathfrak{q}]} \mathbf{L}_2 \subset \mathbb{X}^\otimes$. In particular, the matrix C must be valued in \mathbb{Z} . This implies $\mathbb{Z} \otimes_{\mathbb{Z}[\mathfrak{q}]} \mathbf{L}_1 = \mathbb{Z} \otimes_{\mathbb{Z}[\mathfrak{q}]} \mathbf{L}_2$.

Hence, the \mathbb{Z} -module splitting in the second paragraph is indeed what we wanted. \square

2.3 Projectivity of the module $\mathbb{W}(\lambda)_\mathbb{k}$

Let \mathbb{k} be a field. We equip $\mathring{U}_\mathbb{k}^+ := \bigoplus_{\lambda \in P} U_\mathbb{k}^+ a_\lambda^0$ with the structure of an algebra by setting $a_\lambda^0 a_\mu^0 = \delta_{\lambda, \mu} a_\lambda^0$ ($\lambda, \mu \in P$),

$$E_0^{(m)} a_\lambda^0 = a_{\lambda - m\vartheta}^0 E_0^{(m)}, \quad E_i^{(m)} a_\lambda^0 = a_{\lambda + m\alpha_i}^0 E_i^{(m)} \quad (i \in \mathbf{I}, m \in \mathbb{Z}_{\geq 0}, \lambda \in P),$$

and requires

$$(\xi a_\lambda^0)(\xi' a_\mu^0) = (\xi\xi') a_\mu^0 \quad \text{when} \quad a_\lambda^0 \xi' a_\mu^0 = \xi' a_\mu^0 \quad (\lambda, \mu \in P, \xi, \xi' \in U_\mathbb{k}^+).$$

Similarly, we define

$$\mathring{U}_\mathbb{k}^{\geq 0} := \bigoplus_{\lambda \in P} U_\mathbb{k}^{0, -} U_\mathbb{k}^+ a_\lambda^0 \quad \text{and} \quad \mathring{U}_\mathbb{k}^0 := \bigoplus_{\lambda \in P} U_\mathbb{k}^{0, -} U_\mathbb{k}^{0, +} a_\lambda^0$$

and regard them as subspaces of the completions of $\mathring{U}_\mathbb{k}^{\geq 0}$ and $\mathring{U}_\mathbb{k}^0$ with respect to the idempotents by setting

$$\sum_{\lambda = \bar{\Lambda}} a_\lambda = a_\Lambda^0 \quad (\lambda \in P \subset P^{\text{af}}).$$

The eigendecomposition with respect to $d \in \tilde{\mathfrak{h}}$ now becomes an external gradings of algebras $\mathring{U}_\mathbb{k}^+$ and $\mathring{U}_\mathbb{k}^{\geq 0}$. We have algebra inclusions

$$\mathring{U}_\mathbb{k}^+ \subset \mathring{U}_\mathbb{k}^{\geq 0} \supset \mathring{U}_\mathbb{k}^0.$$

Note that we have a surjective algebra map $\mathring{U}_\mathbb{Z}^{\geq 0} \rightarrow \mathring{U}_\mathbb{Z}^0$ since $a_\Lambda \mathring{U}_\mathbb{Z}^{\geq 0} a_\Gamma \neq 0$ implies $\langle d, \Lambda \rangle \geq \langle d, \Gamma \rangle$ and $a_\Lambda \mathring{U}_\mathbb{Z}^0 a_\Gamma \neq 0$ implies $\langle d, \Lambda \rangle = \langle d, \Gamma \rangle$ for $\Lambda, \Gamma \in P^{\text{af}}$. This induces a surjective algebra map $\mathring{U}_\mathbb{Z}^{\geq 0} \rightarrow \mathring{U}_\mathbb{Z}^0$, that specializes to $\mathring{U}_\mathbb{k}^{\geq 0} \rightarrow \mathring{U}_\mathbb{k}^0$.

We sometimes regard $\mathbb{W}(\lambda)_\mathbb{k}$ ($\lambda \in P_+$) as a graded $\mathring{U}_\mathbb{k}^+$ -module or a graded $\mathring{U}_\mathbb{k}^{\geq 0}$ -module by means of (2.2), whose gradings are given by the d -degrees.

Proposition 2.19. *Let \mathbb{k} be a field and let $\lambda \in P_+$. The module $\mathbb{W}(\lambda)_{\mathbb{k}}$ is the projective cover of $V(\lambda)_{\mathbb{k}}$ in the category of $\dot{U}_{\mathbb{k}}^{\geq 0}$ -modules that are $\dot{U}_{\mathbb{k}}^0$ -integrable and whose P -weights are contained in $\text{Conv } W\lambda \subset P \otimes_{\mathbb{Z}} \mathbb{R}$, where Conv denote the \mathbb{R} -convex hull.*

Lemma 2.20. *For each $\lambda \in P_+$, the module $\mathbb{W}(\lambda)_{\mathbb{k}}$ is free over a polynomial ring and we have a finite-dimensional quotient $W(\lambda)_{\mathbb{k}}$ with $\dim_{\mathbb{k}} a_{\lambda}^0 W(\lambda)_{\mathbb{k}} = 1$.*

Remark 2.21. The modules $\mathbb{W}(\lambda)_{\mathbb{k}}$ and $W(\lambda)_{\mathbb{k}}$ ($\lambda \in P_+$) are analogues of global and local Weyl modules in [19, 62, 18] (cf. Theorem 1.2). By construction, their characters are the same as these defined over \mathbb{C} (see [46, §1] for more discussion).

Proof of Lemma 2.20. Note that the endomorphism τ_{β} in Lemma 2.17 induces an endomorphism of $\mathbf{B}(\mathbb{X}(\varpi_i))$ for each $\beta \in Q$ and $i \in \mathbf{I}$, and hence we know the existence of the injective endomorphisms of $\mathbb{W}(\lambda)_{\mathbb{k}}$ that forms a polynomial algebra whose graded dimension is equal to

$$\text{gdim } \text{End}_{\mathfrak{g}[z]_{\mathbb{C}}} \mathbb{W}(\lambda)_{\mathbb{C}} = \prod_{i \in \mathbf{I}} \prod_{j=1}^{\langle \alpha_i^{\vee}, \lambda \rangle} \frac{1}{1 - q^j},$$

that can be read-off from [5, Proposition 4.13] (or [46, Theorem 1.4]), through Theorem 2.18. By [5, 18, 46], it exhausts the P -weight λ -part of $\mathbb{W}(\lambda)_{\mathbb{k}}$, that generates $\mathbb{W}(\lambda)_{\mathbb{k}}$ as a $\dot{U}_{\mathbb{k}}^{\geq 0}$ -module. Therefore, we conclude the result. \square

The rest of this subsection is devoted to the proof of Proposition 2.19.

We consider the Demazure functor \mathcal{D}_w for $w \in W_{\text{af}}$ with respect to $\dot{U}_{\mathbb{k}}^+$ (cf. [39, 46, 20]). In view of [45, §2.8], the character part of the calculation in [46, Theorem 4.13] carries over to our setting, and hence we have

$$\mathbf{L} \bullet \mathcal{D}_{t_{\beta}}(\mathbb{W}(\lambda)_{\mathbb{k}}) = \mathcal{D}_{t_{\beta}}(\mathbb{W}(\lambda)_{\mathbb{k}}) \cong \mathbb{W}(\lambda)_{\mathbb{k}} \otimes_{\mathbb{k}} \mathbb{k}_{-\langle \beta, w_0 \lambda \rangle \delta} \quad \beta \in Q_{<}^{\vee}.$$

From this, we also derive that

$$\mathbf{L} \bullet \mathcal{D}_{t_{\beta}}(W(\lambda)_{\mathbb{k}}) = \mathcal{D}_{t_{\beta}}(W(\lambda)_{\mathbb{k}}) \cong W(\lambda)_{\mathbb{k}} \otimes_{\mathbb{k}} \mathbb{k}_{-\langle \beta, w_0 \lambda \rangle \delta} \quad \beta \in Q_{<}^{\vee}$$

using the Koszul resolution as in [20, §5.1.4] by Lemma 2.20.

The d -gradings of $\dot{U}_{\mathbb{k}}^+$, $\mathbb{W}(\lambda)$, and $W(\mu)_{\mathbb{k}}$ are concentrated in non-negative degrees. It follows that the d -grading of $\text{Ext}_{\dot{U}_{\mathbb{k}}^+}^{\bullet}(\mathbb{W}(\lambda), W(\mu)_{\mathbb{k}}^*)$ is bounded from the above. Moreover, we have

$$\text{Ext}_{\dot{U}_{\mathbb{k}}^+}^{\bullet}(\mathcal{D}_{t_{\beta} w_0}(\mathbb{W}(\lambda)_{\mathbb{k}}), W(\mu)_{\mathbb{k}}^*) \cong \text{Ext}_{\dot{U}_{\mathbb{k}}^+}^{\bullet}(\mathbb{W}(\lambda)_{\mathbb{k}}, \mathcal{D}_{t_{-w_0 \beta} w_0}(W(\mu)_{\mathbb{k}})^*)$$

for every $\lambda, \mu \in P_+$ and $\beta \in Q_{<}^{\vee}$ by (a repeated application of) [25, Proposition 5.7] (whose argument carries over to our setting). By varying β , we conclude that

$$\text{Ext}_{\dot{U}_{\mathbb{k}}^+}^{\bullet}(\mathbb{W}(\lambda)_{\mathbb{k}}, W(\mu)_{\mathbb{k}}^*) \equiv \{0\} \quad \lambda \neq -w_0 \mu \quad (2.4)$$

since $-\langle \beta, w_0 \lambda \rangle \neq -\langle -w_0 \beta, w_0 \mu \rangle = \langle \beta, \mu \rangle$ for some choice of β .

Consider the simple integrable $\dot{U}_{\mathbb{k}}^0$ -module quotient $L(\lambda)_{\mathbb{k}}$ of $V(\lambda)_{\mathbb{k}}$ for each $\lambda \in P_+$.

Lemma 2.22. *The set $\{L(\lambda)_{\mathbb{k}}\}_{\lambda \in P_+}$ is the complete collection of the isomorphism classes of the irreducible d -graded $\dot{U}_{\mathbb{k}}^0$ -integrable $\dot{U}_{\mathbb{k}}^{\geq 0}$ -modules (up to d -grading shifts).*

Proof. Since $V(\lambda)_{\mathbb{k}}$ surjects onto every irreducible $\dot{U}_{\mathbb{k}}^0$ -modules with highest weight λ ([1, §1.20]), it follows that $V(\lambda)_{\mathbb{k}}$ surjects onto every irreducible d -graded $\dot{U}_{\mathbb{k}}^{\geq 0}$ -module with highest weight λ . Since the P -weight λ -part of $V(\lambda)_{\mathbb{k}}$ is one-dimensional, we conclude the result. \square

We return to the proof of Proposition 2.19. We have

$$\mathrm{gch} W(\lambda)_{\mathbb{k}} \equiv \mathrm{gch} V(\lambda)_{\mathbb{k}} \equiv \mathrm{gch} L(\lambda)_{\mathbb{k}} \pmod{\sum_{\lambda > \mu \in P_+} \mathbb{Z}[q] \mathrm{gch} L(\mu)_{\mathbb{k}}}$$

by [62] (cf. [18]). Therefore, $W(\lambda)_{\mathbb{k}}$ admits a (d -graded) Jordan-Hölder series as a $\dot{U}_{\mathbb{k}}^{\geq 0}$ -module whose irreducible constituents are of the form $\{L(\mu)_{\mathbb{k}}\}_{\mu \leq \lambda}$ (up to d -grading shifts). It follows that

$$\mathrm{Ext}_{\dot{U}_{\mathbb{k}}^+}^{\bullet}(\mathbb{W}(\lambda)_{\mathbb{k}}, V(\mu)_{\mathbb{k}}) = \mathrm{Ext}_{\dot{U}_{\mathbb{k}}^+}^{\bullet}(\mathbb{W}(\lambda)_{\mathbb{k}}, L(\mu)_{\mathbb{k}}) \equiv \{0\} \quad \lambda > \mu \in P_+$$

by a repeated application of the short exact sequences to (2.4). Since the both of $\mathbb{W}(\lambda)_{\mathbb{k}}$ and $L(\lambda)_{\mathbb{k}}$ are $\dot{U}_{\mathbb{k}}^0$ -integrable, we find

$$\mathrm{Ext}_{\dot{U}_{\mathbb{k}}^{\geq 0}}^1(\mathbb{W}(\lambda)_{\mathbb{k}}, L(\mu)_{\mathbb{k}}) \equiv \{0\} \quad \lambda > \mu \in P_+.$$

From these, it suffices to prove

$$\mathrm{Ext}_{\dot{U}_{\mathbb{k}}^{\geq 0}}^1(\mathbb{W}(\lambda)_{\mathbb{k}}, L(\lambda)_{\mathbb{k}}) \equiv \{0\} \quad \lambda \in P_+ \quad (2.5)$$

in order to deduce the assertion.

In view of the fact $\lambda + \alpha_i$ ($i \in \mathbf{I}$) is not a P -weight of $\mathbb{W}(\lambda)_{\mathbb{k}}$, the Drinfeld presentation of $\dot{U}_{\mathbb{k}}^{\geq 0}$ ([5, §3]) forces that the space of $\dot{U}_{\mathbb{k}}^{\geq 0}$ -module endomorphisms of $\mathbb{W}(\lambda)_{\mathbb{k}}$ is generated by the images of the imaginary weight vectors $\tilde{P}_{i,m\delta}$ for $i \in \mathbf{I}$ and $m > 0$ (defined in [5, (3.7)]). In view of its descriptions ([5, Proposition 3.17] or [17, Lemma 4.5]), we find that $\tilde{P}_{i,m}$ acts on $\mathbf{v}_{\lambda} \in \mathbb{W}(\lambda)_{\mathbb{k}}$ by zero if $\langle \alpha_i^{\vee}, \lambda \rangle > m$. From this, we derive that the P -weight λ -part of $\mathbb{W}(\lambda)_{\mathbb{k}}$ is maximal possible as a cyclic module with a cyclic vector of P -weight λ in the category of $\dot{U}_{\mathbb{k}}^{\geq 0}$ -modules that is $\dot{U}_{\mathbb{k}}^0$ -integrable and whose P -weights are contained in $\mathrm{Conv} W\lambda \subset P \otimes_{\mathbb{Z}} \mathbb{R}$. Therefore, (2.5) vanishes and we conclude Proposition 2.19.

2.4 Frobenius splitting of $\mathbf{Q}_{G,J}$

Theorem 2.23. *For each $\Lambda \in P_+^{\mathrm{af}}$, we have a surjective map $L(\Lambda)_{\mathbb{C}} \rightarrow \mathbb{W}^-(\bar{\Lambda})_{\mathbb{C}}$ of $\mathfrak{g}[z^{-1}]_{\mathbb{C}}$ -modules. In addition, this map yields a surjection $L(\Lambda)_{\mathbb{Z}} \rightarrow \mathbb{W}^-(\bar{\Lambda})_{\mathbb{Z}}$ of $U_{\mathbb{Z}}^-$ -modules.*

Proof. By [50, Theorem A], the graded $\mathfrak{g}[z]_{\mathbb{C}}$ -module $\theta^*(L(\Lambda)_{\mathbb{C}})$ admits a filtration by the grading shifts of $\{\mathbb{W}(\mu)_{\mathbb{C}}\}_{\mu \in P_+}$. Since the (d -)degrees of $L(\Lambda)_{\mathbb{C}}$ is concentrated in $\mathbb{Z}_{\leq 0}$ and the degree 0-part of $L(\Lambda)_{\mathbb{C}}$ is $V(\bar{\Lambda})_{\mathbb{C}}$, the first quotient of $\theta^*(L(\Lambda)_{\mathbb{C}})$ in our filtration must be $\theta^*(\mathbb{W}^-(\bar{\Lambda})_{\mathbb{C}})$. Hence, we obtain the surjection $\eta : L(\Lambda)_{\mathbb{C}} \rightarrow \mathbb{W}^-(\bar{\Lambda})_{\mathbb{C}}$ of $\mathfrak{g}[z^{-1}]_{\mathbb{C}}$ -modules.

Since the both modules share the $U_{\mathbb{C}}^-$ -cyclic vector and compatible with the negative global basis, we conclude that the \mathbb{Z} -basis of $\mathbb{W}^-(\bar{\Lambda})_{\mathbb{Z}}$ is obtained as a \mathbb{Z} -basis of $L(\Lambda)_{\mathbb{Z}}$ that is not annihilated by η . Hence, we conclude that η induces a surjection $L(\Lambda)_{\mathbb{Z}} \rightarrow \mathbb{W}^-(\bar{\Lambda})_{\mathbb{Z}}$ of $U_{\mathbb{Z}}^-$ -modules. \square

Corollary 2.24. *Let \mathbb{k} be a field. For $\Lambda \in P_+^{\text{af}}$, we have a $U_{\mathbb{k}}^-$ -module generator set $\{\mathbf{u}_m\}_{m \in \mathbb{Z}_{\geq 0}}$ of $\ker(L(\Lambda)_{\mathbb{k}} \rightarrow \mathbb{W}^-(\bar{\Lambda})_{\mathbb{k}})$ that satisfies:*

- Each element \mathbf{u}_m satisfies $a_{\Lambda_m} \mathbf{u}_m = \mathbf{u}_m$ for some $\Lambda_m \in P_+^{\text{af}}$;
- For each $m \in \mathbb{Z}_{\geq 0}$, we have $\bar{\Lambda}_m \notin \text{Conv } W\bar{\Lambda} \subset P \otimes_{\mathbb{Z}} \mathbb{R}$.

Proof. Note that $L(\Lambda)_{\mathbb{k}}$ has at most countable rank over \mathbb{k} , which implies that the generator set is at most countable. As the both of $L(\Lambda)_{\mathbb{k}}$ and $\mathbb{W}^-(\bar{\Lambda})_{\mathbb{k}}$ admit P^{af} -weight decompositions, we deduce the first assertion. Since the both modules are $U_{\mathbb{k}}^0$ -integrable and $U_{\mathbb{k}}^-$ -cyclic, the second assertion follows by Proposition 2.19. \square

Proposition 2.25. *For each $\Lambda, \Gamma \in P_+^{\text{af}}$, we have the following commutative diagram of $U_{\mathbb{Z}}^-$ -modules:*

$$\begin{array}{ccc} L(\Lambda + \Gamma)_{\mathbb{Z}} & \hookrightarrow & L(\Lambda)_{\mathbb{Z}} \otimes_{\mathbb{Z}} L(\Gamma)_{\mathbb{Z}} \\ \downarrow & & \downarrow \\ \mathbb{W}^-(\overline{\Lambda + \Gamma})_{\mathbb{Z}} & \xrightarrow{\mathfrak{m}} & \mathbb{W}^-(\bar{\Lambda})_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{W}^-(\bar{\Gamma})_{\mathbb{Z}}. \end{array}$$

Moreover, all the maps define direct summands as \mathbb{Z} -modules.

Proof. The injectivity of the top horizontal arrow and the fact that it defines a direct summand as \mathbb{Z} -modules is Corollary 2.12.

The surjectivity of the vertical arrows are Theorem 2.23. Since they are obtained by annihilating parts of \mathbb{Z} -bases, these maps define direct summands as \mathbb{Z} -modules.

Since all the modules are generated by the cyclic vectors $\mathbf{v}_{\Lambda + \Gamma}$ or $\mathbf{v}_{\Lambda} \otimes \mathbf{v}_{\Gamma}$ as $\mathfrak{g}[z^{-1}]_{\mathbb{C}}$ -modules or $\mathfrak{g}[z^{-1}]_{\mathbb{C}}^{\oplus 2}$ -modules, Theorem 1.3 (twisted by θ) implies the injectivity of \mathfrak{m} after extending the scalar to \mathbb{C} . Hence, we deduce

$$\mathfrak{m}(\mathbb{W}^-(\overline{\Lambda + \Gamma})_{\mathbb{Z}}) \subset \mathbb{W}^-(\bar{\Lambda})_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{W}^-(\bar{\Gamma})_{\mathbb{Z}}. \quad (2.6)$$

Therefore, it suffices to prove that (2.6) has torsion-free cokernel (as a \mathbb{Z} -module) to complete the proof. By a repeated use of (2.6), we arrive the setting of Theorem 2.18 in view of Theorem 1.3. Thus, the map \mathfrak{m} defines a direct summand of $\mathbb{W}^-(\bar{\Lambda})_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{W}^-(\bar{\Gamma})_{\mathbb{Z}}$ as \mathbb{Z} -modules. \square

Corollary 2.26. *For each $\lambda, \mu \in P_+$ and $w \in W$, we have the following commutative diagram of $U_{\mathbb{Z}}^+$ -modules:*

$$\begin{array}{ccc} \mathbb{X}(\lambda + \mu)_{\mathbb{Z}} & \hookrightarrow & \mathbb{X}(\lambda)_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{X}(\mu)_{\mathbb{Z}} \\ \uparrow & & \uparrow \\ \mathbb{W}_w(\lambda + \mu)_{\mathbb{Z}} & \hookrightarrow & \mathbb{W}_w(\lambda)_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{W}_w(\mu)_{\mathbb{Z}} \end{array}$$

Moreover, the vertical inclusions are compatible with positive global basis, and the horizontal inclusions define direct summands as \mathbb{Z} -modules. In addition, all the inclusions commute with the automorphism τ_β ($\beta \in Q^\vee$) of $\mathbb{X}(\lambda)_\mathbb{Z}, \mathbb{X}(\mu)_\mathbb{Z}$, and $\mathbb{X}(\lambda + \mu)_\mathbb{Z}$.

Proof. In view of Proposition 2.25, the $w = w_0$ case follows from Lemma 2.17 1). Thanks to Theorem 1.3 and Lemma 2.17 3), we deduce the general case from the $w = w_0$ case. \square

Let $w \in W_{\text{af}}$ and $\mathbf{J} \subset \mathbf{I}$. We define P_+ - and $P_{\mathbf{J},+}$ - graded \mathbb{Z} -modules:

$$R^{\text{af}} := \bigoplus_{\Lambda \in P_+^{\text{af}}} L(\Lambda)_\mathbb{Z}^\vee \quad \text{and} \quad R_w(\mathbf{J}) := \bigoplus_{\lambda \in P_{\mathbf{J},+}} \mathbb{W}_{ww_0}(\lambda)_\mathbb{Z}^\vee.$$

Lemma 2.27. *The \mathbb{Z} -duals of the horizontal maps in Proposition 2.25 equip R^{af} and $R_w(\mathbf{J})$ structures of (P_+ - and $P_{\mathbf{J},+}$ -) graded commutative rings.*

Proof. The maps in Proposition 2.25 are characterized as the d -degree zero maps of cyclic $U_\mathbb{Z}^-$ -modules, that are unique up to a scalar. Therefore, the composition

$$\mathbb{W}(\lambda + \mu + \gamma)_\mathbb{Z} \hookrightarrow \mathbb{W}(\lambda + \mu)_\mathbb{Z} \otimes_\mathbb{Z} \mathbb{W}(\gamma)_\mathbb{Z} \hookrightarrow \mathbb{W}(\lambda)_\mathbb{Z} \otimes_\mathbb{Z} \mathbb{W}(\mu)_\mathbb{Z} \otimes_\mathbb{Z} \mathbb{W}(\gamma)_\mathbb{Z}$$

is the same map as

$$\mathbb{W}(\lambda + \mu + \gamma)_\mathbb{Z} \hookrightarrow \mathbb{W}(\lambda)_\mathbb{Z} \otimes_\mathbb{Z} \mathbb{W}(\mu + \gamma)_\mathbb{Z} \hookrightarrow \mathbb{W}(\lambda)_\mathbb{Z} \otimes_\mathbb{Z} \mathbb{W}(\mu)_\mathbb{Z} \otimes_\mathbb{Z} \mathbb{W}(\gamma)_\mathbb{Z}$$

for every $\lambda, \mu, \gamma \in P_+$ as the images of the cyclic vectors are the same. Taking their restricted duals implies the associativity of the multiplication of $R_e(\mathbf{J})$. In view of Lemma 2.17 1) and Corollary 2.26, we deduce the associativity of the multiplication of $R_w(\mathbf{J})$ for each $w \in W_{\text{af}}$. The associativity of R^{af} is proved similarly (cf. [49]). The commutativity of R^{af} and $R_w(\mathbf{J})$ follow as the $\mathfrak{q} = 1$ coproduct of $U_\mathbb{Z}$ is symmetric ([66, Lemma 3.1.4, §23.1.5]). \square

Corollary 2.28 (of the proof of Lemma 2.27). *For each $w \in W_{\text{af}}$ and $\beta \in Q^\vee$, we have an isomorphism of $(R_w)_\mathbb{Z}$ and $(R_{wt_\beta})_\mathbb{Z}$ as graded commutative rings equipped with $U_\mathbb{Z}^+$ -actions up to grading twists (given by Lemma 2.17). \square*

We set $R := R_e$. Note that $R_w(\mathbf{J}) \subset R_w$ is a subring. We also define

$$R^+(\mathbf{J}) := \bigoplus_{\lambda \in P_{\mathbf{J},+}} \text{Span}_\mathbb{Z} \prod_{i \in \mathbf{I}} (\mathbb{X}(\varpi_i)_\mathbb{Z}^\vee)^{\langle \alpha_i^\vee, \lambda \rangle} \subset \bigoplus_{\lambda \in P_+} \mathbb{X}(\lambda)_\mathbb{Z}^\vee =: \tilde{R},$$

where the multiplication is defined through the projective limit formed by the duals of Corollary 2.26. Here we warn that the (\mathbb{Z} -)rank of some P^{af} -weight space of $\mathbb{X}(\lambda)_\mathbb{Z}$ can be infinity, and hence the inclusion $R^+ \subsetneq \tilde{R}$ has a huge cokernel. Note that the rank of the P^{af} -weight spaces of $\mathbb{X}(\varpi_i)_\mathbb{Z}$ are bounded for each $i \in \mathbf{I}$ ([44, Proposition 5.16]), and hence $R^+(\mathbf{J})$ has only countably many generators of $P_{\mathbf{J},+}$ -degrees $\{\varpi_i\}_{i \in \mathbf{I} \setminus \mathbf{J}}$.

By construction, the rings $R^{\text{af}}, R_w(\mathbf{J})$, and $R^+(\mathbf{J})$ are free over \mathbb{Z} .

For each $\lambda \in P_+$ and $w \in W_{\text{af}}$, we have a unique P^{af} -weight vector

$$\mathbf{v}_{w\lambda}^\vee \in \mathbb{W}_w(\lambda)_\mathbb{Z}^\vee \tag{2.7}$$

with paring 1 with $\mathbf{v}_{w\lambda} \in \mathbb{W}_w(\lambda)_\mathbb{Z}$. This vector $\mathbf{v}_{w\lambda}^\vee$ yields an $U_\mathbb{Z}^+$ -cocyclic vector of $\mathbb{W}_w(\lambda)_\mathbb{Z}^\vee$ for a field \mathbb{k} . By the construction of the ring structure on R_w , we have $\mathbf{v}_{w_0\lambda}^\vee \cdot \mathbf{v}_{w_0\mu}^\vee = \mathbf{v}_{w_0(\lambda+\mu)}^\vee$ in R_w for every $\lambda, \mu \in P_+$.

Lemma 2.29. *For each $w, v \in W_{\text{af}}$ and $J \subset \mathbf{I}$, the ring $R_w(J)$ is a quotient of $R_v(J)$ if $w \leq_{\frac{\infty}{2}} v$. In addition, the ring R_w is a quotient of R_v if and only if $w \leq_{\frac{\infty}{2}} v$.*

Proof. We have $\mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}} \subset \mathbb{W}_{vw_0}(\lambda)_{\mathbb{Z}}$ if and only if $\mathbf{v}_{ww_0\lambda} \in \mathbb{W}_{vw_0}(\lambda)_{\mathbb{Z}}$. Now we apply Lemma 2.15 to deduce the result. \square

Lemma 2.30. *We have the following morphisms of rings with $U_{\mathbb{Z}}^+$ -actions*

$$R^+ \twoheadrightarrow R \hookrightarrow R^{\text{af}},$$

that admit \mathbb{Z} -module splittings, where the $\dot{U}_{\mathbb{Z}}$ -action on R^{af} is twisted by θ .

Proof. Apply Proposition 2.25 and Corollary 2.26. \square

For each $w \in W$ and $J \subset \mathbf{I}$, we set

$$(\mathbf{Q}_{G,J}(w))_{\mathbb{Z}} := \text{Proj } R_w(J) \quad \text{and} \quad (\mathbf{Q}_{G,J}^{\text{rat}})_{\mathbb{Z}} := \bigcup_{w \in W} (\mathbf{Q}_{G,J}(w))_{\mathbb{Z}}.$$

These schemes and indspaces are flat over \mathbb{Z} .

Remark 2.31. In view of (1.5), we have embeddings

$$\begin{array}{ccc} \text{Proj } R_{\mathbb{K}}^+ & \longrightarrow & \prod_{i \in \mathbf{I}} \mathbb{P}_{\mathbb{K}}(\mathbb{X}(\varpi_i)_{\mathbb{K}}^{\wedge}), \\ \uparrow & & \uparrow \\ (\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}} & \longrightarrow & \prod_{i \in \mathbf{I}} \mathbb{P}_{\mathbb{K}}(\mathbb{X}(\varpi_i)_{\mathbb{K}}^{\flat}) \end{array} \quad (2.8)$$

where we set

$$\mathbb{X}(\varpi_i)_{\mathbb{K}}^{\wedge} := \prod_{n \in \mathbb{Z}} \mathbb{X}(\varpi_i)_{n,\mathbb{K}} \quad \text{and} \quad \mathbb{X}(\varpi_i)_{\mathbb{K}}^{\flat} := \bigcup_{m \in \mathbb{Z}} \prod_{n > m} \mathbb{X}(\varpi_i)_{n,\mathbb{K}}$$

for each $i \in \mathbf{I}$. Here all the spaces in (2.8) admit actions of $SL(2, i)_{\mathbb{K}}$ ($i \in \mathbf{I}$), while only the bottom two spaces in (2.8) admit actions of $G[[z]]_{\mathbb{K}}$. Nevertheless, the top two spaces in (2.8) have some advantages since they are θ -stable (unlike the bottom two), and consequently also contain $\theta((\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}})$.

Theorem 2.32 ([49] Corollary B). *Let p be a prime. Then, the ring $R_{\mathbb{F}_p}^{\text{af}}$ admits a Frobenius splitting, that is \mathbf{I} - and \mathbf{I}^- -canonically split.* \square

Theorem 2.33. *Let p be a prime. The ring $R_{\mathbb{F}_p}$ admits a Frobenius splitting, that is \mathbf{I} -canonically split.*

Proof. The \mathbf{I} -canonical Frobenius splitting ϕ of R^{af} gives rise to the following maps, whose composition is the identity:

$$L(\Lambda)_{\mathbb{F}_p} \xrightarrow{\phi^{\vee}} L(p\Lambda)_{\mathbb{F}_p} \longrightarrow L(\Lambda)_{\mathbb{F}_p} \quad \Lambda \in P_+^{\text{af}}.$$

In view of Proposition 2.25, it prolongs to

$$\begin{array}{ccccc} L(\Lambda)_{\mathbb{F}_p} & \xrightarrow{\phi^{\vee}} & L(p\Lambda)_{\mathbb{F}_p} & \longrightarrow & L(\Lambda)_{\mathbb{F}_p} & \Lambda \in P_+^{\text{af}}. \\ \downarrow \pi_{\Lambda} & & \downarrow \pi_{p\Lambda} & & \downarrow & \\ \mathbb{W}^-(\bar{\Lambda})_{\mathbb{F}_p} & \xrightarrow{\phi_{\mathbb{W}}^{\vee}} & \mathbb{W}^-(p\bar{\Lambda})_{\mathbb{F}_p} & \longrightarrow & \mathbb{W}^-(\bar{\Lambda})_{\mathbb{F}_p} & \end{array}$$

The right square is automatic (and is canonically defined) from the adjunction of the Frobenius push-forward (by taking the restricted dual). In order to show that ϕ descends to a Frobenius splitting of $R_{\mathbb{F}_p}$, it suffices to show that the dotted map $\phi_{\mathbb{W}}^{\vee}$ is a well-defined linear map (induced from ϕ^{\vee} and so that the left square is commutative).

By Corollary 2.24, $\ker \pi_{\Lambda}$ is generated by the P -weight $(P \setminus \text{Conv } W\bar{\Lambda})$ -part of $L(\Lambda)_{\mathbb{F}_p}$. By the cyclicity of $L(\Lambda)_{\mathbb{F}_p}$ as $U_{\mathbb{Z}}^{-}$ -modules and Proposition 2.7, we deduce that $\phi^{\vee}(\ker \pi_{\Lambda})$ is contained in the $U_{\mathbb{F}_p}^{-}$ -submodule of $L(p\Lambda)_{\mathbb{F}_p}$ generated by the P -weight $p(P \setminus \text{Conv } W\bar{\Lambda})$ -part of $L(p\Lambda)_{\mathbb{F}_p}$. The latter is contained in $\ker \pi_{p\Lambda}$ by

$$p(P \setminus \text{Conv } W\bar{\Lambda}) \subset P \setminus p\text{Conv } W\bar{\Lambda}.$$

Therefore, we conclude that $\phi_{\mathbb{W}}^{\vee}$ is a well-defined linear map, and hence $\theta^*(R_{\mathbb{F}_p})$ admits a Frobenius splitting induced from ϕ . The unipotent part of the \mathbf{I}^{-} -canonical splitting condition is in common with subrings. It remains to twist the grading given by α_0^{\vee} with that given by $-\vartheta^{\vee}$ and twist the \mathbf{I}^{-} -action into the \mathbf{I} -action by θ to conclude that our Frobenius splitting on $R_{\mathbb{F}_p}$ is \mathbf{I} -canonical. \square

Corollary 2.34. *Let p be a prime, and let $w \in W$. The \mathbf{I} -canonical splitting of $R_{\mathbb{F}_p}$ obtained in Theorem 2.33 induces an \mathbf{I} -canonical splitting of $(R_w)_{\mathbb{F}_p}$.*

Proof. We set $L^{ww_0}(\Lambda)_{\mathbb{Z}} := U_{\mathbb{Z}}^{-} \mathbf{v}_{ww_0\Lambda}$, where $\mathbb{Z} \mathbf{v}_{ww_0\Lambda}$ is the P^{af} -weight $ww_0\Lambda$ -part of $L(\Lambda)_{\mathbb{Z}}$, that is rank one over \mathbb{Z} .

The subspace $\mathbb{W}_{ww_0}^{-}(\lambda)_{\mathbb{Z}} \subset \mathbb{W}^{-}(\lambda)_{\mathbb{Z}}$ is the image of $L^{ww_0}(\Lambda)_{\mathbb{Z}} \subset L(\Lambda)_{\mathbb{Z}}$ (with $\lambda = \bar{\Lambda}$) under Theorem 2.23 as $L^{ww_0}(\Lambda)_{\mathbb{Z}}$ is spanned by a subset of $\mathbf{B}(\Lambda)$ ([43, (0.3)]). Our Frobenius splitting ϕ is obtained from that of $R_{\mathbb{F}_p}^{\text{af}}$, which is compatible with $\bigoplus_{\lambda \in P_+} L^{ww_0}(\Lambda)_{\mathbb{F}_p}^{\vee}$ by [49, Corollary B]. Applying θ , we conclude that ϕ must descend to a Frobenius splitting of $(R_w)_{\mathbb{F}_p}$. \square

Corollary 2.35. *An \mathbf{I} -canonical splitting of $R_{\mathbb{F}_p}$ is unique.*

Proof. The behavior of the vectors in (2.7) (with $w = e$) under an \mathbf{I} -canonical Frobenius splitting is uniquely determined as they form a polynomial ring isomorphic to $\mathbb{F}_p P_+$ such that each of its (P_+) -graded component is a multiplicity-free P^{af} -weight space in $\mathbb{W}(\lambda)_{\mathbb{F}_p}^{\vee}$'s. By Proposition 2.7, this completely determines the behavior of our splitting (through its dual map). \square

Corollary 2.36. *An \mathbf{I} -canonical splitting of $R_{\mathbb{F}_p}$ is compatible with $(R_w)_{\mathbb{F}_p}$ for every $w \in W_{\text{af}}$ such that $w \leq_{\frac{\infty}{2}} e$.*

Proof. By [15, Proposition 4.1.17 and Remark 4.1.18 (i)] and [46, Theorem 4.12] (the algebraic portion of the latter stems from [45, Lemma 2.6], that carries over to this setting; cf. [46, Lemma 4.4 and Theorem 4.7]), we derive that an \mathbf{I} -canonical Frobenius splitting of $(R_w)_{\mathbb{F}_p}$ ($w \in W$) gives rise to an \mathbf{I} -canonical splitting of $R_{\mathbb{F}_p}$ that is compatible with $(R_w)_{\mathbb{F}_p}$ (arguing by restricting to the $SL(2, i)$ -actions for each $i \in \mathbf{I}$). In particular, the \mathbf{I} -canonical splitting of $(R_w)_{\mathbb{F}_p}$ ($w \in W$) also uniquely exist and compatible with that of $R_{\mathbb{F}_p}$ by Corollary 2.35 and Corollary 2.34. By Corollary 2.28, we further deduce that the \mathbf{I} -canonical splitting of $(R_w)_{\mathbb{F}_p}$ ($w \in W_{\text{af}}$) uniquely exists.

Let $w \in W$ such that $s_0 w = s_{\vartheta} w t_{-w^{-1}\vartheta^{\vee}} \leq_{\frac{\infty}{2}} w$. Then, $R_{s_0 w}$ is a quotient of R_w . Again by [15, Proposition 4.1.17 and Remark 4.1.18 (i)], the set of \mathbf{I} -canonical splittings of $(R_{s_0 w})_{\mathbb{F}_p}$ is in bijection with that of $(R_w)_{\mathbb{F}_p}$ compatible

with $(R_{s_0 w})_{\mathbb{F}_p}$. By Corollary 2.28 and the above paragraph, we find that the (unique) \mathbf{I} -canonical splitting of $(R_{s_0 w})_{\mathbb{F}_p}$ is compatible with $(R_w)_{\mathbb{F}_p}$, and hence also compatible with $R_{\mathbb{F}_p}$ and $(R_{t_{-w^{-1}\vartheta^\vee}})_{\mathbb{F}_p}$. This forces the \mathbf{I} -canonical splitting of $R_{\mathbb{F}_p}$ to be compatible with that of $(R_{ut_{-w^{-1}\vartheta^\vee}})_{\mathbb{F}_p}$ for every $u, w \in W$ such that $-w^{-1}\vartheta^\vee \in \Delta_+^\vee$.

The set $(\{-w^{-1}\vartheta^\vee\}_{w \in W} \cap Q_+^\vee)$ is precisely the set of short positive coroots in Δ_+^\vee . This spans Q_+^\vee as monoids by inspection. Since $w \in W_{\text{af}}$ with $w \leq_{\frac{\infty}{2}} e$ is written as $w = ut_\beta$ for some $u \in W$ and $\beta \in Q_+^\vee$ ([74, Lecture 13, Proposition 1]), we conclude the assertion. \square

Theorem 2.37. *Let p be a prime and let $J \subset \mathbf{I}$. The ring $R^+(J)_{\mathbb{F}_p}$ admits a Frobenius splitting that is \mathbf{I} - and \mathbf{I}^- -canonically split. This splitting is compatible with $R_w(J)_{\mathbb{F}_p}$ and the image of $R^+(J)_{\mathbb{F}_p} \subset R_{\mathbb{F}_p}^+$ under the quotient map $R_{\mathbb{F}_p}^+ \twoheadrightarrow \theta^*((R_{ww_0})_{\mathbb{F}_p})$ for each $w \in W_{\text{af}}$.*

Proof. Since the case of $J \neq \emptyset$ follows by the restriction to a part of the P_+ -grading, we concentrate into the case $J = \emptyset$.

The ring structure of $R_{\mathbb{F}_p}^+$ is determined by $R_{\mathbb{F}_p}$ through the application of $U_{\mathbb{Z}}^-$ before taking duals. By Corollary 2.36 (and its proof), it defines an \mathbf{I} -canonical splitting ϕ of $R_{\mathbb{F}_p}^+$ compatible with $(R_w)_{\mathbb{F}_p}$ for each $w \in W_{\text{af}}$. The ring $R_{\mathbb{F}_p}^+$ admits an $SL(2, i)_{\mathbb{F}_p}$ -action that integrates the actions of $E_i^{(n)}$ and $F_i^{(n)}$ ($n \in \mathbb{Z}_{>0}$) for each $i \in \mathbf{I}_{\text{af}}$. By [15, Proposition 2.10], this splitting ϕ is also \mathbf{I}^- -canonical.

The \mathbf{I} -cocyclic P^{af} -weight vector $\mathbf{v}_{ww_0\lambda}^\vee \in \mathbb{W}_{ww_0}(\lambda)_{\mathbb{F}_p}^\vee$ is uniquely characterized by its P^{af} -weight. Hence, we obtain a map

$$\mathbb{X}(\lambda)_{\mathbb{F}_p}^\vee \supset \text{Span}_{\mathbb{Z}} \prod_{i \in \mathbf{I}} (\mathbb{X}(\varpi_i)_{\mathbb{F}_p}^\vee)^{\langle \alpha_i, \lambda \rangle} \twoheadrightarrow \mathbb{W}_{ww_0}(\lambda)_{\mathbb{F}_p}^\vee \twoheadrightarrow \mathbb{F}_p \mathbf{v}_{ww_0\lambda}^\vee.$$

It gives rise to the ring surjections

$$R_{\mathbb{F}_p}^+ \twoheadrightarrow (R_w)_{\mathbb{F}_p} \twoheadrightarrow \bigoplus_{\lambda \in P_+} \mathbb{F}_p \mathbf{v}_{ww_0\lambda}^\vee$$

that is compatible with ϕ by construction in the first surjection and by examining the P^{af} -weights in the second surjection (we denote this composition surjective ring map by ξ). Consider the ideal

$$I(w) := R_{\mathbb{F}_p}^+ \cap \bigcap_{g \in \mathbf{I}^-(\overline{\mathbb{F}}_p)} g(\overline{\mathbb{F}}_p \otimes_{\mathbb{F}_p} \ker \xi) \subset R_{\overline{\mathbb{F}}_p}^+.$$

(Here the action of $\mathbf{I}^-(\overline{\mathbb{F}}_p)$ is obtained by the unipotent one-parameter subgroups $\{\rho_{-\alpha_i}\}_{i \in \mathbf{I}_{\text{af}}}$ defined through the exponentials, that are well-defined as we have all the divided powers. The passage from \mathbb{F}_p to $\overline{\mathbb{F}}_p$ is necessary to ensure the scheme-theoretic invariance since we want an intersection that is equivalent to the geometric \mathbb{G}_a -actions through $\rho_{-\alpha_i}$ for all $i \in \mathbf{I}_{\text{af}}$ by finding the Zariski dense subsets of \mathbb{G}_a . For this purpose, we want the image of each $\rho_{-\alpha_i}$ to be infinite, that cannot be achieved by sending $\mathbb{G}_a(\mathbb{F}_p) \cong \mathbb{F}_p$.) This ideal is the maximal $U_{\overline{\mathbb{F}}_p}^-$ -invariant ideal of $R_{\overline{\mathbb{F}}_p}^+$ that is contained in $\ker \xi$. Let us denote the quotient ring by

$$Q = \bigoplus_{\lambda \in P_+} Q(\lambda) := R_{\mathbb{F}_p}^+ / I(w).$$

By the construction of $I(w)$, we deduce that

$$U_{\mathbb{F}_p}^- \mathbf{v}_{ww_0\lambda} \subset Q(\lambda)^\vee \subset \mathbb{X}(\lambda)_{\mathbb{F}_p}$$

for each $\lambda \in P_+$ (otherwise we can derivate a vector in $I(w)$ to obtain a non-zero element of $\bigoplus_{\lambda \in P_+} \mathbb{F}_p \mathbf{v}_{ww_0\lambda}^\vee$). Since $\theta^*(\mathbb{W}_{ww_0}(-w_0\lambda)_{\mathbb{F}_p})$ is a cyclic $U_{\mathbb{F}_p}^-$ -submodule of $\mathbb{X}(\lambda)_{\mathbb{F}_p}$, we have $U_{\mathbb{F}_p}^- \mathbf{v}_{ww_0\lambda} = \theta^*(\mathbb{W}_{ww_0}(-w_0\lambda)_{\mathbb{F}_p})$. In particular, we deduce a vector space surjection

$$R_{\mathbb{F}_p}^+ / I(w) \twoheadrightarrow \theta^*((R_{ww_0})_{\mathbb{F}_p})$$

(cf. Corollary 2.26). Since the RHS is naturally a ring, we conclude

$$R_{\mathbb{F}_p}^+ / I(w) \cong \theta^*((R_{ww_0})_{\mathbb{F}_p})$$

by the maximality of $I(w)$.

The ideal $I(w) \subset R_{\mathbb{F}_p}^+$ also splits compatibly by ϕ (since ϕ is \mathbf{I}^- -canonical and $\ker \xi$ splits compatibly). In particular, each $\theta^*((R_{ww_0})_{\mathbb{F}_p})$ compatibly split under ϕ as required. \square

Corollary 2.38. *For each $\mathbf{J} \subset \mathbf{I}$, the indscheme $(\mathbf{Q}_{G,\mathbf{J}}^{\text{rat}})_{\mathbb{F}_p}$ admits an \mathbf{I} - and \mathbf{I}^- -canonical Frobenius splitting that is compatible with $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{F}_p}$ for each $w \in W_{\text{af}}$.*

Proof. The condition of the canonical splitting can be checked by line bundle twists, and hence we only need to show whether the Frobenius splitting of $R_w(\mathbf{J})_{\mathbb{F}_p}$ descends to $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{F}_p}$ for each $w \in W_{\text{af}}$. Since $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{F}_p}$ is the quotient of an open subset of $X := \text{Spec } R_w(\mathbf{J})_{\mathbb{F}_p}$ by H (corresponding to the $P_{\mathbf{J},+}$ -grading) and our Frobenius splitting is H -fixed, it suffices to see whether the localization to the non-irrelevant locus preserves the Frobenius splitting. This follows if the localization of $\text{Fr}_* \mathcal{O}_X = \mathcal{O}_{X^{(1)}}$ as a $\text{Fr}_* \mathcal{O}_X$ -module and as a \mathcal{O}_X -module are the same. It holds as the localization by a multiplicative set S is the same as the localization by S^p . \square

Recall that a scheme X defined over a field \mathbb{k} is called weakly normal if every finite bijective birational (\mathbb{k} -)morphism $f : Y \rightarrow X$ from a (reduced) scheme over \mathbb{k} is in fact an isomorphism ([68] and [15, §1.2.3]).

Corollary 2.39. *For each $\mathbf{J} \subset \mathbf{I}$, the indscheme $(\mathbf{Q}_{G,\mathbf{J}}^{\text{rat}})_{\mathbb{F}_p}$ and the schemes $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{F}_p}$ ($w \in W_{\text{af}}$) are reduced. In addition, $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{F}_p}$ is weakly normal.*

Proof. For the first assertion, apply [15, Proposition 1.2.1] to Corollary 2.38. For the second assertion, apply [15, Proposition 1.2.5] to Corollary 2.34. \square

3 Frobenius splitting of quasi-map spaces

We retain the settings of the previous section. In particular, we sometimes work over a ring or a non-algebraically closed field. Moreover, the notational convention explained in the beginning of §2 continue to apply.

3.1 The scheme $\mathcal{Q}'_J(v, w)$ and its Frobenius splitting

Let $v, w \in W_{\text{af}}$ and $J \subset \mathbf{I}$. We set

$$R_w^v(\mathbf{J}) := R^+(\mathbf{J}) / (\ker(R^+(\mathbf{J}) \rightarrow R_w(\mathbf{J})) + \ker(R^+(\mathbf{J}) \rightarrow \theta^*(R_{vw_0}(\mathbf{J}')))),$$

where we have $\ker(R^+(\mathbf{J}) \rightarrow \theta^*(R_{vw_0}(\mathbf{J}')))) = \ker(R^+(\mathbf{J}) \rightarrow \theta^*(R_{vw_0}))$ for

$$\mathbf{J}' := \{i \in \mathbf{I} \mid -w_0\alpha_i = \alpha_j, \exists j \in \mathbf{J}\} \subset \mathbf{I}.$$

By construction, $R_w^v(\mathbf{J})$ is a $P_{\mathbf{J},+}$ -graded ring. We set

$$R_w^v(\mathbf{J}) := \bigoplus_{\lambda \in P_{\mathbf{J},+}} R_w^v(\mathbf{J}, \lambda).$$

Lemma 3.1. *For each $w, v \in W_{\text{af}}$ and $J \subset \mathbf{I}$, the multiplication map*

$$R_w^v(\mathbf{J}, \lambda) \otimes_{\mathbb{Z}} R_w^v(\mathbf{J}, \mu) \rightarrow R_w^v(\mathbf{J}, \lambda + \mu) \quad (\lambda, \mu \in P_{\mathbf{J},+})$$

is surjective.

Proof. We have a quotient

$$R_w(\mathbf{J}) = \bigoplus_{\lambda \in P_{\mathbf{J},+}} \mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}}^{\vee} \twoheadrightarrow \bigoplus_{\lambda \in P_{\mathbf{J},+}} R_w^v(\mathbf{J}, \lambda) = R_w^v(\mathbf{J})$$

of homogeneous rings. Corollary 2.26 implies that the multiplication map of $R_w(\mathbf{J})$ is surjective. Hence, so is the quotient ring. \square

We set

$$\mathcal{Q}'_J(v, w) := \text{Proj } R_w^v(\mathbf{J}),$$

where our definition of Proj is (1.5). In case $v = w_0 t_{\beta}$ for $\beta \in Q^{\vee}$, we may write $R_w^v(\mathbf{J})$ and $\mathcal{Q}'_J(v, w)$ by $R_w^{\beta}(\mathbf{J})$ and $\mathcal{Q}'_J(\beta, w)$, respectively.

Lemma 3.2. *For each $w, v \in W$, the Chevalley involution induces an isomorphism*

$$\mathcal{Q}'(v, w) \xrightarrow{\cong} \mathcal{Q}'(ww_0, vw_0).$$

Proof. Apply Lemma 2.17 to the construction of R_w^v to deduce an isomorphism

$$\theta^* : R_w^v \xrightarrow{\cong} R_{vw_0}^{ww_0}$$

of graded rings, that yields the assertion. \square

Lemma 3.3. *For each $w, v \in W_{\text{af}}$ and $J \subset \mathbf{I}$, the scheme $\mathcal{Q}'_J(v, w)$ is flat over \mathbb{Z} .*

Proof. The ring $R^+(\mathbf{J})$ has a \mathbb{Z} -basis that is dual to $\bigsqcup_{\lambda \in P_{\mathbf{J},+}} \mathbf{B}(\mathbb{X}(\lambda))$. The rings $R_w(\mathbf{J})$ and $\text{Im}(R^+(\mathbf{J}) \rightarrow \theta^*(R_{vw_0}))$ are quotients by subsets of such basis elements by Lemma 2.17. Hence, we have a free \mathbb{Z} -basis of $R_w^v(\mathbf{J})$ as required. \square

Lemma 3.4. *For each $w, v \in W_{\text{af}}$ and $J \subset \mathbf{I}$, the scheme $\mathcal{Q}'_J(v, w)$ is projective (of finite type) over \mathbb{Z} .*

Proof. By [44, Proposition 5.16], we have $\text{rank}_{\mathbb{Z}} R_w^v(\varpi_i) < \infty$ for each $i \in \mathbf{I}$. By Lemma 3.1, this forces $\mathcal{Q}'_{\mathbf{J}}(v, w)$ to be a projective scheme (that usually implies finite type by definition. Here we explicitly include it since our Proj does not yield a finite type scheme in general) as required. \square

Lemma 3.5. *Let $\mathbf{J} \subset \mathbf{I}$. We have $\mathcal{Q}'_{\mathbf{J}}(v, w) \neq \emptyset$ if $v \leq_{\frac{\infty}{2}} w$.*

Proof. We have $R_w^v(\mathbf{J}, \lambda) \neq \{0\}$ if $\mathbf{v}_{vw_0\lambda} \in \mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}}$ by Lemma 2.17 4). Here $\mathbf{v}_{vw_0\lambda} \in \mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}}$ is equivalent to $\mathbb{W}_{vw_0}(\lambda)_{\mathbb{Z}} \subset \mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}}$. Now apply Lemma 2.15 to obtain the assertion. \square

Lemma 3.6. *Let $v, w \in W_{\text{af}}$, $\beta \in Q^{\vee}$, and $\mathbf{J} \subset \mathbf{I}$. We have $\mathcal{Q}'_{\mathbf{J}}(v, w) \cong \mathcal{Q}'_{\mathbf{J}}(vt_{\beta}, wt_{\beta})$.*

Proof. We borrow notation from Lemma 2.17. By the definition of our ring $R_w^v(\mathbf{J})$, the assertion follows if

$$\mathbb{W}_{wt_{\beta}w_0}(\lambda)_{\mathbb{Z}} = \tau_{w_0\beta} \mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}} \quad \text{and} \quad \tau_{w_0\beta} \theta^*(\mathbb{W}_v(-w_0\lambda)_{\mathbb{Z}}) = \theta^*(\mathbb{W}_{vt_{\beta}}(-w_0\lambda)_{\mathbb{Z}})$$

holds for each $w, v \in W_{\text{af}}$, $\beta \in Q^{\vee}$, and $\lambda \in P_+$. These assertions themselves follow by chasing the weights of the cyclic vectors. \square

Lemma 3.7. *Let p be a prime and let $\mathbf{J} \subset \mathbf{I}$. For each $w, v \in W_{\text{af}}$, the ring $R_w^v(\mathbf{J})_{\mathbb{F}_p}$ admits a Frobenius splitting that is compatible with the quotient $R_w^{v'}(\mathbf{J})_{\mathbb{F}_p}$ for $v', w' \in W_{\text{af}}$ such that $v \leq_{\frac{\infty}{2}} v' \leq_{\frac{\infty}{2}} w' \leq_{\frac{\infty}{2}} w$. In particular, the scheme $\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{F}_p}$ is reduced and weakly normal.*

Proof. By construction, $\ker(R^+(\mathbf{J})_{\mathbb{F}_p} \rightarrow (R_w)_{\mathbb{F}_p})$ and $\ker(R^+(\mathbf{J})_{\mathbb{F}_p} \rightarrow \theta^*((R_{vw_0})_{\mathbb{F}_p}))$ are ideals of a ring $R^+(\mathbf{J})_{\mathbb{F}_p}$ that are compatible with the canonical Frobenius splitting of $R^+(\mathbf{J})_{\mathbb{F}_p}$ by Theorem 2.37. Hence, so is their sum. It must be compatible with every quotient of the form $R_w^{v'}(\mathbf{J})_{\mathbb{F}_p}$ with the above condition by Lemma 2.29, Corollary 2.34, Lemma 2.17 4), and Lemma 3.5. This proves the first assertion. We apply [15, Proposition 1.2.1 and 1.2.5] to deduce the second assertion. \square

Corollary 3.8. *Let p be a prime. For each $w, v \in W$, the scheme $\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{F}_p}$ admits a Frobenius splitting compatible with $\mathcal{Q}'_{\mathbf{J}}(v', w')_{\mathbb{F}_p}$ for every $w', v' \in W_{\text{af}}$ such that $v \leq_{\frac{\infty}{2}} v' \leq_{\frac{\infty}{2}} w' \leq_{\frac{\infty}{2}} w$.*

Proof. Apply Theorem 2.5 to Lemma 3.7. \square

Remark 3.9. Unlike the case of Corollary 2.38, the space $\text{Spec } R_w^v(\mathbf{J})_{\mathbb{F}_p}$ is not irreducible in general. In fact, we discard some of the irreducible components of $\text{Spec } R_w^v(\mathbf{J})_{\mathbb{F}_p}$ from Lemma 3.7 to Corollary 3.8.

Corollary 3.10. *The ring $R_w^v(\mathbf{J})$ is reduced. In particular, the scheme $\mathcal{Q}'_{\mathbf{J}}(v, w)$ is reduced.*

Proof. By Lemma 3.3, every non-zero element of $R_w^v(\mathbf{J})$ is annihilated by reduction mod p for only finitely many primes. Now it remains to apply Lemma 3.7. \square

3.2 Modular interpretation of $\mathbf{Q}_{G,\mathbf{J}}^{\text{rat}}$

We have an identification

$$W_{\text{af}} \cong N_{G((z))}(H(\mathbb{K}))/H(\mathbb{K})$$

regardless of the (algebraically closed) base field \mathbb{K} . We denote a lift of $w \in W_{\text{af}}$ in $N_{G((z))}(H(\mathbb{K}))/H(\mathbb{K})$ by \dot{w} .

Lemma 3.11. *For each $w \in W$ and $\mathbf{J} \subset \mathbf{I}$, the scheme $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{K}}$ contains an affine Zariski open $\mathbf{I}_{\mathbb{K}}$ -orbit $\mathcal{O}(\mathbf{J}, w)_{\mathbb{K}}$ that is isomorphic to*

$$\mathbf{I}_{\mathbb{K}} / (H_{\mathbb{K}} \cdot (\text{Ad}(\dot{w}\dot{w}_0)([P(\mathbf{J}), P(\mathbf{J})]((z))) \cap \mathbf{I}_{\mathbb{K}}))$$

as a scheme over \mathbb{K} . (By abuse of notation, here we identify the set of \mathbb{K} -valued points $(\text{Ad}(\dot{w}\dot{w}_0)([P(\mathbf{J}), P(\mathbf{J})]((z))) \cap \mathbf{I}_{\mathbb{K}}$ with its Zariski closure in $\mathbf{I}_{\mathbb{K}}$). It is an open neighbourhood of the unique $(H \times \mathbb{G}_m)_{\mathbb{K}}$ -fixed point of $\mathcal{O}(\mathbf{J}, w)_{\mathbb{K}}$.

Proof. Recall that $\mathbf{v}_{ww_0\lambda}^{\vee} \cdot \mathbf{v}_{ww_0\mu}^{\vee} = \mathbf{v}_{ww_0(\lambda+\mu)}^{\vee}$ for each $\lambda, \mu \in P_+$. Since $\mathbb{W}_{ww_0}(\lambda)$ is compatible with the positive global basis, the ring

$$\mathbb{Z}[w] := \sum_{\lambda \in P_{\mathbf{J},+}} (\mathbf{v}_{ww_0\lambda}^{\vee})^{-1} \mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}}^{\vee} \subset (U_{\mathbb{Z}}^+)^{\vee} \quad (3.1)$$

admits its dual basis. By construction, $\mathbb{Z}[w]$ is the coordinate ring of an affine Zariski open set of $\mathbf{Q}_G(w)_{\mathbb{Z}}$. In addition, it inherits a natural P^{af} -grading from $R_w(\mathbf{J})$. Therefore, $\mathbb{Z}[w]$ defines an open neighbourhood of a $(H \times \mathbb{G}_m)$ -fixed point of $\mathbf{Q}_{G,\mathbf{J}}(w)$ obtained by the linear forms $\{\mathbf{v}_{ww_0\lambda}\}_{\lambda}$. We set $\mathbb{C}[w] := \mathbb{C} \otimes_{\mathbb{Z}} \mathbb{Z}[w]$ and $\mathbb{K}[w] := \mathbb{K} \otimes_{\mathbb{Z}} \mathbb{Z}[w]$.

The defining relation of $(\mathbb{C}[e])^{\vee}$ in terms of the $U_{\mathbb{C}}^+$ -action is

$$U_{\mathbb{C}}^+(\text{Ad}(\dot{w}_0)([\mathfrak{p}(\mathbf{J}), \mathfrak{p}(\mathbf{J})]) \otimes \mathbb{C}[z] \cap \mathfrak{J}).$$

by (the limit of) Chari-Fourier-Khandai [17, Proposition 3.3]. Since $\mathbb{W}_{uw_0}(\lambda)_{\mathbb{Z}} \subset \mathbb{W}_{w_0}(\lambda)_{\mathbb{Z}}$ for each $u \in W$, the defining relation of $(\mathbb{C}[u])^{\vee}$ in terms of the $U_{\mathbb{C}}^+$ -action is

$$U_{\mathbb{C}}^+(\text{Ad}(\dot{u}\dot{w}_0)([\mathfrak{p}(\mathbf{J}), \mathfrak{p}(\mathbf{J})]) \otimes \mathbb{C}[z] \cap \mathfrak{J})$$

by applying the action of $\dot{u} \in N_G(H)_{\mathbb{C}}$ that lifts $u \in W$. In particular, we have

$$\text{Spec } \mathbb{C}[w] \cong \mathbf{I}_{\mathbb{C}} / (H_{\mathbb{C}} \cdot \text{Ad}(\dot{w}\dot{w}_0)([P(\mathbf{J})_{\mathbb{C}}, P(\mathbf{J})_{\mathbb{C}}][z]) \cap \mathbf{I}_{\mathbb{C}}) \quad (3.2)$$

as schemes over \mathbb{C} . We put $\mathbf{I}_{\mathbb{Z}}^1 := [\mathbf{I}_{\mathbb{Z}}, \mathbf{I}_{\mathbb{Z}}]$. Let $\mathbf{L}_{\mathbb{C}}$ and $\mathbf{R}_{\mathbb{C}}$ be pro-unipotent subgroups of $\mathbf{I}_{\mathbb{C}}^1$ whose closed points are

$$\text{Specm } \mathbb{C}[w] \quad \text{and} \quad \text{Ad}(\dot{w}\dot{w}_0)([P(\mathbf{J})_{\mathbb{C}}, P(\mathbf{J})_{\mathbb{C}}][z]) \cap \mathbf{I}_{\mathbb{C}}^1,$$

respectively and they are stable by the natural $(H \times \mathbb{G}_m)_{\mathbb{C}}$ -action on $\mathbf{I}_{\mathbb{C}}$. The isomorphism (3.2) gives rise to an isomorphism

$$m_{\mathbb{C}} : \mathbf{L}_{\mathbb{C}} \times \mathbf{R}_{\mathbb{C}} \xrightarrow{\cong} \mathbf{I}_{\mathbb{C}}^1 \quad (3.3)$$

of schemes over \mathbb{C} , where $m_{\mathbb{C}}$ is the multiplication map as all the groups are pro-unipotent and the isomorphism between the sets of closed points is compatible with truncations by P^{af} -weight considerations. The Hopf algebra $\mathbb{Z}[\mathbf{I}_{\mathbb{Z}}^1] :=$

$(U_{\mathbb{Z}}^+)^{\vee}$ is the coordinate ring of $\mathbf{I}_{\mathbb{Z}}^1$ (cf. [49, Theorem 1.3]). By sending $\mathbb{Z}[\mathbf{I}^1]$ by the restriction morphisms $\mathbb{C}[\mathbf{I}_{\mathbb{C}}^1] \rightarrow \mathbb{C}[\mathbf{L}_{\mathbb{C}}]$ and $\mathbb{C}[\mathbf{I}_{\mathbb{C}}^1] \rightarrow \mathbb{C}[\mathbf{R}_{\mathbb{C}}]$, we have the corresponding group schemes $\mathbf{L}_{\mathbb{Z}}$ and $\mathbf{R}_{\mathbb{Z}}$ over \mathbb{Z} (we denote their coordinate rings as $\mathbb{Z}[\mathbf{L}]$ and $\mathbb{Z}[\mathbf{R}]$, respectively). For each real root $\alpha \in \Delta_{\text{af},+}$, one of the restricted dual rings $\mathbb{C}[\mathbf{L}_{\mathbb{C}}]^{\vee}$ nor $\mathbb{C}[\mathbf{R}_{\mathbb{C}}]^{\vee}$ contains the image of a primitive element of $U_{\mathbb{Z}}^+$ with $\tilde{\mathfrak{h}}$ -weight α (obtained by conjugations of $\{E_i\}_{i \in \mathbf{I}_{\text{af}}}$'s cf. [66, Proposition 40.1.3] or [31, Lemma 6.6]), and hence the corresponding one-parameter subgroup lands in either of $\mathbf{L}_{\mathbb{Z}}$ or $\mathbf{R}_{\mathbb{Z}}$. In view of [5, Theorem 3.13] (or [31, Theorem 5.8]), they generate a closed normal subgroup scheme $\mathbf{N}_{\mathbb{Z}}^-$ of $\mathbf{L}_{\mathbb{Z}}$ that is a projective limit of extensions of \mathbb{G}_a over \mathbb{Z} . (The same is true and exhaust the whole $\mathbf{R}_{\mathbb{Z}}$ if $\mathbf{J} = \emptyset$.)

We examine the action of the imaginary PBW generators $\{\tilde{P}_{i,m\delta}\}_{i \in \mathbf{I} \setminus \mathbf{J}, m > 0}$ (weight of $\tilde{P}_{i,m\delta}$ is $m\delta$) in [5, (3.7)]. By applying them on the direct sum of P -weight λ -parts of $\mathbb{X}(\lambda)_{\mathbb{C}}$ for all $\lambda \in P_{\mathbf{J},+}$, we obtain a quotient group scheme $\mathbf{L}_{\mathbb{Z}} \rightarrow \mathbf{T}_{\mathbb{Z}}$ whose kernel is $\mathbf{N}_{\mathbb{Z}}^-$. By [5, Proposition 3.22], the group scheme $\text{Spec } \mathbb{Z}[\tilde{P}_{i,m\delta} \mid m > 1]^{\vee}$ for each $i \in \mathbf{I}$ is isomorphic to a projective limit of extensions of \mathbb{G}_a (given by truncations with respect to the duals of $\{\tilde{P}_{i,m\delta}\}_{m > N}$ for $N \in \mathbb{Z}_{>0}$), which is flat over \mathbb{Z} . Thus, so is $\mathbf{T}_{\mathbb{Z}}$.

From these, we deduce that $\mathbf{L}_{\mathbb{Z}} \subset \mathbf{I}_{\mathbb{Z}}^1$ defines a group subscheme over \mathbb{Z} such that $\mathbb{Z}[\mathbf{I}_{\mathbb{Z}}^1] \rightarrow \mathbb{Z}[\mathbf{L}]$ splits as \mathbb{Z} -modules. The P -weight decomposition of the coordinate ring defines a splitting $\mathbf{T}_{\mathbb{Z}} \subset \mathbf{L}_{\mathbb{Z}} \subset \mathbf{I}_{\mathbb{Z}}^1$. Since $[L(\mathbf{J}), L(\mathbf{J})]$ defines a product of connected and simply connected simple algebraic group, its (pro-unipotent part of the) Iwahori subgroup (we take the product over simple factors) defines a subgroup scheme $\mathbf{R}_{\mathbb{Z}}^0 \subset \mathbf{R}_{\mathbb{Z}}$ such that

$$\mathbf{R}_{\mathbb{Z}}^0 \cap \mathbf{T}_{\mathbb{Z}} = \text{Spec } \mathbb{Z} \quad (3.4)$$

(scheme-theoretically). Since $[L(\mathbf{J}), L(\mathbf{J})] \subset G$ defines a closed group subscheme over \mathbb{Z} (thanks to the fact that the corresponding modified enveloping algebras share the same \mathbb{Z} -basis by [41, 43, 67]), we deduce that the map $\mathbb{Z}[\mathbf{I}_{\mathbb{Z}}^1] \rightarrow \mathbb{Z}[\mathbf{R}_{\mathbb{Z}}^0]$ must be surjective. The one-parameter subgroups corresponding to the unipotent radical of $P(\mathbf{J})$ generates a subgroup $\mathbf{R}_{\mathbb{Z}}^+ \subset \mathbf{R}_{\mathbb{Z}} \subset \mathbf{I}_{\mathbb{Z}}^1$ such that $\mathbb{Z}[\mathbf{I}_{\mathbb{Z}}^1] \rightarrow \mathbb{Z}[\mathbf{R}_{\mathbb{Z}}^+]$ is surjective. In view of the P -weight decomposition, we deduce that $\mathbf{R}_{\mathbb{Z}}^0 \times \mathbf{R}_{\mathbb{Z}}^+ \cong \mathbf{R}_{\mathbb{Z}}$. From (3.4), we derive $\mathbb{Z}[\mathbf{L}] \cap \mathbb{Z}[\mathbf{R}] = \mathbb{Z}$. Therefore, we have a surjection

$$m_{\mathbb{Z}}^* : \mathbb{Z}[\mathbf{I}^1] \twoheadrightarrow \mathbb{Z}[\mathbf{L}] \otimes_{\mathbb{Z}} \mathbb{Z}[\mathbf{R}].$$

The ideal $\ker m_{\mathbb{Z}}^*$ must be flat over \mathbb{Z} , and hence $\ker m_{\mathbb{C}}^* \neq 0$ if $\ker m_{\mathbb{Z}}^* \neq 0$. We know that $m_{\mathbb{C}}^*$ must be an isomorphism by (3.3). This is a contradiction, and hence $m_{\mathbb{Z}}^*$ must be an isomorphism (cf. [31, Theorem 5.8]).

In view of the flatness of the group schemes over \mathbb{Z} afforded by the above, we conclude that

$$\mathbf{R}_{\mathbb{K}} \subset \text{Ad}(i\tilde{w}_0)([P(\mathbf{J}), P(\mathbf{J})][z]) \cap \mathbf{I}_{\mathbb{K}}^1$$

is in fact an isomorphism as schemes. By construction, $\mathbb{Z}[w]$ is precisely the subring of $\mathbb{Z}[\mathbf{I}_{\mathbb{K}}^1]$ that is invariant under the $\mathbf{R}_{\mathbb{Z}}$ -action. It follows that the image of the composition map

$$\mathbb{Z}[w] \hookrightarrow \mathbb{Z}[\mathbf{I}^1] \xrightarrow{\cong} \mathbb{Z}[\mathbf{L}] \otimes_{\mathbb{Z}} \mathbb{Z}[\mathbf{R}]$$

is equal to $\mathbb{Z}[\mathbf{L}] \otimes_{\mathbb{Z}} \mathbb{Z}$. Hence, we have

$$\mathrm{Spec} \mathbb{K}[w] = \mathbf{L}_{\mathbb{K}} \cong \mathbf{I}_{\mathbb{K}} / (H_{\mathbb{K}} \cdot \mathrm{Ad}(\dot{w}\dot{w}_0)([P(\mathbf{J}), P(\mathbf{J})][z]) \cap \mathbf{I}_{\mathbb{K}}).$$

Therefore, each $\mathbf{Q}_G(w)_{\mathbb{K}}$ contains $\mathrm{Spec} \mathbb{K}[w]$ as a Zariski open \mathbf{I} -orbit, and it admits a unique $(H \times \mathbb{G}_m)_{\mathbb{K}}$ -fixed point as required. \square

Corollary 3.12. *The ring $R(\mathbf{J})_{\mathbb{K}}$ is integral for each $\mathbf{J} \subset \mathbf{I}$.*

Remark 3.13. We refer Proposition A.1 for general case of Corollary 3.12.

Proof of Corollary 3.12. It suffices to prove the case $\mathbf{J} = \emptyset$ since a subring of an integral ring is integral. We borrow notation from the proof of Lemma 3.11. We have an inclusion

$$R_e = \bigoplus_{\lambda \in P_+} \mathbb{W}(\lambda)_{\mathbb{Z}}^{\vee} \hookrightarrow \mathbb{Z}[e] \otimes \bigoplus_{\lambda \in P_+} \mathbb{Z}\mathbf{v}_{w_0\lambda}^{\vee}$$

as the Rees construction of (3.1). Taking tensor product $\mathbb{K} \otimes_{\mathbb{Z}} \bullet$ preserves the inclusion in view of the definition of $\mathbb{Z}[e]$. Hence, $(R_e)_{\mathbb{K}}$ is integral as a subring of an integral ring. \square

Proposition 3.14. *Let $\mathbf{J} \subset \mathbf{I}$. The indscheme $(\mathbf{Q}_{G,\mathbf{J}}^{\mathrm{rat}})_{\mathbb{K}}$ admits a $G((z))$ -action. We have a subset $\mathbf{Q}'_{\mathbb{K}}$ of the set of \mathbb{K} -valued points of $(\mathbf{Q}_{G,\mathbf{J}}^{\mathrm{rat}})_{\mathbb{K}}$ that is in bijection with*

$$G((z)) / (H(\mathbb{K}) \cdot [P(\mathbf{J}), P(\mathbf{J})](z)).$$

Proof. In view of Lemma 3.11, the proof for general \mathbf{J} is completely parallel to the case of $\mathbf{J} = \emptyset$. Hence, we concentrate into the case $\mathbf{J} = \emptyset$ during this proof.

Let $w \in W_{\mathrm{af}}$ and $\beta \in Q^{\vee}$. We have an isomorphism $R_w \cong R_{wt_{\beta}}$ as rings with $U_{\mathbb{Z}}^+$ -action by Corollary 2.28.

In particular, we have an isomorphism $\mathbf{Q}_G(w)_{\mathbb{K}} \cong \mathbf{Q}_G(wt_{\beta})_{\mathbb{K}}$ of schemes with \mathbf{I} -actions for each $\beta \in Q^{\vee}$. This implies that $\mathbf{Q}_G(wt_{\beta})_{\mathbb{K}}$ has a Zariski open subset of the shape

$$\mathbf{I} / (H \cdot (\mathrm{Ad}(\dot{w}t_{\beta}\dot{w}_0)(N((z))) \cap \mathbf{I})) \cong \mathbf{I} / (H \cdot (\mathrm{Ad}(\dot{w}\dot{w}_0)(N((z))) \cap \mathbf{I})), \quad (3.5)$$

where we used that $\mathrm{Ad}(\dot{w}_0)(N((z)))$ is invariant under the $\mathrm{Ad}(\dot{t}_{\beta})$ -action.

The ring $(R_w)_{\mathbb{K}}$ admits an action of $SL(2, i)$ whenever $s_i w \leq_{\frac{\infty}{2}} w$ ($i \in \mathbf{I}_{\mathrm{af}}$) since each $\mathbb{W}_{ww_0}(\lambda)_{\mathbb{K}}$ ($\lambda \in P_+$) admits an action of $SL(2, i)$ by Corollary 2.16. Hence, $\mathrm{Proj} (R_w)_{\mathbb{K}}$ admits an action of $SL(2, i)$ if $s_i w \leq_{\frac{\infty}{2}} w$. In particular, the pro-algebraic group $\mathbf{I}(i)$ acts on $\mathrm{Proj} (R_w)_{\mathbb{K}}$ if $s_i w \leq_{\frac{\infty}{2}} w$. Hence, the ind-limit

$$(\mathbf{Q}_G^{\mathrm{rat}})_{\mathbb{K}} = \bigcup_{w \in W_{\mathrm{af}}} \mathbf{Q}_G(w)_{\mathbb{K}} = \varinjlim_w \mathrm{Proj} (R_w)_{\mathbb{K}}$$

admits an action of $\mathbf{I}(i)$ for each $i \in \mathbf{I}_{\mathrm{af}}$ (that coincide on \mathbf{I}). By rank two calculations, they induce an action of $N_{G((z))}(H(\mathbb{K}))$ on $(\mathbf{Q}_G^{\mathrm{rat}})_{\mathbb{K}}$. The intersections of $N_{G((z))}(H(\mathbb{K}))$ and $\mathbf{I}(\mathbb{K})$ or $\mathbf{I}(i)(\mathbb{K})$ ($i \in \mathbf{I}_{\mathrm{af}}$) inside $G((z))$ define common actions on $(\mathbf{Q}_G^{\mathrm{rat}})_{\mathbb{K}}$. The system of groups $(\mathbf{I}(\mathbb{K}), N_{G((z))}(H(\mathbb{K})), \mathbf{I}(i)(\mathbb{K}); i \in \mathbf{I}_{\mathrm{af}})$ (in the sense of [57, Definition 5.1.6]) admits a map from the system of groups in [57, §6.1.16] for $\tilde{\mathfrak{g}}$ (with exponential maps replaced by one-parameter subgroups). Hence, [57, Theorem 6.1.17] asserts that $G((z))$ acts on $(\mathbf{Q}_G^{\mathrm{rat}})_{\mathbb{K}}$.

The Bruhat decomposition of $SL(2, i)$ asserts that $\mathbb{O}(w)_{\mathbb{K}} \sqcup \mathbb{O}(s_i w)_{\mathbb{K}}$ admits a $SL(2, i)$ -action. This induces an action of $SL(2, i)(\mathbb{K})$ ($i \in \mathbf{I}_{\text{af}}$) on the union

$$\mathbf{Q}'_{\mathbb{K}} := \bigsqcup_{w \in W_{\text{af}}} \mathbb{O}(w)_{\mathbb{K}}(\mathbb{K}) \subset \mathbf{Q}_G^{\text{rat}}(\mathbb{K}).$$

Taking into account the fact that each $\mathbb{O}(w)_{\mathbb{K}}$ admits an \mathbf{I} -action (Lemma 3.11), we conclude that $\mathbf{Q}'_{\mathbb{K}}$ admits an action of $G((z))$ thanks to the Iwasawa decomposition (cf. [38, Theorem 2.5])

$$G((z)) = \bigsqcup_{w \in W_{\text{af}}} \mathbf{I}(\mathbb{K}) \dot{w} \dot{w}_0 H(\mathbb{K}) \cdot N((z))$$

as required. \square

For $\lambda, \mu \in P_+$, we have a unique injective $\dot{U}_{\mathbb{Z}}^0$ -module map

$$V(\lambda + \mu)_{\mathbb{Z}} \longrightarrow V(\lambda)_{\mathbb{Z}} \otimes_{\mathbb{Z}} V(\mu)_{\mathbb{Z}}$$

obtained by sending $\mathbf{v}_{\lambda+\mu}^0$ to $\mathbf{v}_{\lambda}^0 \otimes \mathbf{v}_{\mu}^0$, that is in fact a \mathbb{Z} -direct summand (Corollary 2.12). By extending the scalar, we obtain a unique injective $(\dot{U}_{\mathbb{K}}^0, \mathbb{K}[[z]])$ -bimodule map

$$\eta_{\lambda, \mu} : V(\lambda + \mu)_{\mathbb{K}} \otimes \mathbb{K}[[z]] \longrightarrow (V(\lambda)_{\mathbb{K}} \otimes \mathbb{K}[[z]]) \otimes_{\mathbb{K}[[z]]} (V(\mu)_{\mathbb{K}} \otimes \mathbb{K}[[z]]).$$

Lemma 3.15. *For each $\lambda, \mu, \gamma \in P_+$, we have*

$$\eta_{\lambda+\mu, \gamma} \circ (\text{id} \boxtimes \eta_{\lambda, \mu}) = \eta_{\lambda, \mu+\gamma} \circ (\text{id} \boxtimes \eta_{\mu, \gamma}).$$

Proof. Straight-forward from the construction (cf. Lemme 2.27). \square

Proposition 3.16. *Assume that $\text{char } \mathbb{K} \neq 2$. For each $w \in W_{\text{af}}$ and $\mathbf{J} \subset \mathbf{I}$, we have an \mathbf{I} -equivariant rational map*

$$\psi_w : \mathbf{Q}_{G, \mathbf{J}}(w)_{\mathbb{K}} \dashrightarrow \bigcup_{m \in \mathbb{Z}} \prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}_{\mathbb{K}}(V(\varpi_i)_{\mathbb{K}} \otimes z^m \mathbb{K}[[z]]) = \prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}_{\mathbb{K}}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}((z))),$$

that gives rise to a $G((z))$ -equivariant rational map

$$\psi : (\mathbf{Q}_{G, \mathbf{J}}^{\text{rat}})_{\mathbb{K}} \dashrightarrow \prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}_{\mathbb{K}}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}((z))).$$

In addition, the set $(\text{Im } \psi)(\mathbb{K})$ defines a closed (ind-)subscheme of $\prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}_{\mathbb{K}}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}((z)))$.

Proof. We have a surjective map

$$\mathbb{W}(\varpi_i)_{\mathbb{K}} \rightarrow V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[z]$$

as $\dot{U}_{\mathbb{K}}^{\geq 0}$ -modules since we have the corresponding map over $\dot{U}_{\mathbb{Z}}^{\geq 0}$ such that the P -weight ϖ_i -part is the same (and $V(\varpi_i)_{\mathbb{K}}$ is cyclic as a $\dot{U}_{\mathbb{K}}^0$ -module). The identification of the P -weight ϖ_i -part also implies that this map commutes with the action of τ_{β} ($\beta \in \mathbf{Q}^{\vee}$), and extends to a surjective map

$$\mathbb{X}(\varpi_i)_{\mathbb{K}} \rightarrow V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[z, z^{-1}]$$

of $\dot{U}_{\mathbb{K}}$ -modules.

This gives a rational map

$$\mathbb{P}(\mathbb{W}(\varpi_i)_{\mathbb{K}}) \dashrightarrow \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[z])$$

and its graded completion

$$\mathbb{P}(\mathbb{W}(\varpi_i)_{\mathbb{K}}^{\wedge}) \dashrightarrow \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z]]).$$

Taking Corollary 2.26 into account, we have an embedding:

$$\mathbf{Q}_{G,J}(e)_{\mathbb{K}} \hookrightarrow \prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}(\mathbb{W}(\varpi_i)_{\mathbb{K}}^{\wedge}). \quad (3.6)$$

This yields a rational map

$$\psi_e : \mathbf{Q}_{G,J}(e)_{\mathbb{K}} \dashrightarrow \prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z]])$$

as a composition. This map is $G[[z]]$ -equivariant by construction.

For each $w \in W$, we can choose $\beta \in Q_+^{\vee}$ such that $\mathbf{Q}_{G,J}(w)_{\mathbb{K}} \cong \mathbf{Q}_{G,J}(wt_{\beta})_{\mathbb{K}} \subset \mathbf{Q}_{G,J}(e)_{\mathbb{K}}$ by Corollary 2.28 and Lemma 2.29. Hence we obtain the map ψ_w for every $w \in W_{\text{af}}$ as the composition of the above maps (up to grading shifts). We fix the effect of grading shifts from ψ_w to $\psi_{wt_{\beta}}$ by applying $\tau_{w_0\beta}$ on $\mathbb{W}(\varpi_i)_{\mathbb{K}}^{\wedge}$ for each $i \in \mathbf{I} \setminus \mathbf{J}$ in (3.6). Then, the $\tau_{w_0\beta}$ -action (on the completions of $\{\mathbb{X}(\varpi_i)_{\mathbb{K}}\}_{i \in \mathbf{I} \setminus \mathbf{J}}$) transfers ψ_w to $\psi_{wt_{\beta}}$ for each $\beta \in Q^{\vee}$, and it is compatible with the restrictions to $\mathbf{Q}_G(\bullet)$'s by Lemma 2.17 1). Hence, we obtain the map ψ of indschemes. This map is $G((z))$ -equivariant in our sense. This proves the first assertion.

From now on, we concentrate into the second assertion.

Each $(H \times \mathbb{G}_m)$ -fixed point of $\mathbb{O}(w)_{\mathbb{K}}$ ($w \in W_{\text{af}}$) is contained in the domain of ψ , and their images are distinct by inspection. It follows that $\mathbf{Q}'_{\mathbb{K}}$ is contained in the domain of ψ , and the restriction of ψ to $\mathbf{Q}'_{\mathbb{K}}$ is injective by examining the stabilizer of the $\mathbf{I}(\mathbb{K})$ -actions at each $(H \times \mathbb{G}_m)$ -fixed points.

In view of Theorem 2.18, the maps $\{\eta_{\lambda,\mu}\}_{\lambda,\mu}$ ($\lambda, \mu \in P_{J,+}$) induce a commutative diagram of $\dot{U}_{\mathbb{K}}^{\geq 0}$ -modules:

$$\begin{array}{ccc} \mathbb{W}(\lambda)_{\mathbb{K}} \hookrightarrow & \bigotimes_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{W}(\varpi_i)_{\mathbb{K}}^{\otimes \langle \alpha_i^{\vee}, \lambda \rangle} & \\ \downarrow \kappa_{\lambda} & & \searrow \\ V(\lambda)_{\mathbb{K}} \otimes \mathbb{K}[z] \hookrightarrow & \left(\bigotimes_{i \in \mathbf{I} \setminus \mathbf{J}} V(\varpi_i)_{\mathbb{K}}^{\otimes \langle \alpha_i^{\vee}, \lambda \rangle} \right) \otimes \mathbb{K}[z] & \longleftarrow \bigotimes_{i \in \mathbf{I} \setminus \mathbf{J}} \left(V(\varpi_i)_{\mathbb{K}}^{\otimes \langle \alpha_i^{\vee}, \lambda \rangle} \otimes \mathbb{K}[z] \right). \end{array} \quad (3.7)$$

Here the map κ_{λ} is well-defined by examining the degree 0-part and the action of $E_0 = F_{\vartheta} \otimes z$ (where F_{ϑ} is a non-zero vector in the $-\vartheta$ -weight space of \mathfrak{n}^-). The above commutative diagram also commutes with the translation by τ_{β} ($\beta \in Q^{\vee}$) by construction. Moreover, we have $\kappa_{\lambda}(\mathbf{v}_{w\lambda}) \neq 0$ for each $w \in W_{\text{af}}$. Therefore, the map κ_{λ} must be surjective whenever its d -degree belongs to $\sum_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{Z} \langle \alpha_i^{\vee}, \lambda \rangle$.

For each $i \in \mathbf{I} \setminus \mathbf{J}$, the $\mathbb{Z}[\frac{1}{2}]$ -integral structure of $V(2\varpi_i) \otimes_{\mathbb{C}} \mathbb{C}[z, z^{-1}]$ at the odd degree and even degree must be the same as $\dot{U}_{\mathbb{Z}}$ -modules (as we can connect the extremal weight vectors of even degree part and the odd degree part using the $\mathfrak{sl}(2)$ -strings of length 3). Therefore, for $\lambda = \varpi_i, \varpi_i + \varpi_j, 2\varpi_i$ ($i, j \in \mathbf{I} \setminus \mathbf{J}$), the map κ_{λ} is surjective.

Consider a (representative of the) image

$$\psi(x) = (x_i) \in \prod_{i \in \mathbf{I} \setminus \mathbf{J}} V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}((z))$$

of a \mathbb{K} -valued point $x \in \mathbf{Q}_{G, \mathbf{J}}(e)_{\mathbb{K}}$ under ψ . We consider its lifts $\tilde{x}_i \in \mathbb{W}(\varpi_i)_{\mathbb{K}}^{\wedge}$ and $\tilde{x}_j \in \mathbb{W}(\varpi_j)_{\mathbb{K}}^{\wedge}$. They must belong to

$$\text{Im}(\mathbb{W}(\varpi_i + \varpi_j)_{\mathbb{K}}^{\wedge} \rightarrow (\mathbb{W}(\varpi_i)_{\mathbb{K}} \otimes \mathbb{W}(\varpi_j)_{\mathbb{K}})^{\wedge})$$

in order to satisfy the defining relations of $R_{\mathbb{K}}$. In view of the commutative diagram (3.7) for $\lambda = \varpi_i + \varpi_j$, we deduce an equation

$$x_i \otimes_{\mathbb{K}[[z]]} x_j \in \text{Im} \eta_{\varpi_i, \varpi_j} \quad i, j \in \mathbf{I} \setminus \mathbf{J}. \quad (3.8)$$

Since the relation of the ring $\bigoplus_{\lambda \in P_{\mathbf{I}, +}} V(\lambda)_{\mathbb{K}}^*$ is generated by P -degrees $2\varpi_i$ and $\varpi_i + \varpi_j$ for $i, j \in \mathbf{I}$ ([15, Theorem 3.5.3]), this defines an element of

$$G((z))/[P(\mathbf{J}), P(\mathbf{J})]((z)) \subset \overline{G/[P(\mathbf{J}), P(\mathbf{J})]}(\mathbb{K}((z))) \quad (3.9)$$

through quadratic relations (see [29, §4]), where

$$\overline{G/[P(\mathbf{J}), P(\mathbf{J})]} = \text{Spec } \mathbb{K}[G/[P(\mathbf{J}), P(\mathbf{J})]]$$

is the basic affine space. Therefore, applying some τ_{β} ($\beta \in Q^{\vee}$) if necessary, we conclude that if a \mathbb{K} -valued point x of $\mathbf{Q}_{G, \mathbf{J}}^{\text{rat}}$ belongs to the domain of ψ , then $\psi(x)$ belongs to the set of \mathbb{K} -valued points of the image given by (3.9). Taking Proposition 3.14 (and its $G((z))$ -action) and the equation (3.8) into account, we conclude the second assertion. \square

Corollary 3.17. *Keep the setting of Proposition 3.16. The map ψ induces a bijection of \mathbb{K} -valued points between the domain and the range.*

Proof. By abuse of convention, we regard $\mathbf{Q}'_{\mathbb{K}}$ as subsets of the both of $\mathbf{Q}_G^{\text{rat}}(\mathbb{K})$ and $(\text{Im } \psi)(\mathbb{K})$. Let $x' \in \mathbf{Q}_G^{\text{rat}}(\mathbb{K})$ be a preimage of a $(H \times \mathbb{G}_m)(\mathbb{K})$ -fixed point $x \in \mathbf{Q}'_{\mathbb{K}}$ through ψ . Consider the embedding

$$(\mathbf{Q}_{G, \mathbf{J}}^{\text{rat}})_{\mathbb{K}} \hookrightarrow \prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}(\mathbb{X}(\varpi_i)_{\mathbb{K}}^{\wedge})$$

that prolongs (3.6). Let $x' = (x'_i)_{i \in \mathbf{I} \setminus \mathbf{J}}$ (resp. $x = (x_i)_{i \in \mathbf{I} \setminus \mathbf{J}}$) be the coordinate of x' (resp. x) through the above embedding. We can regard $x'_i \in \mathbb{X}(\varpi_i)_{\mathbb{K}}^{\wedge}$, and it admits a decomposition

$$x'_i = \prod_{\mu} x'_i[\mu],$$

where $\mu \in P$ runs over the P -weights of $\mathbb{X}(\varpi_i)_{\mathbb{K}}^{\wedge}$ (or $\mathbb{W}(\varpi_i)_{\mathbb{K}}^{\wedge}$). Let $d'_i[\mu]$ be the degree of the lowest d -degree non-zero contribution of $x'_i[\mu]$ (or ∞ if $x'_i[\mu] = 0$)

for each $i \in \mathbf{I} \setminus \mathbf{J}$ and $\mu \in P$. Let d_i be the d -degree of x_i for each $i \in \mathbf{I} \setminus \mathbf{J}$ (remember that x is $(H \times \mathbb{G}_m)(\mathbb{K})$ -fixed). Note that we have $x_i[u\varpi_i] = x'_i[u\varpi_i]$ for every $i \in \mathbf{I}$ and $u \in W$ since

$$\mathbb{W}(\varpi_i)_{\mathbb{K}} \quad \text{and} \quad V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[z] \quad \text{shares the same } P\text{-weight } u\varpi_i\text{-parts.} \quad (3.10)$$

For each $\gamma \in Q^\vee$, we have a collection of the automorphisms of the vector spaces $\{\mathbb{X}(\lambda)_{\mathbb{K}}\}_{\lambda \in P_+}$ by shifting the (d) -gradings of the weight μ -parts by $\langle \gamma, \mu \rangle$ ($\mu \in P$). This defines an automorphism of $R_{\mathbb{K}}^+$, and hence defines an automorphism of $\mathbf{Q}_G^{\text{rat}}(\mathbb{K})$. By using this twist for an appropriate $\gamma \in Q^\vee$, we can assume that

$$d'_i[\mu] \gg d'_i[u\varpi_i] = d_i \quad \mu \neq u\varpi_i \quad (3.11)$$

for each $i \in \mathbf{I} \setminus \mathbf{J}$ and fixed $u \in W$ from the fact that every P -weight of $W(\varpi_i)_{\mathbb{K}}$ is $\leq \varpi_i$. We have $x_i \in \mathbb{W}_{ww_0}(\varpi_i)_{\mathbb{K}}$ for some $w \in W_{\text{af}}$ and all $i \in \mathbf{I} \setminus \mathbf{J}$. The inequality (3.11) implies $x'_i \in \mathbb{W}_{ww_0}(\varpi_i)_{\mathbb{K}}^\wedge$ for all $i \in \mathbf{I} \setminus \mathbf{J}$ ([44, Theorem 5.17]). Hence, we have $x' \in \mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{K}}$ for $w \in W_{\text{af}}$ such that x is the unique $(H \times \mathbb{G}_m)$ -fixed point of $\mathbb{O}(\mathbf{J}, w)_{\mathbb{K}}$.

From (3.10), the $(H \times \mathbb{G}_m)$ -action shrinks x' to x . In view of its proof, the Zariski open set $\mathbb{O}(\mathbf{J}, w)_{\mathbb{K}}$ in the Lemma 3.11 is obtained by localizing along the vectors $\{\mathbf{v}_{ww_0\lambda}^\vee\}_{\lambda \in P_{+,+}}$, that pairs nontrivially with x' . Therefore, we have necessarily $x' \in \mathbb{O}(\mathbf{J}, w)_{\mathbb{K}}(\mathbb{K})$. As $\mathbb{O}(\mathbf{J}, w)_{\mathbb{K}}(\mathbb{K}) \subset \mathbf{Q}'_{\mathbb{K}}$ by Proposition 3.14 and its proof, we deduce $x = x'$. Since $\mathbf{Q}'_{\mathbb{K}}$ is stable under the above action of Q^\vee , as well as $G((z))$, we conclude that $x = x'$ holds for a preimage of x' of every $x \in \mathbf{Q}'_{\mathbb{K}}$ as required. \square

Theorem 3.18. *Assume that $\text{char } \mathbb{K} \neq 2$. For each $\mathbf{J} \subset \mathbf{I}$, we have a closed immersion of ind-schemes*

$$(\mathbf{Q}_{G,\mathbf{J}}^{\text{rat}})_{\mathbb{K}} \longrightarrow \prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}((z))).$$

In particular, the set of \mathbb{K} -valued points of the ind-scheme $(\mathbf{Q}_{G,\mathbf{J}}^{\text{rat}})_{\mathbb{K}}$ is in bijection with

$$G((z)) / (H(\mathbb{K}) \cdot [P(\mathbf{J}), P(\mathbf{J})]((z))).$$

Proof. By construction, the locus $E \subset \mathbf{Q}_{G,\mathbf{J}}^{\text{rat}}$ on which ψ (borrowed from Proposition 3.16) is not defined is an ind-subscheme. The map ψ is $G((z))$ -equivariant. It follows that E admits a $G((z))$ -action as ind-schemes. The map (borrowed from the proof of Proposition 3.16)

$$\mathbf{Q}_{G,\mathbf{J}}(e)_{\mathbb{K}} \longrightarrow \mathbb{P}(\mathbb{W}(\varpi_i)_{\mathbb{K}}^\wedge)$$

sends an irreducible component of $(E \cap \mathbf{Q}_{G,\mathbf{J}}(e)_{\mathbb{K}})$ onto a closed subscheme of $\mathbb{P}(\ker(\mathbb{W}(\varpi_i)_{\mathbb{K}}^\wedge \rightarrow V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z]])$ for some $i \in \mathbf{I} \setminus \mathbf{J}$. In particular, $(E \cap \mathbf{Q}_{G,\mathbf{J}}(e)_{\mathbb{K}}) \subset \mathbf{Q}_{G,\mathbf{J}}(e)_{\mathbb{K}}$ is a closed subscheme. Taking into account the $\{\tau_\beta\}_{\beta}$ -actions (Lemma 2.17), we have $E \neq \emptyset$ if and only if $(E \cap \mathbf{Q}_{G,\mathbf{J}}(e)_{\mathbb{K}}) \neq \emptyset$, and ψ is a closed immersion if and only if ψ_e is a closed immersion (cf. Lemma 2.29).

The one-parameter subgroup $a = (\xi, 1) : \mathbb{G}_m \rightarrow (H \times \mathbb{G}_m)$ for $\xi \in Q_{<}^\vee$ attracts every point of $\mathbb{P}(\mathbb{W}(\varpi_i)_{\mathbb{K}})$ into its $(H \times \mathbb{G}_m)$ -fixed points (by setting $t \rightarrow 0$). In particular, E has a $(H \times \mathbb{G}_m)$ -fixed point that is not realized by $\mathbf{Q}'_{\mathbb{K}}$. Since the $(H \times \mathbb{G}_m)$ -fixed point of bounded d -degree is captured by the corresponding

degree terms of $R_{\mathbb{K}}$, it follows that the indscheme E intersects with $\mathcal{Q}'_J(v, e)$ for some $v \in W_{\text{af}}$. The intersection of $\mathcal{Q}'_J(v, e)_{\mathbb{K}}$ and $\mathbf{Q}'_{\mathbb{K}}$ (as the set of \mathbb{K} -valued points) defines a closed subset Y of (a product of) finite-dimensional projective space by Proposition 3.16 and Corollary 3.17. In particular, Y acquires the structure of a proper scheme through ψ . From this view-point, Corollary 3.17 provides a ψ -section of Y that defines a bijection of \mathbb{K} -valued points with a Zariski open subset $\mathcal{Q}'_J(v, e)_{\mathbb{K}} \setminus E$ of $\mathcal{Q}'_J(v, e)_{\mathbb{K}}$. Since $\mathbf{Q}'_{\mathbb{K}}$ admits a homogeneous $G((z))$ -action, we can think of ψ as an everywhere defined section of a vector bundle over the image (whose fiber is a product of $\ker(\mathbb{W}(\varpi_i)_{\mathbb{K}}^{\wedge} \rightarrow V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z]])$'s). It implies that $\mathcal{Q}'_J(v, e)_{\mathbb{K}} \setminus E$ can be seen as an everywhere defined section of a vector bundle over the image (whose fiber is a product of finite-dimensional subspaces of $\ker(\mathbb{W}(\varpi_i)_{\mathbb{K}} \rightarrow V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z]])$'s). Since $\mathcal{Q}'_J(v, e)_{\mathbb{K}}$ is a finite type scheme defined over an algebraically closed field \mathbb{K} , it has a dense subset formed by its \mathbb{K} -valued points ([36, Corollarie 10.4.8]). Thus, the section ψ can be seen as that of schemes. Therefore, $\mathcal{Q}'_J(v, e)_{\mathbb{K}} \setminus E$ is proper by itself. In conclusion, $\mathcal{Q}'_J(v, e)_{\mathbb{K}} \setminus E$ can be seen as a connected component of $\mathcal{Q}'_J(v, e)_{\mathbb{K}}$ (as being open and closed subset) which does not intersect with E . It follows that if $E \neq \emptyset$ as a scheme, then we find that the scheme $\mathcal{Q}'_J(v, e)$ must have at least two connected components. Therefore, the projective coordinate ring $R_e^v(\mathbf{J})_{\mathbb{K}}$ of $\mathcal{Q}'_J(v, e)$ must be non-integral. The same is true if we replace v with a smaller element with respect to \ll_{∞} .

Hence, we can find $f, g \in R_e^v(\mathbf{J})_{\mathbb{K}} \setminus \{0\}$ such that $fg = 0$ for $v \ll_{\infty} e$. Since E and its complement are \mathbb{G}_m -stable, we can assume that f and g are \mathbb{G}_m -eigenfunctions. Since E defines a connected component of $\mathcal{Q}'_J(v, e)$ for every $v \ll_{\infty} e$, we can fix the degrees of f and g for every $v \ll_{\infty} e$. For a fixed degree, the d -graded component of $R_e^v(\mathbf{J})_{\mathbb{K}}$ and $R_e(\mathbf{J})_{\mathbb{K}}$ are in common for $v \ll_{\infty} e$. Since $R_e^v(\mathbf{J})_{\mathbb{K}}$ is a quotient ring of $R_e(\mathbf{J})_{\mathbb{K}}$, we can find \mathbb{G}_m -eigenfunctions $f, g \in R_e(\mathbf{J})_{\mathbb{K}} \setminus \{0\}$ such that $fg = 0$. This implies that the ring $R_{\mathbb{K}}$ is also non-integral. This contradicts with Corollary 3.12, and hence we deduce $E = \emptyset$ as an ind-scheme. Therefore, we conclude that ψ is in fact a genuine morphism (instead of a rational map) of ind-schemes.

Next, we prove that ψ (or rather ψ_e) defines a closed immersion. By Lemma 3.11 and the fact that ψ is a morphism, we deduce that ψ induces an isomorphism between the function fields of $\mathbf{Q}_{G, \mathbf{J}}(e)_{\mathbb{K}}$ and its image under ψ . An irreducible component Z of a reduced scheme $\mathcal{Q}'_J(v, e)_{\mathbb{K}}$ shares a Zariski dense subset with a unique orbit $\mathbb{O}(\mathbf{J}, w)_{\mathbb{K}}$ ($w \in W_{\text{af}}$). Hence, the variety $(Z \cap \mathbb{O}(\mathbf{J}, w)_{\mathbb{K}})$ and its image under ψ are presented by the quotients of a common polynomial ring $\mathbb{K}[\mathbb{O}(\mathbf{J}, w)_{\mathbb{K}}]$ of infinite variables. In view of the d -degree bounds from the above and below offered by $\mathbb{W}_{vw_0}^-(\varpi_i)_{\mathbb{K}}$ and $\mathbb{W}_{w_0}(\varpi_i)_{\mathbb{K}}$ for each $i \in \mathbf{I} \setminus \mathbf{J}$, these two quotients of $\mathbb{K}[\mathbb{O}(\mathbf{J}, w)_{\mathbb{K}}]$ factor through a common polynomial ring of finite variables. Thus, we deduce that ψ also induces an isomorphism between the function fields of Z and $\psi(Z)$. Once we fix an integer d_0 , two rings $R_e^v(\mathbf{J})_{\mathbb{K}}$ and $R_e(\mathbf{J})_{\mathbb{K}}$ share the same (d) -grading d_0 -part for $v \ll_{\infty} e$. Hence, it suffices to prove that ψ restricts to a closed immersion of $\mathcal{Q}'_J(v, e)_{\mathbb{K}}$ for $v \ll_{\infty} e$. Here $R_e^v(\mathbf{J})_{\mathbb{K}}$ is weakly normal by Lemma 3.7. Let R' be the multi-homogeneous coordinate ring of the (reduced induced structure of the) closed subscheme of $\prod_{i \in \mathbf{I} \setminus \mathbf{J}} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}((z)))$ defined by $\psi(\mathcal{Q}'_J(v, e)_{\mathbb{K}})$. Then, if we consider via each maximal integral quotient, we deduce that the weak normalization of R' is precisely $R_e^v(\mathbf{J})_{\mathbb{K}}$ (up to irrelevant locus). This implies that $\psi(\mathcal{Q}'_J(v, e)_{\mathbb{K}})$ defines

a closed subscheme for each $v \in W_{\text{af}}$ as required.

Since ψ is a closed immersion, the rest of assertion follows from Proposition 3.16. These complete the proof. \square

Corollary 3.19 (of the proof of Theorem 3.18). *Let $w \in W_{\text{af}}$ and $\mathbf{J} \subset \mathbf{I}$. Every two rational functions on $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{K}}$ are distinguished by a pair of \mathbb{K} -valued points of $\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}$ for some $v \in W_{\text{af}}$. In particular, the union $\bigcup_{v \in W_{\text{af}}} \mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}$ is Zariski dense in $\mathbf{Q}_{G,\mathbf{J}}(w)_{\mathbb{K}}$.* \square

Corollary 3.20. *The conclusions of Theorem 1.4 and Theorem 1.5 holds when we replace \mathbb{C} with an algebraically closed field \mathbb{K} of characteristic $\neq 2$.*

Proof. In view of Theorem 3.18, all the results in Theorem 1.4 and Theorem 1.5 are consequences of the corresponding set-theoretic considerations. \square

Corollary 3.21. *We have $\mathcal{Q}'(v, w)_{\mathbb{K}} \neq \emptyset$ if and only if $v \leq_{\frac{\infty}{2}} w$.*

Proof. By Lemma 3.5, it remains to prove that $\mathcal{Q}'(v, w)_{\mathbb{K}} = \emptyset$ if $v \not\leq_{\frac{\infty}{2}} w$. By Theorem 3.18 and the definition of $\mathcal{Q}'(v, w)_{\mathbb{K}}$, the $(H \times \mathbb{G}_m)$ -fixed point of $\mathcal{O}(u)_{\mathbb{K}}$ is contained in $\mathcal{Q}'(v, w)_{\mathbb{K}}$ ($v, w \in W_{\text{af}}$) only if $v \leq_{\frac{\infty}{2}} u \leq_{\frac{\infty}{2}} w$. Since $\leq_{\frac{\infty}{2}}$ is a partial order, the condition $v \not\leq_{\frac{\infty}{2}} w$ implies that $\mathcal{Q}'(v, w)_{\mathbb{K}}$ has no $(H \times \mathbb{G}_m)$ -fixed point. Since $\mathcal{Q}'(v, w)_{\mathbb{K}}$ is projective by Lemma 3.4, it carries a $(H \times \mathbb{G}_m)$ -fixed point if it is nonempty. Thus, we conclude $\mathcal{Q}'(v, w)_{\mathbb{K}} = \emptyset$ as required. \square

3.3 Coarse representability of the scheme $\mathbf{Q}_G^{\text{rat}}$

Material in this subsection is rather special throughout this paper, and is irrelevant to the arguments in the later part, such as the normality of quasi-map spaces.

In this subsection, we assume that $\text{char } \mathbb{K} \neq 2$, and we also drop subscripts \mathbb{K} from $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{K}}$ and its subschemes in order to simplify notation.

Let $\text{Aff}_{\mathbb{K}}$ be the category of affine schemes over \mathbb{K} . We identify $\text{Aff}_{\mathbb{K}}^{\text{op}}$ with the category of commutative rings over \mathbb{K} . Let $\text{Zar}_{\mathbb{K}}$ denote a big Zariski site over \mathbb{K} [77, Section 020N]. For $X \in \text{Zar}_{\mathbb{K}}$, the assignment

$$\text{Zar}_{\mathbb{K}}^{\text{op}} \ni U \mapsto \text{Hom}_{\text{Zar}_{\mathbb{K}}}(U, X) \in \text{Sets}$$

defines a sheaf h_X on $\text{Zar}_{\mathbb{K}}$ [77, Definition 00WR].

For the definition on the coarse moduli functors, we refer to [78, Definition 1.10]. However, we employ some modified definition given in the below:

Definition 3.22 (Strict indscheme). Let $\mathfrak{X} = \bigcup_{n \geq 0} X_n$ be an increasing union of schemes in $\text{Zar}_{\mathbb{K}}$. We call $(\mathfrak{X}, \{X_n\}_n)$ (or simply refer as \mathfrak{X}) a strict indscheme if each inclusion $X_k \subset X_{k+1}$ ($k \geq 0$) is a closed immersion.

Definition 3.23 (Filtered sheaf on $\text{Zar}_{\mathbb{K}}$). A filtered (pre)sheaf $(\mathcal{F}, \{\mathcal{F}_n\}_{n \geq 0})$ on $\text{Zar}_{\mathbb{K}}$ is a family of (pre)sheaves such that $\mathcal{F}_k \subset \mathcal{F}_{k+1}$ for each $k \in \mathbb{Z}_{\geq 0}$ and $\mathcal{F} = \bigcup_n \mathcal{F}_n$. Let $(\mathcal{F}, \{\mathcal{F}_n\}_{n \geq 0})$ and $(\mathcal{G}, \{\mathcal{G}_n\}_{n \geq 0})$ be filtered (pre)sheaves on $\text{Zar}_{\mathbb{K}}$. A morphism $f : \mathcal{F} \rightarrow \mathcal{G}$ of (pre)sheaves is said to be continuous if for each $n \in \mathbb{Z}_{\geq 0}$, there is some $m \in \mathbb{Z}_{\geq 0}$ such that

$$f(\mathcal{F}_n) \subset \mathcal{G}_m \quad \text{and} \quad \mathcal{G}_n \cap \text{Im } f \subset f(\mathcal{F}_m).$$

Let $(\mathfrak{F}, \{F_n\}_n)$ be a strict indscheme. Then, we call $h_{\mathfrak{F}} := (\bigcup_n h_{F_n}, \{h_{F_n}\}_{n \geq 0})$ the filtered sheaf associated to \mathfrak{F} .

Definition 3.24 (Coarse ind-representability). Let \mathcal{X} be a filtered (pre)sheaf on $Zar_{\mathbb{K}}$. Let \mathfrak{X} be a strict indscheme over \mathbb{K} . We say that \mathcal{X} is coarsely ind-representable by \mathfrak{X} if the following conditions hold:

- We have a continuous morphism $u : \mathcal{X} \rightarrow h_{\mathfrak{X}}$ of filtered (pre)sheaves;
- We have $\mathcal{X}(\mathbb{k}) = h_{\mathfrak{X}}(\mathbb{k})$ for an overfield $\mathbb{k} \supset \mathbb{K}$;
- Let \mathfrak{Y} be a strict indscheme and we have a continuous morphism $f : \mathcal{X} \rightarrow h_{\mathfrak{Y}}$, then it factors as:

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{u} & h_{\mathfrak{X}} \\ & \searrow f & \downarrow g \\ & & h_{\mathfrak{Y}} \end{array}$$

where g is a morphism of presheaves. It is automatic that g is continuous, and hence is induced by a morphism of indschemes.

We consider the assignment \mathcal{Q} on $Aff_{\mathbb{K}}^{op}$ defined as:

$$Aff_{\mathbb{K}}^{op} \ni R \mapsto \mathcal{Q}(R) := G(R((z)))/(H(R)N(R((z)))) \in \text{Sets}.$$

For each $n \in \mathbb{Z}_{\geq 0}$, we consider an assignment

$$Aff_{\mathbb{K}}^{op} \ni R \mapsto \mathcal{Q}_n(R) := \{g \bmod H(R)N(R((z))) \in \mathcal{Q}(R) \mid (\star)\} \in \text{Sets},$$

where

- (\star) $g_{\mathbf{v}_{\varpi_i}}$ has at worst pole of order n on $V(\varpi_i)_{\mathbb{Z}} \otimes_{\mathbb{Z}} R((z))$ for each $i \in \mathbf{I}$.

The assignments $(\mathcal{Q}, \{\mathcal{Q}_n\}_n)$ define a filtered presheaf on $Zar_{\mathbb{K}}$ that we denote by \mathcal{Q} .

Lemma 3.25. *The indscheme $\mathbf{Q}_G^{\text{rat}}$ defines a filtered sheaf on $Zar_{\mathbb{K}}$ given by a strict indscheme structure.*

Proof. A scheme over \mathbb{K} defines a sheaf over $Zar_{\mathbb{K}}$, and so is its increasing union. In view of Proposition 3.16, the pole order n condition amounts to choose the \mathbf{I} -orbits $\mathcal{O}(ut_{\beta})$ ($u \in W, \beta \in Q^{\vee}$) such that $\langle \beta, \varpi_i \rangle \geq -n$ (for every $i \in \mathbf{I}$), that makes the smaller one to be a closed subscheme of the larger one (cf. Lemma 2.29). \square

Proposition 3.26. *The scheme $\mathbf{Q}_G^{\text{rat}}$ coarsely ind-represents the presheaf \mathcal{Q} .*

Proof. We first construct an injective continuous morphism $\mathcal{Q} \rightarrow h_{\mathbf{Q}_G^{\text{rat}}}$ as filtered presheaves on $Zar_{\mathbb{K}}$ such that $\mathcal{Q}(\mathbb{k}) = h_{\mathbf{Q}_G^{\text{rat}}}(\mathbb{k})$ for an overfield $\mathbb{k} \supset \mathbb{K}$.

For $R \in Aff_{\mathbb{K}}^{op}$, the set $\mathcal{Q}(R)$ is represented by a class of $g \in G(R((z)))$ modulo the right action of $H(R)N(R((z)))$. It defines a point of $\mathbf{Q}_G^{\text{rat}}(R)$ by applying g on $\{\mathbf{v}_{\varpi_i}\}_{i \in \mathbf{I}} \in \prod_{i \in \mathbf{I}} \mathbb{P}_R(V(\varpi_i)_{\mathbb{Z}} \otimes_{\mathbb{Z}} R((z)))$. Since the $G(R((z)))$ -stabilizer of \mathbf{v}_{ϖ_i} is precisely $H(R)N(R((z)))$, we conclude an inclusion $\mathcal{Q}(R) \subset h_{\mathbf{Q}_G^{\text{rat}}}(R)$. By examining the construction, we deduce that this defines an injective continuous morphism of filtered presheaves.

By the Bruhat decomposition, we have

$$G(\mathbb{k}((z)))/N(\mathbb{k}((z))) = (G/N)(\mathbb{k}((z)))$$

for an overfield $\mathbb{k} \supset \mathbb{K}$ (and hence $\mathbb{k}((z))$ is a field). In view of Theorem 3.18, we conclude that $\mathcal{Q} \subset h_{\mathbf{Q}_G^{\text{rat}}}$ is an inclusion of filtered presheaves with $\mathcal{Q}(\mathbb{k}) = h_{\mathbf{Q}_G^{\text{rat}}}(\mathbb{k})$ when $\mathbb{k} \supset \mathbb{K}$ is an overfield.

We verify the versality property. Suppose that we have a strict indscheme $(\mathcal{X}, \{X_n\}_n)$ and we have a continuous morphism $\mathcal{Q} \rightarrow h_{\mathcal{X}}$. By Lemma 3.11, we deduce that each \mathbf{I} -orbit of $\mathbf{Q}_G^{\text{rat}}$ defines a subfunctor of \mathcal{Q}_n for some $n \geq 0$. The zero-th ind-piece $(\mathbf{Q}_G^{\text{rat}})_0$ in Lemma 3.25 is $\mathbf{Q}_G(e)$.

For each $t_\beta \leq_{\frac{\infty}{2}} e$ ($\beta \in Q_+^V$), we find a reduced expression $t_\beta^{-1} = s_{i_1} \cdots s_{i_\ell}$ (that we record as $\mathbf{i} := (i_1, \dots, i_\ell)$) and form a scheme

$$Z(\mathbf{i})^\circ := \mathbf{I}(i_1) \times^{\mathbf{I}} \mathbf{I}(i_2) \times^{\mathbf{I}} \cdots \times^{\mathbf{I}} \mathbb{O}(t_\beta)$$

and the map

$$Z(\mathbf{i})^\circ = \mathbf{I}(i_1) \times^{\mathbf{I}} \mathbf{I}(i_2) \times^{\mathbf{I}} \cdots \times^{\mathbf{I}} \mathbb{O}(t_\beta) \rightarrow \mathbf{Q}_G(e)$$

(see e.g. Kumar [57, Chapter VIII]; cf. [46, §6]). In view of Lemma 3.11, the image of this map contains an open neighbourhood of $\mathbb{O}(t_\beta)$. In view of the $G((z))$ -action (or the various $SL(2, i)$ -actions for $i \in \mathbf{I}_{\text{af}}$) on $\mathbf{Q}_G^{\text{rat}}$ and \mathcal{Q} , we have a morphism

$$f_{\mathbf{i}} : h_{Z(\mathbf{i})^\circ} \longrightarrow \mathcal{Q}_0.$$

By varying \mathbf{i} (and consequently varying $t_\beta \leq_{\frac{\infty}{2}} e$), we deduce that the union of the image of the morphisms $\{f_{\mathbf{i}}\}_{\mathbf{i}}$ exhausts $\mathcal{Q}_0(\mathbb{k})$ for an overfield $\mathbb{k} \supset \mathbb{K}$. From the Yoneda embedding, we derive a map

$$Z(\mathbf{i})^\circ \longrightarrow X_n$$

of schemes for some fixed $n \in \mathbb{Z}$. This map factors through a scheme Z that glues (among \mathbf{i} 's) all the closed points that maps to the same points in \mathcal{Q}_0 . Such a scheme is integral as $Z(\mathbf{i})^\circ$'s are so and the gluing identifies the Zariski open dense subset $\mathbb{O}(e)$ for distinct \mathbf{i} 's. In addition, we have a birational map $\pi : Z \rightarrow \mathbf{Q}_G(e)$, and hence we have

$$Z(\mathbb{k}) = \mathbf{Q}_G(e)(\mathbb{k}) = \mathcal{Q}_0(\mathbb{k}) \quad \text{for an overfield } \mathbb{k} \supset \mathbb{K}.$$

We prove that $Z = \mathbf{Q}_G(e)$ by induction. For each $m \in \mathbb{Z}_{\geq 0}$, let $\mathbf{Q}_G(e)_{< m}$ (resp. $\mathbf{Q}_G(e)_{\leq m}$) be the union of \mathbf{I} -orbits in $\mathbf{Q}_G(e)$ of the shape $\mathbb{O}(v)$ for $\ell_{\frac{\infty}{2}}(v) < m$ (resp. $\leq m$).

Assume that the map π is an isomorphism when restricted to $\mathbf{Q}_G(e)_{< m}$, and we prove the same is true when restricted to $\mathbf{Q}_G(e)_{\leq m}$. The $m = 1$ case is afforded by $\mathbb{O}(e) \subset Z$ already used in the construction of the above.

We have a partial compactification $Z(\mathbf{i})$ of $Z(\mathbf{i})^\circ$ with a map $f_{\mathbf{i}}^+$ given as:

$$Z(\mathbf{i}) := \mathbf{I}(i_1) \times^{\mathbf{I}} \mathbf{I}(i_2) \times^{\mathbf{I}} \cdots \times^{\mathbf{I}} \mathbf{Q}_G(t_\beta) \xrightarrow{f_{\mathbf{i}}^+} \mathbf{Q}_G(e).$$

Note that we have a surjective morphism induced by $\mathbb{O}(t_\beta) \rightarrow \text{Spec } \mathbb{K}$

$$\eta_{\mathbf{i}} : Z(\mathbf{i}) \longrightarrow \mathbf{I}(i_1) \times^{\mathbf{I}} \mathbf{I}(i_2) \times^{\mathbf{I}} \cdots \times^{\mathbf{I}} \text{Spec } \mathbb{K},$$

where we denote the image (the RHS term) by $Z'(\mathbf{i})$. Since $Z'(\mathbf{i})$ is a finite successive \mathbb{P}^1 -fibration, it is proper. The map $f_{\mathbf{i}}^+$ is proper as the product map

$$(\eta_{\mathbf{i}} \times f_{\mathbf{i}}^+) : Z(\mathbf{i}) \hookrightarrow Z'(\mathbf{i}) \times \mathbf{I}(i_1) \cdots \mathbf{I}(i_\ell) \mathbf{Q}_G(t_\beta) = Z'(\mathbf{i}) \times \mathbf{Q}_G(e)$$

is a closed immersion. In view of the isomorphism $\mathbf{Q}_G(t_\beta) \cong \mathbf{Q}_G(e)$, we transplant $\mathbf{Q}_G(e)_{< m}$ to $\mathbf{Q}_G(t_\beta)_{< m}$.

Claim A. For each closed point $x \in \mathbf{Q}_G(e)_{\leq m}$, the scheme

$$(f_{\mathbf{i}}^+)^{-1}(x) \setminus ((f_{\mathbf{i}}^+)^{-1}(x) \cap (\mathbf{I}(i_1) \times^{\mathbf{I}} \mathbf{I}(i_2) \times^{\mathbf{I}} \cdots \times^{\mathbf{I}} \mathbf{Q}_G(t_{\beta})_{< m})) \subset (f_{\mathbf{i}}^+)^{-1}(x)$$

is a closed subscheme that is zero-dimensional. In other words, it is a finite union of points (that is potentially an empty set).

Proof. For each sequence $(j_1, \dots, j_s) \in \mathbf{I}_{\text{af}}^s$ ($s \in \mathbb{Z}_{>0}$) and $v \leq_{\frac{\infty}{2}} e$ such that $\ell_{\frac{\infty}{2}}(v) = \ell$, the image of the map

$$f : \mathbf{I}(j_1) \times^{\mathbf{I}} \mathbf{I}(j_2) \times^{\mathbf{I}} \cdots \times^{\mathbf{I}} \mathbf{I}(j_s) \times^{\mathbf{I}} \mathbf{Q}_G(v) \longrightarrow \mathbf{Q}_G^{\text{rat}}$$

induced by the multiplication is a union of \mathbf{I} -orbits $\mathbb{O}(v')$ with $\ell_{\frac{\infty}{2}}(v') \geq \ell - s$ (as we have $\ell_{\frac{\infty}{2}}(s_i w) \in \{\ell_{\frac{\infty}{2}}(w) \pm 1\}$ for each $i \in \mathbf{I}_{\text{af}}$ and $w \in W_{\text{af}}$ by [74, Lecture 13, Proposition ℓ_s]). In addition, if the image of the map f contains $\mathbb{O}(v')$ for $\ell_{\frac{\infty}{2}}(v') = \ell - s$, then the map f is an isomorphism along $\mathbb{O}(v')$ (as the isomorphism between open subsets). By collecting these for \mathbf{I} -orbits in the closed subset $\mathbf{Q}_G(t_{\beta}) \setminus \mathbf{Q}_G(t_{\beta})_{< m}$ of $\mathbf{Q}_G(t_{\beta})$ in the construction of the (proper) map $f_{\mathbf{i}}^+$, we conclude the result. \square

We return to the proof of Proposition 3.26. By Claim A, we deduce that

$$\overline{(f_{\mathbf{i}}^+)^{-1}(x) \cap (\mathbf{I}(i_1) \times^{\mathbf{I}} \mathbf{I}(i_2) \times^{\mathbf{I}} \cdots \times^{\mathbf{I}} \mathbf{Q}_G(t_{\beta})_{< m})} \subset (f_{\mathbf{i}}^+)^{-1}(x)$$

is a union of connected components of $(f_{\mathbf{i}}^+)^{-1}(x)$ for each closed point $x \in \mathbf{Q}_G(e)_{\leq m}$.

Requiring that regular functions on $(f_{\mathbf{i}}^+)^{-1}(\mathbf{Q}_G(e)_{\leq m})$ to be constant along all the fibers yield sections in $(f_{\mathbf{i}}^+)_* \mathcal{O}_{Z(\mathbf{i})}$. From this (for arbitrary $t_{\beta} \leq_{\frac{\infty}{2}} e$ and \mathbf{i}) and the induction hypothesis, we conclude that $\pi^{-1}(\mathbf{Q}_G(e)_{\leq m}) \subset Z$ is a union of proper schemes over $\mathbf{Q}_G(e)_{\leq m}$ that contains a Zariski open subset $\mathbf{Q}_G(e)_{< m}$. In view of Corollary 2.39 (when $\text{char } \mathbb{K} > 0$) or Proposition A.1 (when $\text{char } \mathbb{K} = 0$; cf. [51, Theorem A]), we deduce that

$$(Z \supset) \quad \pi^{-1}(\mathbf{Q}_G(e)_{\leq m}) \rightarrow \mathbf{Q}_G(e)_{\leq m}$$

defines an isomorphism as schemes (as $\pi^{-1}(\mathbf{Q}_G(e)_{\leq m}) \rightarrow \mathbf{Q}_G(e)_{\leq m}$ is finite bijective, and birational, cf. [77, Section 02LQ]). Therefore, induction on m proceeds and we conclude $Z \cong \mathbf{Q}_G(e)$ as schemes. Thus, we obtain a morphism $\mathbf{Q}_G(e) \rightarrow X_n$ of schemes.

By rearranging $\mathbf{Q}_G(e)$ by the right Q^{\vee} -translations, we deduce a morphism $\mathbf{Q}_G^{\text{rat}} \rightarrow \mathfrak{X}$ as indschemes. This yields a continuous morphism $h_{\mathbf{Q}_G^{\text{rat}}} \rightarrow h_{\mathfrak{X}}$. Therefore, $h_{\mathbf{Q}_G^{\text{rat}}}$ is an initial object in the category of sheaves on $\text{Zar}_{\mathbb{K}}$ ind-representable by strict indschemes that admits a continuous morphism from \mathcal{Q} as required. \square

Corollary 3.27. For each $\mathbf{J} \subset \mathbf{I}$, the scheme $\mathbf{Q}_{G, \mathbf{J}}^{\text{rat}}$ coarsely ind-represents the filtered presheaf $\mathcal{Q}_{\mathbf{J}}$ defined by

$$\text{Aff}_{\mathbb{K}}^{\text{op}} \ni R \mapsto \mathcal{Q}_{\mathbf{J}}(R) := G(R((z))) / (H(R) \cdot [P_{\mathbf{J}}, P_{\mathbf{J}}](R((z)))) \in \text{Sets}.$$

Proof. By construction, we have a continuous morphism of presheaves $\mathcal{Q} \rightarrow \mathcal{Q}_{\mathbf{J}}$ (by transplanting subsheaves \mathcal{Q}_n to $\mathcal{Q}_{\mathbf{J}}$ via this map). Thus, the coarse ind-representability of $\mathbf{Q}_G^{\text{rat}}$ implies that the maximal indscheme X obtained by

gluing points of $\mathbf{Q}_G^{\text{rat}}$ admits a continuous morphism $\mathcal{Q}_J \rightarrow h_X$ coarsely ind-represents the filtered presheaf \mathcal{Q}_J . Every two rational functions on $\mathbf{Q}_G(w)$ ($w \in W_{\text{af}}$) are distinguished by some pair of \mathbb{K} -valued points (Corollary 3.19). Since we have $\mathcal{Q}_J(\mathbb{K}) = \mathbf{Q}_{G,J}^{\text{rat}}(\mathbb{K})$, we conclude that $X = \mathbf{Q}_{G,J}^{\text{rat}}$. \square

3.4 The properties of the schemes $\mathcal{Q}'_J(v, w)$

In the rest of this section, we assume $\text{char } \mathbb{K} \neq 2$.

Lemma 3.28. *Let $J \subset I$. For each $\beta \in Q_+^\vee$, the set of \mathbb{K} -valued points of $\mathcal{Q}'_J(\beta, e)_{\mathbb{K}}$ is in bijection with the collection $\{u_\lambda(z)\}_{\lambda \in P_{J,+}}$ such that*

- We have $u_\lambda(z) \in V(\lambda)_{\mathbb{K}} \otimes \bigoplus_{j=0}^{-\langle w_0 \beta, \lambda \rangle} \mathbb{K} z^j$;
- For each $\lambda, \mu \in P_+$, we have $\eta_{\lambda, \mu}(u_\lambda(z) \otimes u_\mu(z)) = u_{\lambda+\mu}(z)$.

Proof. We have

$$\mathbf{Q}_{G,J}(e)_{\mathbb{K}} = (\mathbf{Q}_{G,J}^{\text{rat}})_{\mathbb{K}} \cap \prod_{i \in I \setminus J} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z]]) \subset \prod_{i \in I \setminus J} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}((z)))$$

by Theorem 3.18. By the symmetry of the construction of $\mathcal{Q}'_J(\beta, e)$ in terms of θ (cf. Remark 2.31), we conclude that

$$\mathcal{Q}'_J(\beta, e)_{\mathbb{K}} = (\mathbf{Q}_{G,J}^{\text{rat}})_{\mathbb{K}} \cap \prod_{i \in I \setminus J} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z]]) \cap \prod_{i \in I \setminus J} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z^{-1}]] z^{-\langle \beta, w_0 \varpi_i \rangle})$$

inside $\prod_{i \in I \setminus J} \mathbb{P}(V(\varpi_i)_{\mathbb{K}} \otimes \mathbb{K}[[z, z^{-1}]])$, that is our degree bound. In view of this, it suffices to remember that the second condition is the same as the Plücker relation that defines $G((z))/H(\mathbb{K}) \cdot [P(J), P(J)]((z))$ in the last two paragraphs of the proof of Proposition 3.16. \square

Let $w, v \in W_{\text{af}}$ and $J \subset I$. For each $\lambda \in P_{J,+}$, we have a line bundle $\mathcal{O}_{\mathcal{Q}'_J(v, w)}(\lambda)$ on $\mathcal{Q}'_J(v, w)$ such that we have a map

$$R_w^v(J, \lambda) \rightarrow \Gamma(\mathcal{Q}'_J(v, w), \mathcal{O}_{\mathcal{Q}'_J(v, w)}(\lambda))$$

that commutes with the multiplications. This yields $\mathcal{O}_{\mathcal{Q}'_J(v, w)}(\lambda)$ for each $\lambda \in P_J$ by tensor products.

Lemma 3.29. *For each $w, v \in W_{\text{af}}$, the line bundle $\mathcal{O}_{\mathcal{Q}'_J(v, w)}(\lambda)$ is very ample if $\langle \alpha_i^\vee, \lambda \rangle > 0$ for every $i \in I \setminus J$.*

Proof. We can assume that $\mathcal{Q}'_J(v, w) \neq \emptyset$ without the loss of generality.

By Lemma 3.1, we have

$$\mathcal{Q}'_J(v, w) \hookrightarrow \prod_{i \in I \setminus J} \mathbb{P}_{\mathbb{Z}}(R_w^v(\varpi_i)^\vee).$$

From this and again by Lemma 3.1, we deduce that the following diagram is commutative:

$$\begin{array}{ccc} \prod_{i \in I \setminus J} \mathbb{P}_{\mathbb{Z}}(R_w^v(J, \varpi_i)^\vee) & \xrightarrow{\xi} & \mathbb{P}_{\mathbb{Z}}(\bigotimes_{i \in I \setminus J} R_w^v(J, \varpi_i)^\vee) \xleftarrow{\kappa} \mathbb{P}_{\mathbb{Z}}(R_w^v(J, \rho_J)^\vee) \\ & \searrow & \nearrow \\ & \mathcal{Q}'_J(v, w) & \end{array}$$

where the ξ is the Veronese embedding and κ is induced from the multiplication map. The ring $R_w^v(\mathbf{J})$ is reduced by Corollary 3.10. Since $\langle \alpha_i^\vee, \rho_{\mathbf{J}} \rangle > 0$ for every $i \in \mathbf{I} \setminus \mathbf{J}$ and the multiplication maps are surjective, a non-zero element of $R_w^v(\mathbf{J}, \rho_{\mathbf{J}})$ is not supported in the irrelevant locus. It follows that $R_w^v(\mathbf{J}, \rho_{\mathbf{J}}) \subset H^0(\mathcal{Q}'_{\mathbf{J}}(v, w), \mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)}(\rho_{\mathbf{J}}))$ and it is spanned by a product of linear functions that separates closed points of $\mathcal{Q}'_{\mathbf{J}}(v, w)$. In particular, $\mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)}(\rho_{\mathbf{J}})$ is very ample. Since we have embeddings

$$R_w^v(\mathbf{J}, \rho_{\mathbf{J}}) \subset R_w^v(\mathbf{J}, \lambda) \subset H^0(\mathcal{Q}'_{\mathbf{J}}(v, w), \mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)}(\lambda))$$

obtained through multiplications corresponding to the duals of extremal weight vectors (so that the image is base-point-free along a $(H \times \mathbb{G}_m)$ -stable Zariski open neighbourhood of each $(H \times \mathbb{G}_m)$ -fixed point of a projective variety), we conclude that $\mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)}(\lambda)$ is also very ample as required. \square

Theorem 3.30. *For each $w = ut_{\beta'} \in W_{\text{af}}$ ($u \in W, \beta' \in Q^\vee$) and $\beta \in Q_+^\vee$, we have an isomorphism $\mathcal{Q}'(\beta, w)_{\mathbb{C}} \cong \mathcal{Q}(\beta - \beta', u)$ as varieties. Moreover, $\mathcal{Q}'(\beta, w)_{\mathbb{K}}$ is irreducible and its dimension is given as*

$$\dim \mathcal{Q}'(\beta, w)_{\mathbb{K}} = 2 \langle \beta - \beta', \rho \rangle + \dim \mathcal{B}(u).$$

Proof. By Lemma 3.7, we know that $(R_v^\beta)_{\mathbb{K}}$ is a quotient of $(R_w^\beta)_{\mathbb{K}}$ for $v \in W_{\text{af}}$ such that $v \leq_{\infty} w$. Hence, we have $\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbf{Q}_G(v)_{\mathbb{K}} = \mathcal{Q}'(\beta, v)_{\mathbb{K}}$. By Lemma 3.6, the scheme $\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbb{O}(v)_{\mathbb{K}}$ is isomorphic to $\mathcal{Q}'(\beta - \gamma, u')_{\mathbb{K}} \cap \mathbb{O}(u')_{\mathbb{K}}$, where $v = u't_\gamma$ ($u' \in W, \gamma \in Q_+^\vee$). By Lemma 3.28 and Theorem 1.8, we have:

- $(\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbb{O}(v)_{\mathbb{K}}) = (\mathcal{Q}'(\beta, v)_{\mathbb{K}} \cap \mathbb{O}(v)_{\mathbb{K}})$ for each $v \in W_{\text{af}}$ such that $v \leq_{\infty} w$;
- The variety $(\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbb{O}(v)_{\mathbb{K}})$ is irreducible for every $v \in W_{\text{af}}$.

We have an equality

$$\dim(\mathcal{Q}'(\beta, w)_{\mathbb{C}} \cap \mathbb{O}(v)_{\mathbb{C}}) = \dim \mathcal{Q}(\beta - \gamma, u') = 2 \langle \beta - \gamma, \rho \rangle + \dim \mathcal{B}(u')$$

for $v = u't_\gamma$ ($u' \in W, \gamma \in Q_+^\vee$) by Lemma 3.28 and Theorem 1.7 (whenever $\mathcal{Q}'(\beta, w)_{\mathbb{C}} \cap \mathbb{O}(v)_{\mathbb{C}} \neq \emptyset$). In addition, Lemma 3.3 implies

- We have $\dim \mathcal{Q}'(\beta, w)_{\mathbb{C}} = \dim \mathcal{Q}'(\beta, w)_{\mathbb{K}}$.

In particular, we have the desired dimension formula if $\mathcal{Q}'(\beta, w)_{\mathbb{K}}$ is irreducible with its Zariski open dense subset $(\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbb{O}(w)_{\mathbb{K}})$.

Since $\mathcal{Q}(\beta, w)$ and $\mathcal{Q}'(\beta, w)_{\mathbb{C}}$ shares a same open subset and the former is irreducible, we have $\mathcal{Q}(\beta, w) = \mathcal{Q}'(\beta, w)_{\mathbb{C}}$ as closed subvarieties of $(\mathbf{Q}_G^{\text{rat}})_{\mathbb{C}}$ if $\mathcal{Q}'(\beta, w)_{\mathbb{C}}$ is irreducible.

Therefore, it suffices to prove that $\mathcal{Q}'(\beta, w)_{\mathbb{K}}$ is irreducible (with its Zariski open dense subset $(\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbb{O}(w)_{\mathbb{K}})$). By Theorem 1.1 and Corollary 3.21, it suffices to prove that

$$(\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbb{O}(sw)_{\mathbb{K}}) \subset \overline{(\mathcal{Q}'(\beta, w)_{\mathbb{K}} \cap \mathbb{O}(w)_{\mathbb{K}})} \quad (3.12)$$

for every $w \in W_{\text{af}}$ and every reflection $s \in W_{\text{af}}$ such that $\ell^{\frac{\infty}{2}}(sw) = \ell^{\frac{\infty}{2}}(w) + 1$ and $w_0 t_\beta \leq_{\infty} sw \leq_{\infty} w$. Here $\mathbf{Q}_G(sw)_{\mathbb{K}} \subset \mathbf{Q}_G(w)_{\mathbb{K}}$ is an irreducible component

of the boundary by Theorem 1.4 3) (cf. Corollary 3.20). This boundary component is these cut out as (a part of) the zero of $\mathbf{v}_{vw_0\lambda}^\vee \in \mathbb{W}_{vw_0}(\lambda)^\vee$ ($\lambda \in P_+$). Thus, we deduce that $\overline{(\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(sw)_\mathbb{K})}$ contains an irreducible component of

$$\overline{(\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(w)_\mathbb{K})} \cap \{f = 0\}$$

for some single equation f (an instance of $\mathbf{v}_{vw_0\lambda}^\vee$'s) if it is nonempty. By the comparison of dimensions, this forces

$$\overline{(\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(w)_\mathbb{K})} \cap \{f = 0\} \cap \mathbb{O}(sw)_\mathbb{K} \subset (\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(sw)_\mathbb{K}) \quad (3.13)$$

to be an irreducible component if the LHS is nonempty. Consider the $(H \times \mathbb{G}_m)$ -invariant curve C that connects the (unique) $(H \times \mathbb{G}_m)$ -fixed points p_{sw} in $\mathbb{O}(sw)_\mathbb{K}$ and p_w in $\mathbb{O}(w)_\mathbb{K}$. The curve C is the closure of the orbit of an one-parameter unipotent subgroup action corresponding to a root in $\Delta_{\text{af},+}$ applied to p_w . In particular, we have $C \subset \mathbb{O}(sw)_\mathbb{K} \sqcup \mathbb{O}(w)_\mathbb{K}$. By the degree bound (from the above) offered by Lemma 3.28, we conclude that $C \subset \overline{(\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(w)_\mathbb{K})}$. Therefore, the LHS of (3.13) is nonempty. Hence, the irreducibility of $(\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(sw)_\mathbb{K})$ forces

$$\overline{(\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(w)_\mathbb{K})} \cap \{f = 0\} \cap \mathbb{O}(sw)_\mathbb{K} = (\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(sw)_\mathbb{K}).$$

Therefore, we conclude (3.12). This implies $\mathcal{Q}(\beta - \beta', u) = \mathcal{Q}'(\beta, w)_\mathbb{C}$ as an irreducible (reduced) closed subvariety of $(\mathbf{Q}_G^{\text{rat}})_\mathbb{C}$, and $\mathcal{Q}'(\beta, w)_\mathbb{K}$ is irreducible in general. Its dimension $\dim \mathcal{Q}(\beta - \beta', u)$ comes from the dimension of $(\mathcal{Q}'(\beta, w)_\mathbb{K} \cap \mathbb{O}(w)_\mathbb{K})$, that is a Zariski open dense subset of $\mathcal{Q}'(\beta, w)_\mathbb{K}$. \square

Corollary 3.31. *For each $w, v \in W_{\text{af}}$, the dimension of an irreducible component of $\mathcal{Q}'(v, w)_\mathbb{K}$ is $\ell^{\frac{\infty}{2}}(v) - \ell^{\frac{\infty}{2}}(w)$ if $\mathcal{Q}'(v, w)_\mathbb{K} \neq \emptyset$ (that is equivalent to $v \leq_{\frac{\infty}{2}} w$ by Corollary 3.21). In particular, $\mathcal{Q}'(v, w)_\mathbb{K}$ is equidimensional.*

Proof. The case $v = w_0 t_\beta$ with $\beta \in Q^\vee$ follows from Theorem 3.30. We have $\mathcal{Q}'(w, w)_\mathbb{K} = \text{Spec } \mathbb{K}$ ($w \in W_{\text{af}}$) since $\dim \mathbb{W}_{ww_0}(\lambda)_\mathbb{K} \cap \mathbb{W}_{ww_0}^-(\lambda)_\mathbb{K} = 1$ for every $\lambda \in P_+$ by the P^{af} -weight comparison.

Let $v = ut_\beta$ ($u \in W$). In view of the proof of Theorem 3.30, the variety $\mathcal{Q}'(v, w)_\mathbb{K}$ is obtained from $\mathcal{Q}'(w_0 t_\beta, w)_\mathbb{K}$ by a $(\ell(w_0) - \ell(u))$ -successive hyperplane cuts by the θ -twists of the \mathbf{I} -stable boundaries of $\mathbf{Q}_G(\bullet)_\mathbb{K}$ (cf. Remark 2.31). Each of these hyperplane cuts lowers the dimension of an irreducible component by at most one if the intersection is nonempty. Since $\ell(w_0) - \ell(u) = \ell^{\frac{\infty}{2}}(w_0 t_\beta) - \ell^{\frac{\infty}{2}}(v)$, the dimension inequality \geq always hold.

Every irreducible component of $\mathcal{Q}'(v, w)_\mathbb{K}$ is $(H \times \mathbb{G}_m)$ -stable by construction. Since we have only one $(H \times \mathbb{G}_m)$ -fixed point in each \mathbf{I} -orbit of $(\mathbf{Q}_G^{\text{rat}})_\mathbb{K}$, all but one $(H \times \mathbb{G}_m)$ -fixed point of $\mathbf{Q}_G(w)_\mathbb{K}$ ($w \in W_{\text{af}}$) lies in the boundary. Thus, if an irreducible component of $\mathcal{Q}'(v, w)_\mathbb{K}$ does not meet any \mathbf{I} -stable boundary divisors of $\mathbf{Q}_G(w)_\mathbb{K}$ and the θ -twists of any \mathbf{I} -stable boundary divisors of $\mathbf{Q}_G(vw_0)_\mathbb{K}$, then such an irreducible component does not contain a $(H \times \mathbb{G}_m)$ -fixed point unless $v = w$. This is a contradiction (to the properness of $\mathcal{Q}'(v, w)_\mathbb{K}$). Hence, every irreducible component of $\mathcal{Q}'(v, w)_\mathbb{K}$ meets its boundary cut out by an \mathbf{I} -stable boundary divisor of $\mathbf{Q}_G(w)_\mathbb{K}$ or the θ -twist of an \mathbf{I} -stable boundary divisor of $\mathbf{Q}_G(vw_0)_\mathbb{K}$.

Assume to the contrary to deduce contradiction. Then, we have $\mathcal{Q}'(v, w)_\mathbb{K}$ whose irreducible components have dimension $(\ell^{\frac{\infty}{2}}(v) - \ell^{\frac{\infty}{2}}(w))$, but one of its

irreducible component is in fact contained in $\mathcal{Q}'(v', w')_{\mathbb{K}}$ with $\mathcal{Q}'(v', w')_{\mathbb{K}} \subsetneq \mathcal{Q}'(v, w)_{\mathbb{K}}$ for $v', w' \in W_{\text{af}}$, and hence it gives an irreducible component of $\mathcal{Q}'(v', w')_{\mathbb{K}}$ with its dimension $> (\ell^{\frac{\infty}{2}}(v') - \ell^{\frac{\infty}{2}}(w'))$. From this, we can lower the dimension of an irreducible component of $\mathcal{Q}'(v', w')_{\mathbb{K}}$ by intersecting with a divisor to raise v' or lower w' successively to reach to the case $v' = w'$ (as in the second paragraph). Since dimension drops at most one in each step, we find that $\dim \mathcal{Q}'(w, w)_{\mathbb{K}} > 0$ for some $w \in W_{\text{af}}$. This is a contradiction, and hence this case does not happen.

These imply that the dimension equality always hold as required. \square

Remark 3.32. The analogous assertions for Theorem 3.30 and Corollary 3.31 holds for the case $\mathbf{J} \subsetneq \mathbf{I}$ (cf. Corollaries 4.21 and 4.22).

Theorem 3.33. *For each $w, v \in W_{\text{af}}$, $\mathbf{J} \subset \mathbf{I}$, and $\lambda \in P_{\mathbf{J},+}$, we have*

$$H^i(\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}}(\lambda)) \cong \begin{cases} R_w^v(\lambda)_{\mathbb{K}} & (i = 0, \lambda \in P_{\mathbf{J},++}) \\ \{0\} & (i \neq 0) \end{cases}.$$

Moreover, if $w', v' \in W_{\text{af}}$ satisfies $\mathcal{Q}'_{\mathbf{J}}(v', w') \subset \mathcal{Q}'_{\mathbf{J}}(v, w)$ and $\lambda \in P_{\mathbf{J},+}$, then the restriction map induces a surjection

$$H^0(\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}}(\lambda)) \twoheadrightarrow H^0(\mathcal{Q}'_{\mathbf{J}}(v', w')_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v', w')_{\mathbb{K}}}(\lambda)).$$

Remark 3.34. The strict dominance condition in Theorem 3.33 cannot be removed naively. For $G = SL(3)$, $\mathbb{K} = \mathbb{C}$, and $w = e$, we have

$$\dim \mathcal{Q}'_{\{1\}}(2\alpha_1^{\vee}, e)_{\mathbb{C}} = 8 > 7 = \dim \mathcal{Q}'(2\alpha_1^{\vee}, e)_{\mathbb{C}}.$$

This results in the non-injectivity of the pullback map

$$H^0(\mathcal{Q}'_{\{1\}}(2\alpha_1^{\vee}, e)_{\mathbb{C}}, \mathcal{O}_{\mathcal{Q}'_{\{1\}}(2\alpha_1^{\vee}, e)_{\mathbb{C}}}(3\varpi_1)) \twoheadrightarrow H^0(\mathcal{Q}'(2\alpha_1^{\vee}, e)_{\mathbb{C}}, \mathcal{O}_{\mathcal{Q}'(2\alpha_1^{\vee}, e)_{\mathbb{C}}}(3\varpi_1)),$$

where the LHS is $R_e^{2\alpha_1^{\vee}}(3\varpi_1)_{\mathbb{C}} = R_e^{2\alpha_1^{\vee}}(\{1\}, 3\varpi_1)_{\mathbb{C}} \cong S^3(\mathbb{C}^3 \oplus \mathbb{C}^3 z \oplus \mathbb{C}^3 z^2)^{\vee}$, while the RHS is its quotient by the G -invariants of $(\mathbb{C}^3 \otimes \mathbb{C}^3 z \otimes \mathbb{C}^3 z^2)^{\vee} \subset S^3(\mathbb{C}^3 \oplus \mathbb{C}^3 z \oplus \mathbb{C}^3 z^2)^{\vee}$. In fact, the ring $(R_e^{2\alpha_1^{\vee}})_{\mathbb{C}}$ is reduced but not integral, and the extra irreducible component is contained in the irrelevant locus.

Proof of Theorem 3.33. We first observe that it is enough to show the surjectivity part of the assertion only for the $w = w'$ case since the $v = v'$ case follows by applying θ (and the rest of the cases follow by a repeated applications of the two cases).

Since $\mathbb{O}(\mathbf{J}, w)_{\mathbb{F}_p} \subset \mathbf{Q}_{G, \mathbf{J}}(w)_{\mathbb{F}_p}$ is affine, the union Z of codimension one \mathbf{I} -orbits contains the support of an ample divisor. The reduced union of the irreducible components of Z affords a Weil divisor D (by Corollary A.2). Since our Frobenius splitting of $\mathbf{Q}_{G, \mathbf{J}}(w)_{\mathbb{F}_p}$ is \mathbf{I} -canonical and it is compatible with the \mathbf{I} -orbit closures (Corollary 2.38), we deduce that $\mathbf{Q}_{G, \mathbf{J}}(w)_{\mathbb{F}_p}$ is D -split in the sense of [14, Definition 3] (cf. [75, Definition 1.2]). Each irreducible component of Z intersects properly with $\mathcal{Q}'_{\mathbf{J}}(v', w)_{\mathbb{F}_p} \subset \mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{F}_p}$ ($v' \in W_{\text{af}}$) whenever the intersection is nonempty by Corollary 3.31 (and Remark 3.32). Thus, we deduce that $\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{F}_p}$ admits a Frobenius D -splitting compatible with $\mathcal{Q}'_{\mathbf{J}}(v', w)_{\mathbb{F}_p}$ ($v' \in W_{\text{af}}$). Therefore, the cohomology vanishing part of the assertion and the

surjectivity part of the assertion (for $w' = w$) follow from [75, Proposition 1.13 (ii)]. The cohomology vanishing part of the assertion lifts to $\text{char } \mathbb{K} = 0$ by [15, Proposition 1.6.2]. The surjectivity part of the assertion lifts to $\text{char } \mathbb{K} = 0$ by [15, Corollary 1.6.3] and Lemma 3.3.

It remains to calculate $H^0(\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}}(\lambda))$ for each $\lambda \in P_{\mathbb{J}, ++}$. Here $H^0(\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}}(\lambda))$ is obtained as the degree λ -part of the (graded) normalization of the ring $(R_w^v(\mathbb{J}))_{\mathbb{K}}$. For each prime p , our ring $(R_w^v(\mathbb{J}))_{\mathbb{F}_p}$ is weakly normal by Lemma 3.7. Let us consider the $P_{\mathbb{J}, +}$ -graded quotient $R' = \bigoplus_{\lambda \in P_{\mathbb{J}, +}} R'(\lambda)$ of $R_w^v(\mathbb{J})_{\mathbb{F}_p}$ that annihilates all the irrelevant irreducible components. Since $R_w^v(\mathbb{J})_{\mathbb{F}_p}$ is a reduced ring, we have $R_w^v(\mathbb{J})_{\mathbb{F}_p} = R'(\lambda)$ for $\lambda \in P_{\mathbb{J}, ++}$. The ring R' is generated by $\bigoplus_{i \in \mathbb{I} \setminus \mathbb{J}} R'(\varpi_i)$ as $R_w^v(\mathbb{J})_{\mathbb{F}_p}$ is so. We have

$$H^0(\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{F}_p}, \mathcal{O}_{\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{F}_p}}(\lambda)) \cong R_w^v(\mathbb{J})_{\mathbb{F}_p}(\lambda) \quad (3.14)$$

for sufficiently large $\lambda \in P_{\mathbb{J}, +}$ (see e.g. [37, II Excecise 5.9]). In other words, (3.14) hold for $m\lambda$, where $\lambda \in P_{\mathbb{J}, ++}$ is arbitrary and $m \gg 0$.

Let $R^{\sharp}(\lambda) := H^0(\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{F}_p}, \mathcal{O}_{\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{F}_p}}(\lambda))$ for $\lambda \in P_{\mathbb{J}, ++}$. We set $R^{\sharp}[\lambda] := \mathbb{F}_p 1 \oplus \bigoplus_{m \geq 1} R^{\sharp}(m\lambda)$ and $R'[\lambda] := \bigoplus_{m \geq 0} R'(m\lambda) \subset R'$ for $\lambda \in P_{\mathbb{J}, ++}$. Both are naturally rings. Then, we have a ring extension $R'[\lambda] \subset R^{\sharp}[\lambda]$ such that their degree $m \gg 0$ -parts are the same.

We prove $R'(\lambda) = R^{\sharp}(\lambda)$. The ring $R'[\lambda]$ is weakly normal as we have an inclusion $R'[\lambda] \hookrightarrow R_w^v(\mathbb{J})_{\mathbb{F}_p}$ (and the reasoning that $R_w^v(\mathbb{J})_{\mathbb{F}_p}$ is weakly normal equally applies to $R'[\lambda]$). Since it is generated by $R'(\lambda)$, we deduce that two rings $R'[\lambda]$ and $R^{\sharp}[\lambda]$ share the same spectrum (the irrelevant locus of $R'[\lambda]$ is one point as it is a $\mathbb{Z}_{\geq 0}$ -graded ring, and other points are just the same). This forces $R'(\lambda) = R^{\sharp}(\lambda)$ by the weak normality of $R'[\lambda]$.

This yields the H^0 -part of the assertion for $\text{char } \mathbb{K} > 0$ through the extension of scalars. The H^0 -part of the assertion for $\text{char } \mathbb{K} = 0$ is obtained by taking the generic specialization of the base scheme $\text{Spec } \mathbb{Z}$ of $\mathcal{Q}'_{\mathbb{J}}(v, w)$. \square

Corollary 3.35. *Let $w, v \in W_{\text{af}}$ and $\mathbb{J} \subset \mathbb{I}$. Assume that the map $\Pi_{\mathbb{J}} : \mathcal{Q}'(v, w)_{\mathbb{K}} \rightarrow \mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}$ defined through the projective coordinate ring is surjective. We have*

$$\mathbb{R}^{>0}(\Pi_{\mathbb{J}})_* \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}} \cong \{0\} \text{ and } (\Pi_{\mathbb{J}})_* \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}} \cong \mathcal{O}_{\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}}.$$

Remark 3.36. Since the map $\mathbf{Q}_G^{\text{rat}} \rightarrow \mathbf{Q}_{G, \mathbb{J}}^{\text{rat}}$ is surjective, we can replace v with vt_{β} for some $\beta \in \sum_{j \in \mathbb{J}} \mathbb{Z}_{\geq 0} \alpha_j^{\vee}$ to obtain a surjection:

$$\Pi_{\mathbb{J}} : \mathcal{Q}'(vt_{\beta}, w)_{\mathbb{K}} \rightarrow \mathcal{Q}'_{\mathbb{J}}(vt_{\beta}, w)_{\mathbb{K}} = \mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}.$$

For $v = w_0$ and $w = e$, there is an optimal choice of β , that is sometimes referred to as the Peterson-Woodward threshold ([79]).

Proof of Corollary 3.35. Thanks to [57, Lemma A.31] and Lemma 3.29, it suffices to prove

$$H^{>0}(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\lambda)) \cong \{0\} \text{ and } H^0(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\lambda)) \cong H^0(\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'_{\mathbb{J}}(v, w)_{\mathbb{K}}}(\lambda))$$

for each $\lambda \in P_{J,+++}$. The $H^{>0}$ -part of the assertion follow from Theorem 3.33. The H^0 -part of the assertion follow as in the latter part of the proof of Theorem 3.33 since the surjectivity of Π_J guarantees that every non-zero element of $R_w^v(\lambda)_{\mathbb{F}_p}$ is supported outside of the irrelevant locus by pullback and (graded) normalization of the ring $R_w^v(J)_{\mathbb{F}_p}$ is the degree $P_{J,+}$ -part of the (graded) normalization of the ring $(R_w^v)_{\mathbb{F}_p}$ by the degree reasons. \square

Corollary 3.37. *Let $\lambda \in P_+$ and $w, v \in W_{\text{af}}$. We set $J := \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$. Assume that $\Pi_J(\mathcal{Q}'(v, w)_{\mathbb{K}}) = \mathcal{Q}'_J(v, w)_{\mathbb{K}}$. Then, we have*

$$H^0(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\lambda))^\vee = \mathbb{W}_{ww_0}(\lambda)_{\mathbb{K}} \cap \theta(\mathbb{W}_v(-w_0\lambda)_{\mathbb{K}}).$$

Proof. Combine Theorem 3.33 for $\mathcal{Q}'_J(v, w)_{\mathbb{K}}$ and Corollary 3.35. \square

Lemma 3.38. *Let $w, v \in W_{\text{af}}$ and $J \subset I$. For each $i \in I_{\text{af}}$ such that $s_i w >_{\frac{\infty}{2}} w$ and $s_i v >_{\frac{\infty}{2}} v$, the variety $\mathcal{Q}'_J(v, w)_{\mathbb{K}}$ is B_i -stable. In addition, we have a morphism*

$$\pi_i : SL(2, i) \times^{B_i} \mathcal{Q}'_J(v, w)_{\mathbb{K}} \rightarrow \mathcal{Q}'_J(v, s_i w)_{\mathbb{K}}.$$

The map π_i is a \mathbb{P}^1 -fibration if $\mathcal{Q}'_J(v, w)_{\mathbb{K}}$ is $SL(2, i)$ -stable (this never happens under our assumption when $J = \emptyset$). In general, the fiber of π_i is either a point or \mathbb{P}^1 if it is nonempty.

Proof. For each $\lambda \in P_{J,+++}$, the module $\mathbb{W}_v(-w_0\lambda)_{\mathbb{K}}$ is $\mathbf{I}(i)$ -stable by Corollary 2.16 and (1.2) and the module $\mathbb{W}_{ww_0}(\lambda)_{\mathbb{K}}$ is \mathbf{I} -stable. In particular,

$$\mathbb{W}_{ww_0}(\lambda)_{\mathbb{K}} \cap \theta(\mathbb{W}_v(-w_0\lambda)_{\mathbb{K}}) \cong \mathbb{W}_{ww_0}(\lambda)_{\mathbb{K}} \cap \mathbb{W}_{vw_0}^-(\lambda)_{\mathbb{K}}$$

is B_i -stable. It follows that the ring $R_w^v(J)_{\mathbb{K}}$ is B_i -stable. Hence the scheme $\mathcal{Q}'_J(v, w)_{\mathbb{K}}$ is also B_i -stable.

The map π_i is a \mathbb{P}^1 -fibration if $\mathcal{Q}'_J(v, w)_{\mathbb{K}}$ is $SL(2, i)$ -stable since we have $\mathcal{Q}'_J(\beta, w)_{\mathbb{K}} = \mathcal{Q}'_J(\beta, s_i w)_{\mathbb{K}}$ (that in turn holds if and only if $\mathbb{W}_{s_i w w_0}(\lambda)_{\mathbb{K}} = \mathbb{W}_{ww_0}(\lambda)_{\mathbb{K}}$ for $\lambda \in P_{J,+++}$ since $\mathbb{W}_{vw_0}^-(\lambda)_{\mathbb{K}}$ is $SL(2, i)$ -stable; hence it never happens when $J = \emptyset$ by Lemma 2.15) in this case. The fiber of the map π_i is either a point or \mathbb{P}^1 if $\mathcal{Q}'_J(v, w)_{\mathbb{K}}$ is not $SL(2, i)$ -stable as the corresponding statement holds true for $\mathbf{Q}_{G, J}(w)_{\mathbb{K}} \subset \mathbf{Q}_{G, J}(s_i w)_{\mathbb{K}}$ by a set-theoretic consideration (and it carries over to any B_i -stable locus). Therefore, we conclude the result. \square

Proposition 3.39. *Let $w, v \in W_{\text{af}}$ and $J \subset I$. For each $i \in I_{\text{af}}$ such that $s_i w >_{\frac{\infty}{2}} w$ and $s_i v >_{\frac{\infty}{2}} v$, we have a surjective map*

$$\pi_i : SL(2, i) \times^{B_i} \mathcal{Q}'_J(v, w)_{\mathbb{K}} \rightarrow \mathcal{Q}'_J(v, s_i w)_{\mathbb{K}}$$

such that $(\pi_i)_ \mathcal{O}_{SL(2, i) \times^{B_i} \mathcal{Q}'_J(v, w)_{\mathbb{K}}} \cong \mathcal{O}_{\mathcal{Q}'_J(v, s_i w)_{\mathbb{K}}}$ and*

$$\mathbb{R}^{>0}(\pi_i)_* \mathcal{O}_{SL(2, i) \times^{B_i} \mathcal{Q}'_J(v, w)_{\mathbb{K}}} \cong \{0\}.$$

Proof. For each $\lambda \in P_{J,+++}$, the λ -graded component

$$R_w^v(J, \lambda)^\vee = \mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}} \cap \theta^*(\mathbb{W}_v(-w_0\lambda)_{\mathbb{Z}}) \subset \mathbb{X}(\lambda)_{\mathbb{Z}}$$

of the coordinate ring of $\mathcal{Q}'_J(v, w)$ is obtained as the \mathbb{Z} -span of a subset of positive global basis of $\mathbb{W}_{ww_0}(\lambda)_{\mathbb{Z}}$.

We set $\mathcal{Q}_J^+(v, w)_{\mathbb{K}} := SL(2, i) \times^{B_i} \mathcal{Q}'_J(v, w)_{\mathbb{K}}$ for simplicity in the below.

In view of [72, Theorem 4.2.1] (cf. Corollary 2.16), the basis elements of $\theta^*(\mathbb{W}_v(-w_0\lambda)_{\mathbb{Z}})$ afforded by the negative global basis, regarded as a subset of the $\tilde{\mathfrak{g}}$ -crystal $\mathbf{B}(\mathbb{X}(\lambda))$, decomposes into the disjoint union of connected $\mathfrak{sl}(2, i)$ -crystals (that is the set of labels of a basis of an irreducible $\mathfrak{sl}(2)$ -module equipped with combinatorial operations \tilde{e}_i and \tilde{f}_i corresponding to $E_i^{(\bullet)}$ and $F_i^{(\bullet)}$, see e.g. [41, Definition 2.3.1]; it is the i -string in [42, 45]). Moreover, the basis elements of $\mathbb{W}_{s_i w w_0}(\lambda)_{\mathbb{Z}}$ afforded by the positive global basis, regarded as a subset of the $\tilde{\mathfrak{g}}$ -crystal $\mathbf{B}(\mathbb{X}(\lambda))$, also decomposes into the disjoint union of $\mathfrak{sl}(2, i)$ -crystals ([45, 46]), and the set S of basis elements corresponding to each connected $\mathfrak{sl}(2, i)$ -crystal satisfies

$$S \cap \mathbb{W}_{w w_0}(\lambda)_{\mathbb{Z}} = S \quad \text{or} \quad \{\mathbf{v}\}, \quad (3.15)$$

where $\mathbf{v} \in S$ is the highest weight vector as a $\mathfrak{sl}(2, i)$ -crystal, that has dominant weight as a weight of $\mathfrak{sl}(2, i)$ ([45, Lemma 2.6]). In particular, we have

$$S \cap R_w^v(\mathbf{J}, \lambda)^{\vee} = S \quad \text{or} \quad \{\mathbf{v}\} \quad \text{or} \quad \emptyset.$$

Therefore, a reinterpretation of [42, (0.5)] using Theorem 3.33 (cf. [39, 46]) reads as

$$\begin{aligned} \text{ch } H^0(\mathcal{O}_{\mathcal{Q}'_J(v, s_i w)_{\mathbb{K}}}(\lambda)) &= \text{ch } H^0(\mathcal{O}_{\mathcal{Q}_J^+(v, w)_{\mathbb{K}}}(\lambda)) \\ 0 &= \text{ch } H^1(\mathcal{O}_{\mathcal{Q}_J^+(v, w)_{\mathbb{K}}}(\lambda)). \end{aligned}$$

for each $\lambda \in P_{\mathbf{J}, ++}$. Since the zero-th part of the Demazure functor is the same as taking the maximal integrable $SL(2, i)$ -inflation of a (weight semisimple) B_i -module, we conclude that inclusion $R_w^v(\mathbf{J}, \lambda)_{\mathbb{K}}^{\vee} \subset R_{s_i w}^v(\mathbf{J}, \lambda)_{\mathbb{K}}^{\vee}$ induces a natural inclusion

$$H^0(\mathcal{Q}'_J(v, s_i w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'_J(\beta, s_i w)_{\mathbb{K}}}(\lambda)) \hookrightarrow H^0(\mathcal{Q}_J^+(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}_J^+(v, w)_{\mathbb{K}}}(\lambda)), \quad (3.16)$$

that is an isomorphism by the character comparison. Thus, Lemma 3.29 implies that

$$(\pi_i)_* \mathcal{O}_{\mathcal{Q}_J^+(v, w)_{\mathbb{K}}} \cong \mathcal{O}_{\mathcal{Q}'_J(v, s_i w)_{\mathbb{K}}} \quad \text{and} \quad \mathbb{R}^1(\pi_i)_* \mathcal{O}_{\mathcal{Q}_J^+(\beta, w)_{\mathbb{K}}} \cong \{0\}.$$

Since we have $\mathbb{R}^{\geq 2}(\pi_i)_* \mathcal{O}_{\mathcal{Q}_J^+(v, w)_{\mathbb{K}}} \cong \{0\}$ by the dimension reason, we conclude the result. \square

A particular case of Proposition 3.39 is worth noting:

Corollary 3.40. *Let $\beta \in Q^{\vee}$, $w \in W$, and $\mathbf{J} \subset \mathbf{I}$. For each $i \in \mathbf{I}$ such that $s_i w < w$, we have a surjective map*

$$\pi_i : P_i \times^B \mathcal{Q}'_J(\beta, w)_{\mathbb{K}} \rightarrow \mathcal{Q}'_J(\beta, s_i w)_{\mathbb{K}}$$

such that $\mathbb{R}^{\bullet}(\pi_i)_* \mathcal{O}_{P_i \times^B \mathcal{Q}'_J(\beta, w)_{\mathbb{K}}} \cong \mathcal{O}_{\mathcal{Q}'_J(\beta, s_i w)_{\mathbb{K}}}$. \square

3.5 Lifting to/from characteristic zero

Theorem 3.41. *Let \mathfrak{X} be a Noetherian scheme flat over \mathbb{Z} . If $\mathfrak{X}_{\mathbb{F}_p}$ is weakly normal for $p \gg 0$, then $\mathfrak{X}_{\mathbb{C}}$ is also weakly normal.*

Proof. Since the weak normalization commutes with localization [68, Theorem IV.3], we can argue locally. Let (S, \mathfrak{m}) be a local ring of $\mathfrak{X}_{\overline{\mathbb{Q}}}$ and let S^- be the weak normalization of S ([80, Remark 1]). By the Noetherian hypothesis, we can invert finitely many primes and take a finite algebraic extension of \mathbb{Z} to obtain a ring A such that we have a commutative ring S_A over A and its ideal \mathfrak{m}_A with the following properties:

- We have $(S_A \otimes_A \overline{\mathbb{Q}}, \mathfrak{m}_A \otimes_A \overline{\mathbb{Q}}) \cong (S, \mathfrak{m})$;
- The A -modules S_A , \mathfrak{m}_A , and S_A/\mathfrak{m}_A are torsion-free;
- The specialization of A to the algebraic closure of a finite field yields a weakly normal (local) ring ([68, Theorem V.2]).

As A is a Dedekind domain, we find that S_A , \mathfrak{m}_A , and S_A/\mathfrak{m}_A are flat over A .

We have $S^- = S[f_1, \dots, f_n]$, where f_1, \dots, f_n are integral elements. By multiplying with elements in $\overline{\mathbb{Q}}$, we can assume that f_1, \dots, f_n are integral over S_A . By inverting additional primes in \mathbb{Z} if necessary (to assume that the denominator of f_i in $\text{Frac}(S)$ does not vanish along specializations and achieve the conditions in the followings), we can further assume that $S_A^- := S_A[f_1, \dots, f_n]$ is flat over A and it is integral for every specialization of A to a field.

The ring $(S_A^-/\mathfrak{m}_A S_A^-) \otimes_A \mathbb{C}$ is a finite-dimensional local commutative \mathbb{C} -algebra by the weak normality assumption on S^- (see [80, Remark 1]). In particular, the multiplication action of each element of $(S_A^-/\mathfrak{m}_A S_A^-) \otimes_A \mathbb{C}$ have a unique eigenvalue. Hence, if we present $(S_A^-/\mathfrak{m}_A S_A^-) = A[X_1, \dots, X_m]/\sim$ (where \sim contains the minimal polynomials of the A -valued matrix X_i), then the minimal polynomial of X_i is of the form $(T - a_i)^{m_i}$ ($a_i \in A$). Therefore, we can assume that each X_i is nilpotent by changing X_i with $X_i + a_i$ if necessary. Hence, we conclude that $(X_1, \dots, X_m) \subset (S_A^-/\mathfrak{m}_A S_A^-)$ have eigenvalues zero after specializing to $\overline{\mathbb{F}}_p$.

By assumption, $(S_A) \otimes_A \overline{\mathbb{F}}_p$ is weakly normal for every possible prime p and every ring homomorphism $A \rightarrow \overline{\mathbb{F}}_p$. Hence, the specialization $(S_A^-/\mathfrak{m}_A S_A^-) \otimes_A \overline{\mathbb{F}}_p$ must contain $\overline{\mathbb{F}}_p$ as its ring direct summand (if it is non-zero). This forces $\text{rank}(S_A^-/\mathfrak{m}_A S_A^-) = 1$ since we cannot have two linearly independent idempotents that have distinct eigenspaces by the previous paragraph.

Therefore, we deduce that $S_A^- = S_A$, that implies (S, \mathfrak{m}) is itself weakly normal. In view of [68, Theorem V.2 and Corollary V.3], we conclude the assertion. \square

Corollary 3.42. *For each $J \subset I$, $w, v \in W_{\text{af}}$, the scheme $\mathcal{Q}'_J(v, w)_{\mathbb{C}}$ is weakly normal.*

Proof. Apply Theorem 3.41 to Lemma 3.7. \square

Proposition 3.43. *For each $J \subset I$, $v, w \in W$, the variety $\mathcal{Q}'_J(v, w)_{\overline{\mathbb{F}}_p}$ is normal for $p \gg 0$ provided if $\mathcal{Q}'_J(v, w)_{\mathbb{C}}$ is normal. The same is true for the irreducibility.*

Proof. Since $\mathcal{Q}'_J(v, w)$ is defined over \mathbb{Z} , the scheme $\mathcal{Q}'_J(v, w)_{\mathbb{C}}$ is a scalar extension of $\mathcal{Q}'_J(v, w)_{\overline{\mathbb{Q}}}$. By [77, Lemma 038P], we deduce $\mathcal{Q}'_J(v, w)_{\overline{\mathbb{Q}}}$ is normal. We apply [77, Lemma 0364] to derive the irreducibility of $\mathcal{Q}'_J(v, w)_{\overline{\mathbb{Q}}}$. Now apply [36, Proposition 9.9.4 and Théorème 9.7.7] to $\mathcal{Q}'_J(v, w)_{\mathbb{Z}} \rightarrow \text{Spec } \mathbb{Z}$. \square

4 Normality of quasi-map spaces

In this section, we continue to work under the setting of the previous section with an exception that $\mathbb{K} = \mathbb{C}$. Also, a point of a scheme (over \mathbb{C}) means a closed point unless stated otherwise.

4.1 Graph space resolution of $\mathcal{Q}(\beta)$

We refer to [56, 30, 6, 23, 33] for precise explanations of the material in this subsection. For each non-negative integer n and $\beta \in Q_+^\vee$, we set $\mathcal{GB}_{n,\beta}$ to be the space of stable maps of genus zero curves with n -parked points to $(\mathbb{P}^1 \times \mathcal{B})$ of bidegree $(1, \beta)$, that is also called the graph space of \mathcal{B} . A point of $\mathcal{GB}_{n,\beta}$ represents a genus zero curve C with n -marked points, together with a map to \mathbb{P}^1 of degree one (obtained by composing the map from C to $\mathbb{P}^1 \times \mathcal{B}$ with the first projection of the target). Hence, we have a unique \mathbb{P}^1 -component of C that maps isomorphically onto \mathbb{P}^1 . We call this component the main component of C and denote it by C_0 . The space $\mathcal{GB}_{n,\beta}$ is a normal projective variety by [30, Theorem 2] that have at worst quotient singularities arising from the automorphism of curves (and hence it is smooth as an orbifold). The natural $(H \times \mathbb{G}_m)$ -action on $(\mathbb{P}^1 \times \mathcal{B})$ induces a natural $(H \times \mathbb{G}_m)$ -action on $\mathcal{GB}_{n,\beta}$.

We have a morphism $\pi_{n,\beta} : \mathcal{GB}_{n,\beta} \rightarrow \mathcal{Q}(\beta)$ that factors through $\mathcal{GB}_{0,\beta}$ (Givental's main lemma [34]; see [23, §8.3]). Let $\mathbf{e}_j : \mathcal{B}_{n,\beta} \rightarrow \mathcal{B}$ ($1 \leq j \leq n$) be the evaluation at the j -th marked point, and let $\mathbf{ev}_j : \mathcal{GB}_{n,\beta} \rightarrow \mathcal{B}$ be the j -th evaluation map to $\mathbb{P}^1 \times \mathcal{B}$ composed with the second projection.

Since $\mathcal{Q}(\beta)$ is irreducible (Theorem 1.7), [23, §8.3] asserts that $\mathcal{GB}_{n,\beta}$ is irreducible (as a special feature of flag varieties, see [30, §1.2] and [53]).

4.2 The variety $\mathcal{Q}(\beta, v, w)$

Let $\mathcal{GB}_{2,\beta}^b$ denote the subvariety of $\mathcal{GB}_{2,\beta}$ consisting of points such that the first marked point projects to $0 \in \mathbb{P}^1$, and the second marked point projects to $\infty \in \mathbb{P}^1$ through the projection of quasi-stable curves C to their main component $C_0 \cong \mathbb{P}^1$. Let us denote the restrictions of \mathbf{ev}_i ($i = 1, 2$) and $\pi_{2,\beta}$ to $\mathcal{GB}_{2,\beta}^b$ by the same letter. By [11, 12], $\mathcal{GB}_{2,\beta}^b$ gives a resolution of singularities of $\mathcal{Q}(\beta)$ (in an orbifold sense).

Recall that each Schubert cell $\mathbb{O}_{\mathcal{B}}(w)$ contains a unique H -fixed point p_w . For each $w \in W$, we set

$$\mathbb{O}_{\mathcal{B}}^{\text{op}}(w) := N^- p_w \subset \mathcal{B}, \quad \text{and} \quad \mathcal{B}^{\text{op}}(w) := \overline{\mathbb{O}_{\mathcal{B}}^{\text{op}}(w)} = \overline{N^- p_w} \subset \mathcal{B}.$$

For $w, v \in W$, we define

$$\mathcal{GB}_{2,\beta}^b(w, v) := \mathbf{ev}_1^{-1}(\mathcal{B}(w)) \cap \mathbf{ev}_2^{-1}(\mathcal{B}^{\text{op}}(v)) \subset \mathcal{GB}_{2,\beta}^b$$

and

$$\mathcal{B}_{2,\beta}(w, v) := \mathbf{e}_1^{-1}(\mathcal{B}(w)) \cap \mathbf{e}_2^{-1}(\mathcal{B}^{\text{op}}(v)) \subset \mathcal{B}_{2,\beta}.$$

Theorem 4.1 (Buch-Chaput-Mihalcea-Perrin [16]). *For each $v, w \in W$ and $\beta \in Q_+^\vee$, the variety $\mathcal{GB}_{2,\beta}^b(w, v)$ is either empty or a unirational (and hence connected and irreducible) variety that has rational singularities. The same is true for $\mathcal{B}_{2,\beta}(w, v)$. In particular, they are normal.*

Proof. Since the both cases are parallel, we concentrate into the case $\mathcal{GB}_{2,\beta}^b(w, v)$.

As $\mathbb{P}^1 \times \mathcal{B}$ is a homogeneous variety under the group action of $(SL(2) \times G)$, Kim-Phandaripande [53, Theorem 2 and Theorem 3] applies and hence $\mathcal{GB}_{2,\beta}$ is a rational variety. Then, a pair of Schubert subvarieties (with respect to a pair of opposite Borel subgroups of $SL(2) \times G$) of $\mathbb{P}^1 \times \mathcal{B}$ presented as $\{0\} \times \mathcal{B}(w)$ and $\{\infty\} \times \mathcal{B}^{\text{op}}(v)$ is used to define $\mathcal{GB}_{2,\beta}^b(w, v)$. Hence, [16, Proposition 3.2 c)] implies that $\mathcal{GB}_{2,\beta}^b(w, v)$ is either empty or unirational (and hence connected). Since a pair of Schubert varieties with respect to the opposite Borel subgroups forms the dense subset of the pair of translations of Schubert varieties by applying the G -action, it must contain a pair of Schubert varieties in general position. Therefore, [16, Corollary 3.1] implies that $\mathcal{GB}_{2,\beta}^b(w, v)$ has rational singularity. The last assertion is a well-known property of rational singularities [55, Definition 5.8]. \square

Corollary 4.2. *In the setting of Theorem 4.1, we additionally assume that $\mathcal{GB}_{2,\beta}^b(w, v)$ is nonempty. Then, we have*

$$\dim \mathcal{GB}_{2,\beta}^b(w, v) = \dim \mathcal{Q}(\beta) - \ell(w) - \ell(vw_0).$$

Proof. By [16, Proposition 3.2 b)], the dimension count of (a Zariski open subset of) $\mathcal{GB}_{2,\beta}^b(w, v)$ can be borrowed from [54, Theorem 2] applied to the $(SL(2) \times G)$ -action on $\mathbb{P}^1 \times \mathcal{B}$. In addition, the map $\pi_{2,\beta}$ is birational. Therefore, we have

$$\begin{aligned} \dim \mathcal{GB}_{2,\beta}^b(w, v) &= \dim \mathcal{Q}(\beta) - \text{codim}_{\mathcal{B}} \mathcal{B}(w) - \text{codim}_{\mathcal{B}} \mathcal{B}^{\text{op}}(v) \\ &= \dim \mathcal{Q}(\beta) - \ell(w) - \ell(vw_0) \end{aligned}$$

as required. \square

Proposition 4.3. *For each $v, w \in W$ and $\beta \in Q_+^\vee$, the variety $\mathcal{Q}'(vt_\beta, w)_{\mathbb{C}}$ is irreducible.*

Proof. We define

$$\mathring{\mathcal{Q}}'(vt_\beta, w)_{\mathbb{C}} := \mathcal{Q}'(vt_\beta, w)_{\mathbb{C}} \setminus \left(\left(\bigcup_{w' <_{\frac{\infty}{2}} w} \mathcal{Q}'(vt_\beta, w')_{\mathbb{C}} \right) \cup \bigcup_{v' >_{\frac{\infty}{2}} v} \mathcal{Q}'(v't_\beta, w)_{\mathbb{C}} \right).$$

In view of Corollary 3.31, every irreducible component of $\mathcal{Q}'(vt_\beta, w)_{\mathbb{C}}$ has dimension $\ell_{\frac{\infty}{2}}(vt_\beta) - \ell_{\frac{\infty}{2}}(w)$ if it is nonempty. We have $\mathcal{Q}'(vt_\beta, w)_{\mathbb{C}} \neq \emptyset$ if and only if $vt_\beta \leq_{\frac{\infty}{2}} w$ by Corollary 3.21. In particular, we find that $\mathring{\mathcal{Q}}'(vt_\beta, w)_{\mathbb{C}} \neq \emptyset$ if $vt_\beta \leq_{\frac{\infty}{2}} w$, and all of its irreducible components have dimension

$$\ell_{\frac{\infty}{2}}(vt_\beta) - \ell_{\frac{\infty}{2}}(w) = \ell(v) - \ell(w) + 2 \langle \beta, \rho \rangle \quad (4.1)$$

in that case. A point of $\mathring{\mathcal{Q}}'(vt_\beta, w)_{\mathbb{C}}$ can be seen as a quasi-map of degree β with no defects at 0 and ∞ (Theorem 3.30). In particular, $\mathcal{GB}_{2,\beta}^b(w, v) \neq \emptyset$ if $\mathring{\mathcal{Q}}'(vt_\beta, w)_{\mathbb{C}} \neq \emptyset$. In case $\mathring{\mathcal{Q}}'(vt_\beta, w)_{\mathbb{C}} \neq \emptyset$, we have

$$\begin{aligned} \dim \mathcal{GB}_{2,\beta}^b(w, v) &= \dim \mathcal{Q}(\beta) - \ell(w) - \ell(vw_0) \\ &= 2 \langle \beta, \rho \rangle + \dim \mathcal{B} - \ell(w_0) + \ell(v) - \ell(w) = (4.1), \end{aligned} \quad (4.2)$$

where the first equality follows from Corollary 4.2, and the second equality follows from Theorem 1.7. By the surjectivity of $\pi_{2,\beta}$ and (4.2), we find that the rational map $\mathcal{GB}_{2,\beta}^b(w, v) \dashrightarrow \mathring{Q}'(vt_\beta, w)_\mathbb{C}$ is surjective and generically finite (it is in fact birational by Theorem 4.7 and Proposition 4.11). In view of Theorem 4.1, the variety $\mathcal{GB}_{2,\beta}^b(w, v)$ is irreducible if it is nonempty. Therefore, $\mathring{Q}'(vt_\beta, w)_\mathbb{C} \neq \emptyset$ implies that $\mathring{Q}'(vt_\beta, w)_\mathbb{C}$ is irreducible. Again by Corollary 3.31, we conclude that $Q'(vt_\beta, w)_\mathbb{C}$ has a unique irreducible component as required. \square

In view of Proposition 4.3, we set $Q_J(\beta, v, w) := Q'_J(vt_\beta, w)_\mathbb{C}$ for each $J \subset \mathbf{I}$, $\beta \in Q_+^\vee$ and $v, w \in W$ in the below (see also Corollary 4.21). We have $Q(\beta, w_0, w) = Q(\beta, w)$ by Theorem 3.30.

Corollary 4.4 (of proof of Proposition 4.3). *For each $\beta \in Q_+^\vee$ and $v, w \in W$, the scheme $Q(\beta, v, w)$ is nonempty if and only if $vt_\beta \leq_{\frac{\infty}{2}} w$. If it is nonempty, then it has dimension $\ell(v) - \ell(w) + 2\langle \beta, \rho \rangle$. Moreover, the space $Q(\beta, v, w)$ (viewed as a subspace of $Q(\beta)$) contains a quasi-map with no defects at 0 and ∞ . \square*

Remark 4.5. Corollary 4.4 removes the condition $\gamma \gg 0$ in [29, Lemma 8.5.2].

Thanks to Proposition 4.3 and its proof, we deduce that the map $\pi_{2,\beta}$ restricts to a $(H \times \mathbb{G}_m)$ -equivariant birational proper map

$$\pi_{\beta,w,v} : \mathcal{GB}_{2,\beta}^b(w, v) \rightarrow Q(\beta, v, w)$$

by inspection.

4.3 From Givental's main lemma

For each $w, v \in W$, we define subvarieties of $\mathcal{B}_{2,\beta}$ as:

$$\mathcal{B}_{2,\beta}[w] := \mathbf{e}_1^{-1}(p_w) \quad \text{and} \quad \mathcal{B}_{2,\beta}[w, v] := \mathbf{e}_1^{-1}(p_w) \cap \mathbf{e}_2^{-1}(p_v).$$

Similarly, we set

$$\mathcal{GB}_{2,\beta}^b[w, v] := \mathbf{ev}_1^{-1}(p_w) \cap \mathbf{ev}_2^{-1}(p_v) \subset \mathcal{GB}_{2,\beta}^b \quad \text{and} \quad \mathcal{B}_{1,\beta}[w] := \mathbf{e}_1^{-1}(p_w) \subset \mathcal{B}_{1,\beta}[w].$$

Lemma 4.6. *For each $x, y \in \mathcal{B}$ and $\beta \in Q_+^\vee$, there exists $w \in W$ such that*

$$\mathbf{e}_1^{-1}(x) \cap \mathbf{e}_2^{-1}(y) \cong \mathcal{B}_{2,\beta}[w, w_0].$$

The same is true for \mathbf{ev} and \mathcal{GB} .

Proof. We consider only the case of \mathbf{e} and \mathcal{B} as the other case is completely parallel. Since $(x, y) \in \mathcal{B} \times \mathcal{B}$ and the G -action on \mathcal{B} is transitive, we can assume $y = p_{w_0}$. Since we have $\mathbf{Stab}_G y = B$, we can further rearrange $x = p_w$ for some $w \in W$ by (1.4). \square

Theorem 4.7 (Givental's main lemma [34], see [23] §8). *Let $\beta \in Q_+^\vee$. Let $(f, D) \in Q(\beta)$ be a quasi-map with its defect $D = \sum_{x \in \mathbb{P}^1(\mathbb{C})} \beta_x \otimes [x]$. Then, we have*

$$\pi_{2,\beta}^{-1}(f, D) \cong \mathcal{B}_{2,\beta_0}[w] \times \mathcal{B}_{2,\beta_\infty}[w] \times \prod_{x \in \mathbb{P}^1(\mathbb{C}) \setminus \{0, \infty\}} \mathcal{B}_{1,\beta_x}[w] \subset \mathcal{GB}_{2,\beta}^b.$$

In particular, the map $\pi_{2,\beta}$ is an isomorphism along the locus with $D = 0$. \square

Remark 4.8. In Theorem 4.7, the first marked points of (the stable maps in) $\mathcal{GB}_{2,\beta}^b$ (the marked points at $0 \in \mathbb{P}^1$; i.e. the image of \mathbf{ev}_1) is identified with the second marked points of (the stable maps in) $\mathcal{B}_{2,\beta_0}[w]$ (i.e. the image of \mathbf{e}_2), and the second marked points of $\mathcal{GB}_{2,\beta}^b$ (the marked points at $\infty \in \mathbb{P}^1$) is identified with the second marked points of $\mathcal{B}_{2,\beta_\infty}[w]$. (Other marked points are used to glue pieces of stable maps together.)

Lemma 4.9 ([29] Lemma 8.5.1). *For each $\beta \in Q_+^\vee$ such that $\langle \beta, \alpha_i \rangle \geq 1$ for all $i \in \mathbf{I}$, the evaluation map*

$$\mathbf{ev} := (\mathbf{ev}_1 \times \mathbf{ev}_2) : \mathcal{GB}_{2,\beta}^b \longrightarrow \mathcal{B} \times \mathcal{B}$$

is surjective.

Proof. Taking into account the fact that $\mathcal{GB}_{2,\beta}^b$ is projective, it suffices to prove that the tangent map associated to \mathbf{ev} is surjective on a dense open subset of $\mathcal{GB}_{2,\beta}^b$.

Since the map $\pi_{2,\beta}$ is birational by Theorem 4.7, we replace the problem with the case of a genuine map $f : \mathbb{P}^1 \rightarrow \mathcal{B}$. Thanks to [29, Proposition 3.5], $\mathcal{Q}(\beta)$ and hence $\mathcal{GB}_{2,\beta}^b$ are smooth at f . Moreover, its tangent space is described as

$$H^0(\mathbb{P}^1, f^*T\mathcal{B}), \text{ and } H^1(\mathbb{P}^1, f^*T\mathcal{B}) = \{0\}$$

and the filtration of $T\mathcal{B}$ as G -equivariant line bundles yields the following associated graded

$$\bigoplus_{\alpha \in \Delta_+} H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(\langle \beta, \alpha \rangle)).$$

(Here we used $\langle \beta, \alpha_i \rangle \geq -1$ for each $i \in \mathbf{I}$.) In particular, we have

$$\dim H^0(\mathbb{P}^1, f^*T\mathcal{B}) = \sum_{\alpha \in \Delta_+} \dim H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(\langle \beta, \alpha \rangle)).$$

The effect of fixing the image of two points $0, \infty \in \mathbb{P}^1$ corresponds to imposing divisor twist by $\mathcal{O}_{\mathbb{P}^1}(-[0] - [\infty])$. We have

$$\begin{aligned} \dim H^0(\mathbb{P}^1, f^*T\mathcal{B} \otimes_{\mathcal{O}_{\mathbb{P}^1}} \mathcal{O}_{\mathbb{P}^1}(-[0] - [\infty])) &\leq \dim \bigoplus_{\alpha \in \Delta_+} H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(\langle \beta, \alpha \rangle - 2)) \\ &= \dim \bigoplus_{\alpha \in \Delta_+} H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(\langle \beta, \alpha \rangle)) - 2|\Delta_+| \\ &= \dim \mathcal{Q}(\beta) - 2 \dim \mathcal{B} \\ &= \dim \mathcal{GB}_{2,\beta}^b - 2 \dim \mathcal{B}. \end{aligned}$$

Here the first inequality comes from the short exact sequences, the second equality is $\langle \beta, \alpha_i \rangle \geq 1$ for each $i \in \mathbf{I}$, the third one is the smoothness of $\mathcal{Q}(\beta)$ at f and $|\Delta_+| = \dim \mathcal{B}$, and the fourth one is the birationality of $\pi_{2,\beta}$ (restricted to $\mathcal{GB}_{2,\beta}^b \subset \mathcal{GB}_{2,\beta}$).

From this, we deduce that $d\pi_{2,\beta}$ is generically surjective as required. \square

Proposition 4.10. *For each $\beta \in Q_+^\vee$ such that $\langle \beta, \alpha_i \rangle \geq 1$ for all $i \in \mathbf{I}$ and $w, v \in W$, the scheme $\mathcal{GB}_{2,\beta}^b[w, v]$ is connected and nonempty.*

Proof. The map \mathbf{ev} borrowed from Lemma 4.9 is G -equivariant, and $(\mathcal{B} \times \mathcal{B})$ admits an open dense G -orbit \mathbf{O} by the Bruhat decomposition. Since $\mathcal{G}\mathcal{B}_{2,\beta}^b$ is an irreducible variety, so is its Zariski open set $\mathbf{ev}^{-1}(\mathbf{O})$. If we consider $x \in \mathbf{O}$, then the irreducible components of $\mathbf{ev}^{-1}(x)$ (that must be finite as we consider varieties, that are finite types) must be permuted by $\mathbf{Stab}_G x \cong H$. Since H is connected, we cannot have a non-trivial action. Therefore, the irreducible component of $\mathbf{ev}^{-1}(x)$ must be unique. Thanks to the Stein factorization theorem [77, Theorem 03H0], the map \mathbf{ev} factors through the normalization Y of $(\mathcal{B} \times \mathcal{B})$ inside (the function field of) $\mathcal{G}\mathcal{B}_{2,\beta}^b$ and the map $\mathcal{G}\mathcal{B}_{2,\beta}^b \rightarrow Y$ has connected fiber. Since the general fiber of \mathbf{ev} is connected, so is the map $Y \rightarrow (\mathcal{B} \times \mathcal{B})$. It follows that $Y \rightarrow (\mathcal{B} \times \mathcal{B})$ is a birational map (that is also finite by the Stein factorization theorem). This implies that Y and $(\mathcal{B} \times \mathcal{B})$ share the same function field. Thus, we conclude that $Y \cong (\mathcal{B} \times \mathcal{B})$ by the normality of $(\mathcal{B} \times \mathcal{B})$. This particularly means that every fiber of \mathbf{ev} is connected. Therefore, the assertion follows by choosing $(H \times H)$ -fixed points in $(\mathcal{B} \times \mathcal{B})$ as particular cases. \square

Proposition 4.11. *For each $\beta \in Q_+^\vee$ such that $\langle \beta, \alpha_i \rangle \geq 1$ for all $i \in \mathbf{I}$ and $w, v \in W$, the scheme $\mathcal{B}_{2,\beta}[w, v]$ is connected and nonempty. For each $\beta \in Q_+^\vee$ and $w \in W$, the schemes $\mathcal{B}_{1,\beta}[w]$ and $\mathcal{B}_{2,\beta}[w]$ are connected and nonempty.*

Proof. We prove the first assertion. We have a rational map $\mathcal{G}\mathcal{B}_{2,\beta}^b[w, v] \rightarrow \mathcal{B}_{2,\beta}[w, v]$ obtained by forgetting the map to \mathbb{P}^1 . Moreover, the locus this map is not defined consists of stable maps whose main components have degree 0 and have only two marked points. By modifying the universal family by adjoining such two marked points in such a degree 0 main components, we conclude that $\mathcal{G}\mathcal{B}_{2,\beta}^b[w, v] \rightarrow \mathcal{B}_{2,\beta}[w, v]$ extends to a map of topological spaces. By examining the condition to be a stable map [30, §1.1], we deduce that this map is surjective onto the image. Therefore, the connectedness and the nonemptiness of $\mathcal{G}\mathcal{B}_{2,\beta}^b[w, v]$ implies these of $\mathcal{B}_{2,\beta}[w, v]$, that is the first assertion.

The second assertion is straight-forward from the irreducibility of $\mathcal{B}_{n,\beta}$, together with the fact that \mathbf{e}_1 is G -equivariant fiber bundle over \mathcal{B} (see the proof of Proposition 4.10). \square

Lemma 4.12. *Let $\beta \in Q_+^\vee$. Let $(f, D) \in \mathcal{Q}(\beta)$ be a quasi-map with its defect D with the following properties:*

- $D = \sum_{x \in \mathbb{P}^1(\mathbb{C})} \beta_x \otimes [x]$;
- $\langle \beta_0, \alpha_i \rangle \geq 1$ and $\langle \beta_\infty, \alpha_i \rangle \geq 1$ for each $i \in \mathbf{I}$.

Then, $\pi_{\beta,w,v}^{-1}(f, D)$ is connected for every $w, v \in W$ if it is nonempty.

Proof. By Proposition 4.11 and Theorem 4.7, we can forget about the contribution of $\mathcal{B}_{1,\beta_x}[w]$ when $x \neq 0, \infty$. By our assumption and Proposition 4.11, we know that

$$\mathcal{B}_{2,\beta_0}[u, w'] \neq \emptyset \text{ and } \mathcal{B}_{2,\beta_\infty}[u, v'] \neq \emptyset$$

and are nonempty and connected for each $u, w', v' \in W$. Since \mathcal{B}_{2,β_0} and $\mathcal{B}_{2,\beta_\infty}$ are proper, we always find a limit point with respect to the H -action. It follows that $\mathcal{B}_{2,\beta_0}[u] \cap \mathbf{e}_2^{-1}(\mathcal{B}(u'))$ (resp. $\mathcal{B}_{2,\beta_\infty}[u] \cap \mathbf{e}_2^{-1}(\mathcal{B}^{\text{op}}(u'))$) is connected for each $u, u' \in W$ as we can connect every two points by appropriately sending to/from H -limit points that are contained in a connected component of the form $\mathcal{B}_{2,\beta_0}[u, w']$ ($w' \in W$). Thanks to Theorem 4.7, we conclude the assertion. \square

4.4 Normality of $\mathcal{Q}(\beta, v, w)$

Let $\mathcal{Q}^+(\beta, v, w)$ be the normalization of $\mathcal{Q}(\beta, v, w)$ for each $\beta \in Q_+^\vee$ and $v, w \in W$. We denote the normalization map by $\eta_{\beta, v, w} : \mathcal{Q}^+(\beta, v, w) \rightarrow \mathcal{Q}(\beta, v, w)$.

Proposition 4.13. *For each $\beta \in Q_+^\vee$ and $w, v \in W$, the variety $\mathcal{Q}(\beta, v, w)$ is normal if and only if every fiber of $\pi_{\beta, v, w}$ is connected.*

Proof. As $\mathcal{G}\mathcal{B}_{2, \beta}^b(v, w)$ is normal and $\pi_{\beta, v, w}$ is proper, we know that

$$(\pi_{\beta, v, w})_* \mathcal{O}_{\mathcal{G}\mathcal{B}_{2, \beta}^b(v, w)} \cong \mathcal{O}_{\mathcal{Q}^+(\beta, v, w)}.$$

The properness of $\pi_{\beta, v, w}$ also implies that $\mathcal{O}_{\mathcal{Q}^+(\beta, v, w)}$ is a coherent sheaf on $\mathcal{Q}(\beta, v, w)$. For each closed point x of $\mathcal{Q}(\beta, v, w)$, we set

$$\Theta(x) := \dim_{\mathbb{C}} \mathcal{O}_{\mathcal{Q}^+(\beta, v, w)} \otimes_{\mathcal{O}_{\mathcal{Q}(\beta, v, w)}} \mathbb{C}_x. \quad (4.3)$$

By the Stein factorization theorem, the map $\eta_{\beta, v, w}$ is finite. By Corollary 3.42 and §4.2, we know that the variety $\mathcal{Q}(\beta, v, w)$ is weakly normal. From this, we deduce that $\Theta(x) = \#\eta_{\beta, v, w}^{-1}(x)$ (cf. [80, Remark 1]). Moreover, it counts the number of irreducible components of the fiber of $\eta_{\beta, v, w}$.

The coherence of $\mathcal{O}_{\mathcal{Q}^+(\beta, v, w)}$ implies that the RHS of (4.3) is an upper-semicontinuous function on $\mathcal{Q}(\beta, v, w)$, and hence so is Θ .

If we have $\Theta \equiv 1$ on $\mathcal{Q}(\beta, v, w)$, then we have $\mathcal{Q}^+(\beta, v, w) = \mathcal{Q}(\beta, v, w)$ by the weak normality of the latter (cf. [80, Remark 1]). Therefore, the if part of the assertion follows.

If we have $\Theta \neq 1$ on $\mathcal{Q}(\beta, v, w)$, then we have $\mathcal{Q}^+(\beta, v, w) \neq \mathcal{Q}(\beta, v, w)$. Hence, the only if part of the assertion follows.

These complete the proof of Proposition 4.13. \square

Corollary 4.14 (Braverman-Finkelberg). *For each $\beta \in Q_+^\vee$, the variety $\mathcal{Q}(\beta)$ is normal.*

Remark 4.15. Our proof of Corollary 4.14 is independent of [11] (however based on common former papers [29, 23]). Hence, we obtain a new proof of the normality of $\mathcal{Q}(\beta)$ and $\mathcal{Z}(\beta, w_0)$. Together with Theorem 3.33, Corollary 4.14 also makes the contents in [51] logically independent of [11] (cf. Appendix A).

Proof of Corollary 4.14. Recall that $\mathcal{Q}(\beta) = \mathcal{Q}(\beta, w_0, e)$. We borrow the upper semi-continuous function Θ that counts the number of connected components of the fiber of $\eta_{\beta, w_0, e}$ from (4.3) in the proof of Proposition 4.13.

By Proposition 4.13, it suffices to prove that $\Theta \equiv 1$ on $\mathcal{Q}(\beta)$. In other words, it suffices to prove that the fiber $\eta_{\beta, w_0, e}^{-1}(x)$ is connected for each $x \in \mathcal{Q}(\beta)$.

By Theorem 4.7, we deduce that the set of connected components of $\eta_{\beta, w_0, e}^{-1}(x)$ is in bijection with the set of connected components of $\prod_{y \in \mathbb{P}^1(\mathbb{C})} \mathcal{B}_{\beta_y}[w]$. By Proposition 4.11, this latter space is connected.

Therefore, we conclude the result. \square

Theorem 4.16. *For each $\beta \in Q_+^\vee$ and $w \in W$, the varieties $\mathcal{Q}(\beta, w_0, w_0)$ and $\mathcal{Q}(\beta, e, w_0)$ are normal.*

Proof. We set $(v, w) = (w_0, w_0)$ or (e, w_0) . We borrow the upper semi-continuous function Θ that counts the number of connected components of the fiber of $\eta_{\beta, v, w}$ from (4.3) in the proof of Proposition 4.13.

By Proposition 4.13, it suffices to prove that $\Theta \equiv 1$ on $\mathcal{Q}(\beta, v, w)$ by assuming the contrary to deduce contradiction. For each $x \in \mathcal{Q}(\beta, v, w)$ such that $\Theta(x) \geq 2$, the fiber $\eta_{\beta, w_0, e}^{-1}(x)$ is disconnected.

By our choice of (v, w) and Theorem 4.7 (cf. Proposition 4.11 and Corollary 4.14), we deduce that the set of connected components of $\eta_{\beta, v, w}^{-1}(x)$ is in bijection with the set of connected components of $\mathcal{B}_{2, \beta_0}[u, w_0]$ or $\mathcal{B}_{2, \beta_0}[u, w_0] \times \mathcal{B}_{2, \beta_\infty}[u', w_0]$ for some $u, u' \in W$.

To see whether this is the case, we specialize to the case of $(v, w) = (w_0, w_0)$ (to guarantee that the contribution at $\infty \in \mathbb{P}^1$ in Theorem 4.7 is the same as the points in $\mathbb{C}^\times \subset \mathbb{P}^1$, that is connected by Proposition 4.11). We can choose $\beta_0 < \beta'_0 \in Q_+^\vee$ such that $\langle \alpha_i^\vee, \beta'_0 \rangle \geq 1$ for every $i \in \mathbf{I}$. By Proposition 4.11, we deduce that the fiber of the locus \mathcal{Z} of $\mathcal{Q}(\beta, w_0)$ whose points have defect β'_0 along 0 is connected (if it is nonempty) for every $\beta \in Q^\vee$. Hence we have $\Theta(y) = 1$ for each $y \in \mathcal{Z}$.

For each $\tilde{x} \in \mathcal{G}\mathcal{B}_{2, \beta}^b$ such that $x = \pi_{2, \beta}(\tilde{x}) \in \mathcal{Q}(\beta, w_0)$ has defect β_0 at 0, we can replace β with $\beta + \beta'_0 - \beta_0$ and add additional irreducible components C' to the main component \mathbb{P}^1 of \tilde{x} (as quasi-stable curves) outside of $0 \in \mathbb{P}^1$ (whose images to \mathcal{B} have their degrees sum up to $(\beta'_0 - \beta_0)$). (This does not change the defect of x at 0, and also does not change $\Theta(x)$.) Then, we shrink all the inserting points of C' on \mathbb{P}^1 to 0 to obtain $\tilde{x}' \in \mathcal{G}\mathcal{B}_{2, \beta}^b$ in the limit, that exists by the valuative criterion of properness as $\mathcal{G}\mathcal{B}_{2, \beta}^b$ is projective. By examining the images of this family on $\mathcal{Q}(\beta, w_0)$ via $\pi_{2, \beta}$, we deduce $y = \pi_{2, \beta}(\tilde{x}') \in \mathcal{Z}$. Therefore, the semi-continuity of Θ implies that $\Theta(x) \leq \Theta(y) = 1$, that is $\Theta(x) = 1$. Hence, $\mathcal{B}_{2, \beta_0}[w, w_0]$ must be connected. This is a contradiction, and we conclude that $\Theta \equiv 1$ (for general $(v, w) \in \{(e, w_0), (w_0, w_0)\}$ by the two paragraphs ahead).

Therefore, Proposition 4.13 implies the result. \square

Corollary 4.17. *Let $\beta \in Q_+^\vee$ and $w \in W$. For each $i \in \mathbf{I}$ such that $s_i w < w$, we have a surjective map*

$$\pi_i : P_i \times^B \mathcal{Q}(\beta, w) \rightarrow \mathcal{Q}(\beta, s_i w)$$

such that $(\pi_i)_ \mathcal{O}_{P_i \times^B \mathcal{Q}(\beta, w)} \cong \mathcal{O}_{\mathcal{Q}(\beta, s_i w)}$ and $\mathbb{R}^{>0}(\pi_i)_* \mathcal{O}_{P_i \times^B \mathcal{Q}(\beta, w)} \cong \{0\}$.*

Proof. Combine Corollary 3.40 with Theorem 3.30 (and take the generic localizations over \mathbb{Z}). \square

Corollary 4.18. *For each $\beta \in Q_+^\vee$ and $w \in W$, the variety $\mathcal{Q}(\beta, w)$ is normal.*

Proof. The case $w = w_0$ is in Theorem 4.16. Assume that the assertion holds for w . Let $i \in \mathbf{I}$ such that $s_i w < w$. Then, Corollary 4.17 implies that $\mathcal{O}_{\mathcal{Q}(\beta, s_i w)}$ is isomorphic to the normal sheaf of rings $(\pi_i)_* \mathcal{O}_{P_i \times^B \mathcal{Q}(\beta, w)}$. Hence, the assertion holds for $s_i w < w$. This proceeds the induction and we conclude the result. \square

Corollary 4.19. *For each $\beta \in Q_+^\vee$ and $w, v \in W$, the subspace*

$$\mathbf{e}_1^{-1}(p_v) \cap \mathbf{e}_2^{-1}(\mathcal{B}(w)) \subset \mathcal{B}_{2, \beta}$$

is connected.

Proof. This space appears in the fiber of $\pi_{\beta, w_0, w}$ along the constant quasimap $\mathbb{P}^1 \rightarrow \{p_v\} \subset \mathcal{B}$ with its defect concentrated in $0 \in \mathbb{P}^1$. Hence, the assertion follows from Corollary 4.18 and Proposition 4.13. \square

Theorem 4.20. *For each $\beta \in Q_+^\vee$ and $w, v \in W$, the scheme $\mathcal{Q}(\beta, v, w)$ is normal.*

Proof. The combination of Theorem 4.7 and Corollary 4.19 implies that every fiber of $\pi_{\beta, v, w}$ is connected. Thus, Proposition 4.13 implies the result. \square

Corollary 4.21. *For each $J \subset I$, $\beta \in Q_+^\vee$ and $w, v \in W$, the variety $\mathcal{Q}_J(\beta, v, w)$ is irreducible and normal.*

Proof. By Corollary 3.35 and Remark 3.36, we can rearrange the map Π_J to be surjective with connected fibers. Hence, Proposition 4.3 implies the irreducibility of $\mathcal{Q}_J(\beta, v, w)$. By the Stain factorization theorem applied to the composition map $\Pi_J \circ \pi_{\beta, v, w}$, we deduce that the normalization of $\mathcal{Q}_J(\beta, v, w)$ is bijective to $\mathcal{Q}_J(\beta, v, w)$. Therefore, the weak normality of $\mathcal{Q}_J(\beta, v, w)$ (Corollary 3.42) implies the result. \square

We set $W^J := W/w_0W_Jw_0$, and identify it with the set of their minimal length representatives in W . Recall that $2\rho_J$ is the sum of the positive roots that belong to the unipotent radical of $\mathfrak{p}(J)$.

Corollary 4.22. *For each $J \subset I$, $\beta \in Q_+^\vee$ such that $-w_0\beta \in Q_{J,+}^\vee$, $w \in W^J$, and $v \in W$ such that $w_0v \in W^J$, the variety $\mathcal{Q}_J(\beta, v, w)$ has dimension*

$$\dim G/P(J) - 2 \langle w_0\beta, \rho_J \rangle + \ell(v) - \ell(w_0) - \ell(w) \quad (4.4)$$

if it is nonempty. In case $v = w_0$, the variety $\mathcal{Q}_J(\beta, v, w)$ is always nonempty. In addition, $\mathcal{Q}_J(\beta, w_0, e)$ is the collection of DP-data as in Definition 1.6 with the data and compatibility conditions only for $\lambda \in P_{J,+}$.

Remark 4.23. If we replace the labels of $\mathcal{Q}_J(\beta, v, w) = \mathcal{Q}'_J(vt_\beta, w)_\mathbb{C}$ with the labels of parabolic semi-infinite Bruhat order as in [61, §10.3], then we can write the condition of $\mathcal{Q}_J(\beta, v, w)$ to be nonempty by the order relation just as in Corollary 3.21.

Proof of Corollary 4.22. By [7], the collection of DP-data in Definition 1.6 with the compatibility condition imposed only for $\lambda \in P_{J,+}$ (without non-defining points) represents the (closure of the) space of maps $\mathbb{P}^1 \rightarrow G/P(J)$ such that the image of $[\mathbb{P}^1]$ in \mathcal{B} is $-w_0\beta \in H_2(\mathcal{B}, \mathbb{Z})$. Thus, we conclude the last assertion.

The dimension of $\mathcal{Q}_J(\beta, w_0, e)$ is given by $\dim G/P(J) - 2 \langle w_0\beta, \rho_J \rangle$ since the first Chern class of the tangent bundle of $G/P(J)$ is $2\rho_J$ (cf. [29, Proposition 3.5]). In particular, the dimension of $\mathcal{Q}_J(\beta, w_0, e)$ is (4.4) in this case.

Since imposing a hyperplane section in the image of $G/P(J)$ through the (generically defined) evaluation map at $0 \in \mathbb{P}^1$ (that is surjective as being G -equivariant)

$$\mathcal{Q}_J(\beta, w_0, e) \dashrightarrow G/P(J)$$

lowers the dimension exactly by one, we deduce that $\mathcal{Q}_J(\beta, w_0, w)$ has an irreducible component of expected dimension. From this and Corollary 4.21, we conclude that $\mathcal{Q}_J(\beta, w_0, w)$ is nonempty and its dimension is given by (4.4).

We consider the dimension estimate in the general case.

Since $\mathcal{Q}_{\mathbf{J}}(\beta, s_i v, w) \subsetneq \mathcal{Q}_{\mathbf{J}}(\beta, v, w)$ ($i \in \mathbf{I}$) implies $s_i v >_{\frac{\infty}{2}} v$ ($\Leftrightarrow s_i v < v$) when $w_0 s_i v \in W^{\mathbf{J}}$, we find that they are the images of $\mathcal{Q}(\beta', s_i v, w) \subsetneq \mathcal{Q}(\beta', v, w)$ for some $\beta' \in \beta + \sum_{j \in \mathbf{J}} \mathbb{Z}_{\geq 0} \alpha_j^{\vee}$ (see Remark 3.36). In particular, $\mathcal{Q}_{\mathbf{J}}(\beta, v, w)$ has dimension at least (4.4) in general by induction from the case $v = w_0$.

Let Z denote the space of genus zero stable maps to $\mathbb{P}^1 \times G/P(\mathbf{J})$ of bidegree $(1, \beta)$ with two marked points such that the first marked points and second marked points correspond to $0, \infty \in \mathbb{P}^1$, respectively (parabolic analogue of $\mathcal{GB}_{2, \beta}^b$). Then, we have a resolution $\pi : Z \rightarrow \mathcal{Q}_{\mathbf{J}}(\beta, w_0, e)$ (that shares genuine maps $\mathbb{P}^1 \rightarrow G/P(\mathbf{J})$ in their common open dense subset). Let $\eta : \mathcal{B} = G/B \rightarrow G/P(\mathbf{J})$ be the projection. By our choices of v, w , we find that

$$\text{codim}_{\mathcal{B}} \mathcal{B}(w) = \text{codim}_{G/P(\mathbf{J})} \eta(\mathcal{B}(w)) \quad \text{and} \quad \text{codim}_{\mathcal{B}} \mathcal{B}^{\text{op}}(v) = \text{codim}_{G/P(\mathbf{J})} \eta(\mathcal{B}^{\text{op}}(v)),$$

respectively. We have the evaluation map of two points $\mathbf{z} : Z \rightarrow (G/P(\mathbf{J}))^2$. We set $Z_{v, w} := \mathbf{z}^{-1}(\eta(\mathcal{B}(w)) \times \eta(\mathcal{B}^{\text{op}}(v)))$. By [16, Propostion 3.2 b)] (see also the proof of Corollary 4.2), we find that

$$\begin{aligned} \dim Z_{v, w} &= \dim Z - \text{codim}_{G/P(\mathbf{J})} \eta(\mathcal{B}(w)) - \text{codim}_{G/P(\mathbf{J})} \eta(\mathcal{B}^{\text{op}}(v)) \\ &= \dim Z - \text{codim}_{\mathcal{B}} \mathcal{B}(w) - \text{codim}_{\mathcal{B}} \mathcal{B}^{\text{op}}(v) = (4.4) \end{aligned}$$

and $Z_{v, w}$ is irreducible (if it is non-empty). Since $G/P(\mathbf{J})$ is homogeneous, we find that $G \cdot \pi(Z_{w_0, w}) \subset Z$ is Zariski dense. Hence, $Z_{w_0, w}$ contains an open dense subset whose points are genuine maps $\mathbb{P}^1 \rightarrow G/P(\mathbf{J})$. In conjunction with the irreducibility of $Z_{w_0, w}$ and $\mathcal{Q}_{\mathbf{J}}(\beta, w_0, w)$, we find $\pi^{-1}(\mathcal{Q}_{\mathbf{J}}(\beta, w_0, w)) = Z_{w_0, w}$. Similarly, we have $\pi^{-1}(\mathcal{Q}_{\mathbf{J}}(\beta, v, e)) = Z_{v, e}$. Therefore, we conclude that $Z_{v, w}$ surjects onto $\mathcal{Q}_{\mathbf{J}}(\beta, v, w)$ by the restriction of π . In particular, $\mathcal{Q}_{\mathbf{J}}(\beta, v, w)$ has dimension at most (4.4) in general.

Therefore, our dimension estimate must be strict in general. \square

Corollary 4.24. *For each $\mathbf{J} \subset \mathbf{I}$ and $w, v \in W_{\text{af}}$, the scheme $\mathcal{Q}'_{\mathbf{J}}(v, w)_{\overline{\mathbb{F}}_p}$ is irreducible and normal for $p \gg 0$.*

Proof. Apply Proposition 3.43 to Corollary 4.21 (cf. Lemma 3.6). \square

Corollary 4.25. *Let \mathbb{K} be an algebraically closed field of characteristic 0 or $p \gg 0$. For each $\mathbf{J} \subset \mathbf{I}$ and $w, v \in W_{\text{af}}$, the scheme $\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}$ is projectively normal with respect to a line bundle $\mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}}(\lambda)$ ($\lambda \in P_{\mathbf{J}, ++}$).*

Proof. In view of Lemma 3.1, Lemma 3.3, and Theorem 3.33, the multiplication of the section ring afforded by $\{\mathcal{O}_{\mathcal{Q}'_{\mathbf{J}}(v, w)_{\mathbb{K}}}(m\lambda)\}_{m \geq 0}$ is surjective. Therefore, Corollary 4.21 and Corollary 4.24 implies the result. \square

Appendix A Some properties of $\mathbf{Q}_G(w)$ in positive characteristic

We work in the setting of §3.2. Let \mathbb{K} be an algebraically closed field of characteristic $\neq 2$. The aim of this appendix is two fold: One is to show that our scheme $\mathbf{Q}_{G, \mathbf{J}}(w)_{\mathbb{K}}$ is (projectively) normal. The other is to present an analogue of the Kempf vanishing theorem for $\mathbf{Q}_G(w)_{\mathbb{K}}$. The both results are proved in [51] for the case $\mathbb{K} = \mathbb{C}$. Our proof of the former is new and does not depend on [51] (when $w = e$), while our proof of the latter depends on the results and arguments in [51] in an essential way.

Proposition A.1. *For each $w \in W_{\text{af}}$ and $\mathbf{J} \subset \mathbf{I}$, the ring $R_w(\mathbf{J})_{\mathbb{K}}$ is normal.*

Proof. We first prove the case $w = e$ and $J = \emptyset$. Let $\mathring{\mathbf{Q}}_G$ denote the open $G[[z]]$ -orbit of $\mathbf{Q}_G(e)_{\mathbb{K}}$ obtained by the G -translation of Lemma 3.11.

We have an inclusion

$$\mathbb{W}(\lambda)_{\mathbb{K}}^{\vee} \subset \Gamma(\mathring{\mathbf{Q}}_G, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda)) \quad (\text{A.1})$$

since $R_{\mathbb{K}}$ is integral (Corollary 3.12). We also have an inclusion

$$\Gamma(\mathring{\mathbf{Q}}_G, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda)) \hookrightarrow \Gamma(\mathbb{O}(e)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda)) \quad \lambda \in P_+.$$

Thanks to Lemma 3.11 and its proof, we deduce

$$\Gamma(\mathbb{O}(e)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda)) \cong \mathbb{K}[\mathbf{I}/(\mathbf{I} \cap HN((z)))] \otimes_{\mathbb{K}} \mathbb{K}_{-w_0\lambda},$$

that is cocyclic as a $U_{\mathbb{K}}^+$ -module. Since the G -action on $\Gamma(\mathring{\mathbf{Q}}_G, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda))$ is algebraic, we deduce that

$$(\mathbb{K}[\mathbf{I}/(\mathbf{I} \cap HN((z)))] \otimes_{\mathbb{K}} \mathbb{K}_{-w_0\lambda})^{\vee} \twoheadrightarrow \Gamma(\mathring{\mathbf{Q}}_G, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda))^{\vee}$$

is a $\mathring{U}_{\mathbb{K}}^0$ -integrable quotient. By Proposition 2.19, we conclude a surjection

$$\mathbb{W}(\lambda)_{\mathbb{K}} \twoheadrightarrow \Gamma(\mathring{\mathbf{Q}}_G, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda))^{\vee}.$$

Compared with (A.1), we conclude the isomorphism

$$\Gamma(\mathring{\mathbf{Q}}_G, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda)) \xrightarrow{\cong} \Gamma(\mathbf{Q}_G(e)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda)) \quad \lambda \in P_+$$

since the space of sections supported on a dense open Zarsiki subset must be larger than (or equal to) the space of sections supported on the whole space for an integral scheme. In other words, $R_{\mathbb{K}}$ is the maximal $\mathring{U}_{\mathbb{K}}^0$ -integrable $U_{\mathbb{K}}^+$ -stable subring of

$$S_{\mathbb{K}} := \bigoplus_{\lambda \in P_+} \Gamma(\mathbb{O}(e)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(e)_{\mathbb{K}}}(\lambda)) = \bigoplus_{\lambda \in P_+} \mathbb{K}[\mathbf{I}/(\mathbf{I} \cap HN((z)))] \otimes_{\mathbb{K}} \mathbb{K}_{-w_0\lambda}.$$

The ring $S_{\mathbb{K}}$ is integrally closed as it is a polynomial ring (of countably many variables). Hence, the integral closure R^+ of the ring $R_{\mathbb{K}}$ is also a subring of $S_{\mathbb{K}}$. In view of the $\mathring{U}_{\mathbb{K}}^0$ -integrability, the ring $R_{\mathbb{K}}$ admits an algebraic G -action. By the canonical nature of the normalization, we deduce that R^+ admits an algebraic G -action (note that we can approximate $R_{\mathbb{K}}$ as a union of Noetherian rings with algebraic G -actions). However, $R_{\mathbb{K}}$ is already the maximal $\mathring{U}_{\mathbb{K}}^0$ -integrable $U_{\mathbb{K}}^+$ -stable subring of $S_{\mathbb{K}}$ from the above. Hence, we have necessarily $R^+ = R_{\mathbb{K}}$. Thus, the case of $w = e$ and $J = \emptyset$ follows.

The case of arbitrary $w \in W_{\text{af}}$ follows from the case of $w = e$ as in [46, §4]. As the fraction field of $R_w(J)_{\mathbb{K}}$ is a subfield of the fraction field of $(R_w)_{\mathbb{K}}$ and $P_{J,+} \subset P_+$ forms (the set of integral points of) a face, we conclude the general case by restriction. \square

Corollary A.2. *The scheme $\mathbf{Q}_{G,J}(w)_{\mathbb{K}}$ is normal.* \square

Corollary A.3. *For each $\lambda \in P_{J,+}$, we have*

$$H^0(\mathbf{Q}_{G,J}(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_{G,J}(w)_{\mathbb{K}}}(\lambda)) = \mathbb{W}_{w w_0}(\lambda)_{\mathbb{K}}^{\vee}.$$

Proof. As finding H^0 from $R_w(J)_{\mathbb{K}}$ can be seen as finding graded pieces of the normalization, the assertion follows from Proposition A.1. \square

Proposition A.4 ([51] Proposition 5.1). *Assume that $\text{char } \mathbb{K} \neq 2$. Let $w \in W$ and $J \subset \mathbf{I}$. Then, an \mathbf{I} -equivariant line bundle on $\mathbf{Q}_{G,J}(w)_{\mathbb{K}}$ is a character twist of $\{\mathcal{O}_{\mathbf{Q}_{G,J}(w)_{\mathbb{K}}}(\lambda)\}_{\lambda \in P_J}$.*

Proof. Taking Corollary 3.20 and Corollary A.2 into account, the proof in [51, Proposition 5.1] works in this setting. \square

Theorem A.5. *Let $w \in W$ and $J \subset I$. For each $\lambda \in P_J$, we have*

$$H^{>0}(\mathbf{Q}_{G,J}(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_{G,J}(w)_{\mathbb{K}}}(\lambda)) = \{0\}.$$

Proof. Assume that $w = e$. We consider the subring defined by

$$T_J := \bigoplus_{\lambda \in P_{J,+}} T_J(\lambda) \subset \bigoplus_{\lambda \in P_{J,+}} \mathbb{W}(\lambda)_{\mathbb{K}}^{\vee} = \bigoplus_{\lambda \in P_{J,+}} \mathbb{W}_{w_0}(\lambda)_{\mathbb{K}}^{\vee} = R_J,$$

where $T_J(\lambda) \subset \mathbb{W}(\lambda)_{\mathbb{K}}^{\vee}$ is the P -weight $(-\lambda)$ -part of $\mathbb{W}(\lambda)_{\mathbb{K}}^{\vee}$. They are generated by the duals of $\tilde{P}_{i,m\delta} \mathbf{v}_{\varpi_i} \in \mathbb{W}(\varpi_i)_{\mathbb{K}}$ ($i \in I \setminus J, m \in \mathbb{Z}_{>0}$; see [5, (3.18)] for the definition of $\tilde{P}_{i,m\delta}$). In view of [5, Proposition 3.18], the ring T_J is a polynomial ring. The actions of $\{\tilde{P}_{i,m\delta}\}_{i \in (I \setminus J), m > 0}$ define endomorphisms of the module $\mathbb{W}(\lambda)_{\mathbb{K}}$ ($\lambda \in P_{J,+}$), each of which is either an injection or zero (cf. the proof of Proposition 2.19). It follows that the action of T_J on R_J is torsion-free.

Thanks to [51, Corollary 4.29], the character comparison forces the torsion-free action of T_J on R_J to be free. Therefore, the proof of [51, Theorem 4.30] works in a verbatim way. This proves the case $w = e$. The general case can be easily deduced from the $w = e$ case as in [51, Corollary 4.31]. \square

Appendix B An application of the Pieri-Chevalley formula

We work in the setting of §3.4. The aim of this appendix is to present a method (Theorem B.6) to describe the global sections of nef line bundles on $\mathcal{Q}'(v, w)_{\mathbb{K}}$ for $v, w \in W_{\text{af}}$ and an algebraically closed field \mathbb{K} of characteristic $\neq 2$.

For each $\mu \in P_+$, we have an extremal weight module $\mathbb{X}(\mu)_{\mathbb{K}}$ and its global base $\mathbf{B}(\mathbb{X}(\mu))$ borrowed from Theorem 2.14.

Lemma B.1. *Let $J \subset I$, and let $\mu \in P_{J,+}$. A subset of $\mathbf{B}(\mathbb{X}(\mu))$ spans*

$$\begin{aligned} H^0(\mathbf{Q}_G(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(w)_{\mathbb{K}}}(\mu))^{\vee} &= H^0(\mathbf{Q}_{G,J}(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_{G,J}(w)_{\mathbb{K}}}(\mu))^{\vee} \subset \mathbb{X}(\mu)_{\mathbb{K}} \quad \text{and} \\ \theta(H^0(\mathbf{Q}_G(vw_0)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(vw_0)_{\mathbb{K}}}(-w_0\mu))^{\vee}) \\ &= \theta(H^0(\mathbf{Q}_{G,\theta(J)}(vw_0)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_{G,\theta(J)}(vw_0)_{\mathbb{K}}}(-w_0\mu))^{\vee}) \subset \mathbb{X}(\mu)_{\mathbb{K}} \end{aligned}$$

for each $w, v \in W_{\text{af}}$.

Proof. Taking Corollary A.3 into account, the assertion follows from Theorem 2.14. \square

Corollary A.3, combined with Theorem 3.33, yields

$$H^0(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\mu))^{\vee} \hookrightarrow H^0(\mathbf{Q}_G(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(w)_{\mathbb{K}}}(\mu))^{\vee} \subset \mathbb{X}(\mu)_{\mathbb{K}} \quad (\text{B.2})$$

for each $w, v \in W_{\text{af}}$.

Lemma B.2. *Let $\mu \in P_{++}$. For each $b \in \mathbf{B}(\mathbb{X}(\mu))$, there exist unique elements $\kappa(b), \iota(b) \in W_{\text{af}}$ with the following properties:*

1. *We have $b \in H^0(\mathbf{Q}_G(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(w)_{\mathbb{K}}}(\mu))^{\vee}$ for $w \in W_{\text{af}}$ if and only if $\kappa(b) \leq_{\infty} w$;*
2. *We have $b \in \theta(H^0(\mathbf{Q}_G(vw_0)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(vw_0)_{\mathbb{K}}}(-w_0\mu))^{\vee})$ for $v \in W_{\text{af}}$ if and only if $\iota(b) \geq_{\infty} v$.*

Proof. In view of Corollary A.3, the first assertion is a rephrasing of Lemma 2.15. The second assertion is obtained from the first assertion in view of Lemma 2.17 and (1.2). \square

Corollary B.3. *The functions κ and ι play the same role as the same named functions in [51, (2.17)] (with opposite convention on the order \leq_{∞}).*

Proof. Compare Lemma B.2 with [51, Theorem 2.8]. \square

Corollary B.4. *Let $\mu \in P_{++}$ and $v, w \in W_{\text{af}}$. The space $H^0(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\mu))^\vee$ is spanned by the subset of $b \in \mathbf{B}(\mathbb{X}(\mu))$ that satisfies $w \geq_{\frac{\infty}{2}} \kappa(b) \geq_{\frac{\infty}{2}} \iota(b) \geq_{\frac{\infty}{2}} v$.*

Proof. This is a rephrasing of Theorem 3.33 for $\mathbf{J} = \emptyset$. \square

Theorem B.5 ([51] Theorem 5.8 and its proof). *Let $w \in W_{\text{af}}$ and $\lambda \in P_+$. There exists a unique collection of elements $a_w^u(\lambda) \in \mathbb{Z}[q^{-1}][H]$ ($u \in W_{\text{af}}$) such that*

$$a_w^w(\lambda) = e^{-w w_0 \lambda}, \quad a_w^u(\lambda) = 0 \quad \text{if } u \not\leq_{\frac{\infty}{2}} w,$$

and

$$\text{gch} \Gamma(\mathbf{Q}_G(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(w)_{\mathbb{K}}}(\lambda + \mu)) = \sum_{u \in W_{\text{af}}} a_w^u(\lambda) \text{gch} \Gamma(\mathbf{Q}_G(u)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(u)_{\mathbb{K}}}(\mu))$$

for every $\mu \in P_{++}$. \square

The goal of this appendix is to prove the following:

Theorem B.6. *Let $w, v \in W_{\text{af}}$ and $\lambda \in P_+$. Let $\{a_w^v(\lambda)\}_{v \in W_{\text{af}}}$ be the collection of elements in Theorem B.5. Then, we have*

$$\text{gch} \Gamma(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\lambda)) = \sum_{u \geq_{\frac{\infty}{2}} v} a_w^u(\lambda).$$

Proof. The proof of the numerical part of Theorem B.5 is [51, Theorem 3.5] (proved in [51, §8.1]) and it counts the elements of $\mathbf{B}(\mathbb{X}(\lambda + \mu))$ that contributes $\Gamma(\mathbf{Q}_G(w)_{\mathbb{K}}, \mathcal{O}_{\mathbf{Q}_G(w)_{\mathbb{K}}}(\lambda + \mu))^\vee = \mathbb{W}_{w w_0}(\lambda + \mu)_{\mathbb{K}}$ in two ways. In particular, we can additionally impose the condition $\iota(\bullet) \geq_{\frac{\infty}{2}} v$ for $\mathbf{B}(\mathbb{X}(\lambda + \mu))$ to deduce that

$$\text{gch} \Gamma(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\lambda + \mu)) = \sum_{u \leq_{\frac{\infty}{2}} v} a_w^u(\lambda) \text{gch} \Gamma(\mathcal{Q}'(u, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(u, w)_{\mathbb{K}}}(\mu))$$

for every $\mu \in P_{++}$ in view of Corollary B.4. The $(H \times \mathbb{G}_m$ -equivariant) Euler characteristic of $\mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\mu)$ ($\mu \in P$) is a rational function on the characters of H , and we can specialize to $\mu = 0$. Now we apply Theorem 3.33 to deduce

$$\chi(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\mu)) = \text{gch} \Gamma(\mathcal{Q}'(v, w)_{\mathbb{K}}, \mathcal{O}_{\mathcal{Q}'(v, w)_{\mathbb{K}}}(\mu))$$

for every $\mu \in P_+$. This implies the desired equality. \square

Remark B.7. **1)** In view of Corollary 3.35 and Remark 3.36, Theorem B.6 describes the space of global sections of $\mathcal{O}_{\mathcal{Q}'_J(v, w)_{\mathbb{K}}}(\lambda)$ for every $w, v \in W_{\text{af}}$, $\mathbf{J} \subset \mathbf{I}$, and $\lambda \in P_{\mathbf{J}, +}$. **2)** In conjunction with Theorem 3.33, Theorem B.6 can be seen as an analogue of Lakshmibai-Littelmann [59, Theorem 34] for semi-infinite flag manifolds. **3)** Thanks to [51, §3], we have a combinatorial rule to express $a_w^u(\lambda)$'s.

Acknowledgments

The author would like to thank Prakash Belkale, Ievgen Makedonskyi, Leonardo Mihalcea, Katsuyuki Naoi, and Daisuke Sagaki for helpful discussions. This work was supported in part by JSPS KAKENHI Grant Number JP26287004 and JP19H01782.

References

- [1] H. H. Andersen, P. Polo, and K. Wen. Representations of quantum algebras. *Invent. Math.*, 104:1–59, 1991.
- [2] S. Arkhipov, R. Bezrukavnikov, A. Braverman, D. Gaitsgory, and I. Mirković. Modules over the small quantum group and semi-infinite flag manifold. *Transform. Groups*, 10(3–4):279–362, 2005.
- [3] Arnaud Beauville and Yves Laszlo. Conformal blocks and generalized theta functions. *Comm. Math. Phys.*, 164(2):385–419, 1994.
- [4] Jonathan Beck, Vyjayanthi Chari, and Andrew Pressley. An algebraic characterization of the affine canonical basis. *Duke Math. J.*, 99(3):455–487, 1999.
- [5] Jonathan Beck and Hiraku Nakajima. Crystal bases and two-sided cells of quantum affine algebras. *Duke Math. J.*, 123(2):335–402, 2004.
- [6] Kai A. Behrend and Yuri I Manin. Stacks of stable maps and Gromov-Witten invariants. *Duke Mathematical Journal*, 85(1):1–60, 1996.
- [7] A. Braverman, M. Finkelberg, D. Gaitsgory, and I. Mirković. Intersection cohomology of Drinfeld’s compactifications. *Selecta Math. (N.S.)*, 8(3):381–418, 2002.
- [8] A. Braverman and D. Gaitsgory. Geometric Eisenstein series. *Invent. Math.*, 150(2):287–384, 2002.
- [9] Alexander Braverman. Spaces of quasi-maps into the flag varieties and their applications. In *International Congress of Mathematicians. Vol. II*, pages 1145–1170. Eur. Math. Soc., Zürich, 2006.
- [10] Alexander Braverman, Boris Feigin, Michael Finkelberg, and Leonid Rybnikov. A finite analog of the AGT relation I: Finite W -algebras and quasimaps’ spaces. *Comm. Math. Phys.*, 308(2):457–478, 2011.
- [11] Alexander Braverman and Michael Finkelberg. Semi-infinite Schubert varieties and quantum K -theory of flag manifolds. *J. Amer. Math. Soc.*, 27(4):1147–1168, 2014.
- [12] Alexander Braverman and Michael Finkelberg. Weyl modules and q -Whittaker functions. *Math. Ann.*, 359(1-2):45–59, 2014.
- [13] Alexander Braverman and Michael Finkelberg. Twisted zastava and q -Whittaker functions. *J. London Math. Soc.*, 96(2):309–325, 2017, arXiv:1410.2365.
- [14] M. Brion and V. Lakshmibai. A geometric approach to standard monomial theory. *Represent. Theory*, 7:651–680, 2003.
- [15] Michel Brion and Shrawan Kumar. *Frobenius splitting methods in geometry and representation theory*, volume 231 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 2005.
- [16] Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, and Nicolas Perrin. Finiteness of cominuscule quantum K -theory. *Ann. Sci. Éc. Norm. Supér. (4)*, 46(3):477–494, 2013.
- [17] Vyjayanthi Chari, Ghislain Fourier, and Tanusree Khandai. A categorical approach to weyl modules. *Transform. Groups*, 15:517–549, 2010.
- [18] Vyjayanthi Chari and Bogdan Ion. BGG reciprocity for current algebras. *Compos. Math.*, 151(7):1265–1287, 2015.
- [19] Vyjayanthi Chari and Andrew Pressley. Weyl modules for classical and quantum affine algebras. *Represent. Theory*, 5:191–223 (electronic), 2001.
- [20] Ivan Cherednik and Syu Kato. Nonsymmetric Rogers-Ramanujan sums and thick Demazure modules. *Adv. Math.*, 374:Article Number 107335, 2020.
- [21] Neil Chriss and Victor Ginzburg. *Representation theory and complex geometry*. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, 2010. Reprint of the 1997 edition.
- [22] Gerd Faltings. Algebraic loop groups and moduli spaces of bundles. *J. Eur. Math. Soc. (JEMS)*, 5(1):41–68, 2003.

- [23] Boris Feigin, Michael Finkelberg, Alexander Kuznetsov, and Ivan Mirković. Semi-infinite flags. II. Local and global intersection cohomology of quasimaps' spaces. In *Differential topology, infinite-dimensional Lie algebras, and applications*, volume 194 of *Amer. Math. Soc. Transl. Ser. 2*, pages 113–148. Amer. Math. Soc., Providence, RI, 1999.
- [24] Boris Feigin and Edward Frenkel. Affine Kac-Moody algebras and semi-infinite flag manifold. *Comm. Math. Phys.*, 128:161–189, 1990.
- [25] Evgeny Feigin, Syu Kato, and Ievgen Makedonskyi. Representation theoretic realization of non-symmetric Macdonald polynomials at infinity. *J. reine angew. Math.*, 764:181–216, 2020. arXiv: 1703.04108.
- [26] Evgeny Feigin and Ievgen Makedonskyi. Vertex algebras and coordinate rings of semi-infinite flags. *Comm. Math. Phys.*, 369(1):221–244, 2019.
- [27] Evgeny Feigin and Ievgen Makedonskyi. Semi-infinite Plücker relations and Weyl modules. *Int. Math. Res. Not. IMRN*, 2020(14):4357–4394, 2020, arXiv:1709.05674.
- [28] Evgeny Feigin, Ievgen Makedonskyi, and Daniel Orr. Generalized Weyl modules and nonsymmetric q -Whittaker functions. *Adv. Math.*, 339:997–1033, 2018.
- [29] Michael Finkelberg and Ivan Mirković. Semi-infinite flags. I. Case of global curve \mathbf{P}^1 . In *Differential topology, infinite-dimensional Lie algebras, and applications*, volume 194 of *Amer. Math. Soc. Transl. Ser. 2*, pages 81–112. Amer. Math. Soc., Providence, RI, 1999.
- [30] William Fulton and Rahul Pandharipande. Notes on stable maps and quantum cohomology. In Robert Lazarsfeld János Kollár and David Morrison, editors, *Algebraic Geometry: Santa Cruz 1995*. Amer. Mathematical Society, 1995.
- [31] Howard Garland. The arithmetic theory of loop algebras. *J. Algebra*, 53:480–551, 1978.
- [32] Alexander Givental. Homological geometry and mirror symmetry. *Proceedings of the International Congress of Mathematicians 1994*, pages 472–480, 1995.
- [33] Alexander Givental and Yuan-Pin Lee. Quantum K -theory on flag manifolds, finite-difference Toda lattices and quantum groups. *Invent. Math.*, 151(1):193–219, 2003.
- [34] Alexander B. Givental. Equivariant Gromov-Witten invariants. *Internat. Math. Res. Notices*, 13:613–663, 1996.
- [35] Ian Grojnowski and George Lusztig. A comparison of bases of quantized enveloping algebras. In *Linear algebraic groups and their representations (Los Angeles, CA, 1992)*, volume 153 of *Contemp. Math.*, pages 11–19. Amer. Math. Soc., Providence, RI, 1993.
- [36] Alexander Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III. *Inst. Hautes Études Sci. Publ. Math.*, (28):255, 1966.
- [37] Robin Hartshorne. *Algebraic geometry.*, volume 52 of *Graduate Text in Mathematics*. Springer-Verlag, 1977.
- [38] N. Iwahori and H. Matsumoto. On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups. *Inst. Hautes Études Sci. Publ. Math.*, (25):5–48, 1965.
- [39] Anthony Joseph. On the Demazure character formula. *Ann. Sci. École Norm. Sup. (4)*, 18(3):389–419, 1985.
- [40] Victor G. Kac. *Infinite-dimensional Lie algebras*. Cambridge University Press, Cambridge, third edition, 1990.
- [41] Masaki Kashiwara. On crystal bases of the q -analogue of universal enveloping algebras. *Duke Mathematical Journal*, 63(2):465–516, 1991.
- [42] Masaki Kashiwara. The crystal base and Littelmann's refined Demazure character formula. *Duke Math. J.*, 71(3):839–858, 1993.
- [43] Masaki Kashiwara. Crystal bases of modified quantized enveloping algebra. *Duke Math. J.*, 73(2):383–413, 1994.
- [44] Masaki Kashiwara. On level-zero representations of quantized affine algebras. *Duke Math. J.*, 112(1):117–175, 2002.
- [45] Masaki Kashiwara. Level zero fundamental representations over quantized affine algebras and Demazure modules. *Publ. Res. Inst. Math. Sci.*, 41(1):223–250, 2005.
- [46] Syu Kato. Demazure character formula for semi-infinite flag varieties. *Math. Ann.*, 371(3):1769–1801, 2018, arXiv:1605.0279.

- [47] Syu Kato. Loop structure on equivariant K -theory of semi-infinite flag manifolds. arXiv:1805.01718v6, 2018.
- [48] Syu Kato. On quantum K -groups of partial flag manifolds. arXiv:1906.09343, 2019.
- [49] Syu Kato. Frobenius splitting of thick flag manifolds of Kac-Moody algebras. *Int. Math. Res. Not. IMRN*, 2020(17):5401–5427, 2020. arXiv:1707.03773.
- [50] Syu Kato and Sergey Loktev. A Weyl module stratification of integrable representations. *Comm. Math. Phys.*, 368:113–141, 2019. arXiv:1712.03508.
- [51] Syu Kato, Satoshi Naito, and Daisuke Sagaki. Equivariant K -theory of semi-infinite flag manifolds and the Pieri-Chevalley formula. *Duke Math. J.*, 169(13):2421–2500, 2020.
- [52] George R. Kempf. Linear systems on homogeneous spaces. *Ann. of Math. (2)*, 103(3):557–591, 1976.
- [53] B. Kim and R. Pandharipande. The connectedness of the moduli space of maps to homogeneous spaces. In *Symplectic geometry and mirror symmetry (Seoul, 2000)*, pages 187–201. World Sci. Publ., River Edge, NJ, 2001.
- [54] Steven L. Kleiman. The transversality of a general translate. *Compositio Math.*, 28(3):287–297, 1974.
- [55] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [56] M. Kontsevich and Yu. Manin. Gromov-Witten classes, quantum cohomology, and enumerative geometry. *Comm. Math. Phys.*, 164(3):525–562, 1994.
- [57] Shrawan Kumar. *Kac-Moody groups, their flag varieties and representation theory*, volume 204 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 2002.
- [58] Shrawan Kumar and Peter Littelmann. Algebraization of Frobenius splitting via quantum groups. *Ann. of Math. (2)*, 155(2):491–551, 2002.
- [59] V. Lakshmibai and P. Littelmann. Richardson varieties and equivariant K -theory. *J. Algebra*, 260(1):230–260, 2003.
- [60] Thomas Lam, Changzheng Li, Leonardo C. Mihalcea, and Mark Shimozono. A conjectural Peterson isomorphism in K -theory. *J. Algebra*, 513:326–343, 2018.
- [61] Thomas Lam and Mark Shimozono. Quantum cohomology of G/P and homology of affine Grassmannian. *Acta Mathematica*, 204(1):49–90, 2010, arXiv:0705.1386v1.
- [62] Cristian Lenart, Satoshi Naito, Daisuke Sagaki, Anne Schilling, and Mark Shimozono. A uniform model for Kirillov-Reshetikhin crystals II. Alcove model, path model, and $P=X$. 2014, arXiv:1402.2203.
- [63] Peter Littelmann. Contracting modules and standard monomial theory for symmetrizable Kac-Moody algebras. *J. Amer. Math. Soc.*, 11(3):551–567, 1998.
- [64] George Lusztig. Hecke algebras and Jantzen’s generic decomposition patterns. *Adv. in Math.*, 37(2):121–164, 1980.
- [65] George Lusztig. Canonical bases in tensor products. *PNAS*, 89(17):8177–8179, 1992.
- [66] George Lusztig. *Introduction to Quantum Groups*, volume 110 of *Progress in Mathematics*. 110. Birkhäuser, 1994.
- [67] George Lusztig. Study of a \mathbf{z} -form of the coordinate ring of a reductive group. *J. Amer. Math. Soc.*, 22:739–769, 2009.
- [68] Mirella Manaresi. Some properties of weakly normal varieties. *Nagoya Math. J.*, 77:61–74, 1980.
- [69] Olivier Mathieu. Formules de caractères pour les algèbres de Kac-Moody générales. *Astérisque*, 159–160:1–267, 1988.
- [70] I. Mirković and K. Vilonen. Geometric Langlands duality and representations of algebraic groups over commutative rings. *Ann. of Math. (2)*, 166(1):95–143, 2007.
- [71] Mircea Mustata. Jet schemes of locally complete intersection canonical singularities. *Inventiones mathematicae*, 424:397–424, 2001.

- [72] Satoshi Naito and Daisuke Sagaki. Demazure submodules of level-zero extremal weight modules and specializations of Macdonald polynomials. *Math. Zeit.*, 283:937–978, 2016, arXiv:1404.2436.
- [73] Katsuyuki Naoi. Tensor products of Kirillov-Reshetikhin modules and fusion products. *Int. Math. Res. Not. IMRN*, 2017(18):5667–5709, 2017.
- [74] Dale Peterson. Quantum cohomology of G/P . Lecture at MIT, 1997.
- [75] Annamalai Ramanathan. Equations defining Schubert varieties and Frobenius splitting of diagonals. *Publ. Math. IHES*, 65:61–90, 1987.
- [76] R. W. Richardson. Intersections of double cosets in algebraic groups. *Indag. Math. (N.S.)*, 3:69–77, 1992.
- [77] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2018.
- [78] Eckart Viehweg. *Quasi-projective moduli for polarized manifolds*, volume 30 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1995.
- [79] Christopher T. Woodward. On D. Peterson’s comparison formula for Gromov-Witten invariants of G/P . *Proc. Amer. Math. Soc.*, 133(6):1601–1609, 2005.
- [80] Hiroshi Yanagihara. Some results on weakly normal ring extension. *J. Math. Soc. Japan*, 35(4):649–661, 1983.
- [81] Xinwen Zhu. An introduction to affine Grassmannians and the geometric Satake equivalence. In *Geometry of moduli spaces and representation theory*. Amer. Math. Soc., Providence, RI, 2017.