

# Kyoto University


Kyoto-Math 2012-07

## *KO*-THEORY OF EXCEPTIONAL FLAG MANIFOLDS

by

Daisuke Kishimoto and Akihiro Ohsita

August 2012



京都大学理学部数学教室  
Department of Mathematics  
Faculty of Science  
Kyoto University  
Kyoto 606-8502, JAPAN

# KO-THEORY OF EXCEPTIONAL FLAG MANIFOLDS

DAISUKE KISHIMOTO AND AKIHIRO OHSITA

ABSTRACT. The  $KO$ -theory of the flag manifold  $G/T$  is determined by calculating the Atiyah-Hirzebruch spectral sequence when  $G$  is one of the exceptional Lie groups  $G_2, F_4, E_6$ , where  $T$  is a maximal torus of  $G$ .

## 1. INTRODUCTION

This work is a continuation of [KH1], [KH2], [KKO] and [K] in which the  $KO$ -theory of various homogeneous spaces are calculated by the Atiyah-Hirzebruch spectral sequence. In [KKO], Kono and the authors calculated the  $KO$ -theory of the classical flag manifolds. Here, we mean by the classical (resp. exceptional) flag manifold the compact classical (resp. exceptional) group divided by its maximal torus. We will denote a maximal torus of a compact, connected Lie group  $G$  by  $T$ . We will calculate the  $KO$ -theory of the exceptional flag manifold  $G/T$  for  $G = G_2, F_4, E_6$ . Recently, a connection between Witt groups and  $KO$ -theory of homogeneous spaces such as Grassmannians and flag manifolds was found [Z], [Y1], [Y2], and so our calculation has applications not only in topology but also in this direction. Our main result is the following.

**Theorem 1.1.** *The  $KO$ -theory of  $G/T$  for  $G = G_2, F_4, E_6$  is given as*

$$KO^{2n-1}(G/T) \cong (\mathbb{Z}/2)^{s_n} \quad \text{and} \quad KO^{2n}(G/T) \cong (\mathbb{Z}/2)^{s_{n+1}} \oplus \mathbb{Z}^t$$

for  $n \in \mathbb{Z}/4$ , where  $t, s_n$  are as in the following table.

$G$	$t$	$s_0$	$s_{-1}$	$s_{-2}$	$s_{-3}$
$G_2$	6	1	2	1	0
$F_4$	576	2	4	6	4
$E_6$	25920	2	4	6	4

The organization of the paper is as follows. In §2, we recall from [KH1] and [KH2] useful lemmas in calculating the Atiyah-Hirzebruch spectral sequence converging to the  $KO$ -theory. We also recall some basic facts on the self-conjugate  $K$ -theory. In §3, we consider the homotopy fiber of a certain cohomology class  $BT^6$  studied in [KI1] and related spaces. Results in this section will be used in calculating the  $KO$ -theory of  $F_4/T$  and  $E_6/T$ . In §4, we determine the  $KO$ -theory of  $G_2/T$ . In §5, we first calculate the  $KO$ -theory of  $F_4/U$  for some maximal rank subgroup  $U$  of  $F_4$ . After this, we determine the  $KO$ -theory of  $F_4/T$ . In §6, we calculate the  $KO$ -theory of  $E_6/T$  in a similar method for  $F_4/T$ .

---

1991 *Mathematics Subject Classification.* Primary 55N15; Secondary 14M15, 55T25.

## 2. ATIYAH-HIRZEBRUCH SPECTRAL SEQUENCE

2.1. **KO-theory.** Recall that the coefficient of  $KO$ -theory is given as

$$KO^* = \mathbb{Z}[\eta, \lambda, \beta, \beta^{-1}] / (2\eta, \eta^3, \eta\lambda, \lambda^2 - 4\beta)$$

for  $|\eta| = -1, |\lambda| = -4, |\beta| = -8$ . Let  $(E_r(X), d_r)$  be the Atiyah-Hirzebruch spectral sequence

$$E_2^{p,q}(X) \cong H^p(X; KO^q) \implies KO^*(X).$$

It is shown in [F] that the second differential  $d_2$  is given as

$$(2.1) \quad d_2^{p,q} = \begin{cases} \text{Sq}^2 \pi_2 & q \equiv 0 \pmod{8} \\ \text{Sq}^2 & q \equiv -1 \pmod{8} \\ 0 & \text{otherwise,} \end{cases}$$

where  $\pi_2$  is the modulo 2 reduction. We now suppose the following condition of a space  $X$ .

$$(2.2) \quad H^{2n}(X; \mathbb{Z}) \text{ is a free abelian group and } H^{2n+1}(X; \mathbb{Z}) = 0 \text{ for } n \geq 0.$$

Then for  $\text{Sq}^2 \text{Sq}^2 = \text{Sq}^3 \text{Sq}^1 = 0$ ,  $(H^*(X; \mathbb{Z}/2), \text{Sq}^2)$  is a chain complex. We denote the cohomology of  $(H^*(X; \mathbb{Z}/2), \text{Sq}^2)$  by  $H^*(X; \text{Sq}^2)$  and call it the  $\text{Sq}^2$ -cohomology of  $X$ . It follows from (2.1) that there is an isomorphism

$$(2.3) \quad \iota : E_3^{p,-1}(X) \xrightarrow{\cong} H^p(X; \text{Sq}^2).$$

The following useful lemma is proved in [KH1] and [KH2].

**Lemma 2.1.** *Let  $X$  be a CW-complex satisfying (2.2). Suppose  $r$  is the smallest integer such that  $d_r \neq 0$  for  $r \geq 3$ . Then the following holds.*

- (1)  $r \equiv 2 \pmod{8}$ .
- (2) *If  $p$  is the smallest integer such that  $d_r^{p,q} \neq 0$ , there exists  $x \in E_r^{p,0}(X)$  satisfying that  $d_r(\eta x) \neq 0$  and  $\iota(\eta x)$  is indecomposable in  $H^p(X; \text{Sq}^2)$ .*
- (3) *Let  $x$  be as in (2). Suppose there is a map  $X \times X \rightarrow X$  by which  $H^*(X; \text{Sq}^2)$  becomes a Hopf algebra. Then  $d_r x$  is primitive in  $H^*(X; \text{Sq}^2)$ .*

Let us consider an extension of  $E_\infty(X)$  to  $KO^*(X)$ .

**Lemma 2.2.** *Let  $X$  be a finite CW-complex satisfying (2.2). Then there exist integers  $s_n, t_n$  for  $n \in \mathbb{Z}/4$  and isomorphisms*

$$KO^{2n-1}(X) \cong (\mathbb{Z}/2)^{s_n} \quad \text{and} \quad KO^{2n}(X) \cong (\mathbb{Z}/2)^{s_{n+1}} \oplus \mathbb{Z}^{t_n}.$$

*Proof.* By assumption, the complex  $K$ -theory  $K^{-1}(X) = 0$ , and by the Atiyah-Hirzebruch spectral sequence  $(E_r(X), d_r)$ , one sees that  $KO^{2n-1}(X)$  is a torsion group. Then since the composite  $KO^*(X) \xrightarrow{\mathbf{c}} K^*(X) \xrightarrow{\mathbf{r}} KO^*(X)$  is the 2-power map for the complexification  $\mathbf{c}$  and

the realization  $\mathbf{r}$ , it follows that  $KO^{2n-1}(X) \cong (\mathbb{Z}/2)^{s_n}$  for some integer  $s_n$ . There is the Bott exact sequence

$$\cdots \rightarrow K^{*-1}(X) \rightarrow KO^{*+1}(X) \xrightarrow{\eta} KO^*(X) \xrightarrow{c} K^*(X) \rightarrow \cdots .$$

Since  $K^0(X)$  is a free abelian group and  $K^{-1}(X) = 0$  by assumption,  $\eta : KO^{2n-1}(X) \rightarrow KO^{2n}(X)$  is an isomorphism on the torsion part. Thus the proof is completed.  $\square$

We calculate integers  $s_n, t_n$  in Lemma 2.2. Define formal series  $f_X(t)$  and  $g_X(t)$  as

$$(2.4) \quad f_X(t) = \sum_{p \geq 0} \dim_{\mathbb{Q}} H^p(X; \mathbb{Q}) t^p \quad \text{and} \quad g_X(t) = \sum_{p \geq 0} \dim_{\mathbb{Z}/2} E_{\infty}^{p,-1}(X) t^p.$$

By [MT], the polynomial  $f_X(t)$  for  $G = G_2/T, F_4/T, E_6/T$  is given as

$$(2.5) \quad f_X(t) = \begin{cases} \frac{(1-t^4)(1-t^{12})}{(1-t^2)^2} & X = G_2/T, \\ \frac{(1-t^4)(1-t^{12})(1-t^{16})(1-t^{24})}{(1-t^2)^4} & X = F_4/T, \\ \frac{(1-t^4)(1-t^{10})(1-t^{12})(1-t^{16})(1-t^{18})(1-t^{24})}{(1-t^2)^6} & X = E_6/T. \end{cases}$$

**Lemma 2.3.** *Let  $X$  be a finite CW-complex satisfying (2.2), and let  $s_n, t_n$  be as in Lemma 2.2. Then it holds that*

$$t_0 = t_{-2} = \frac{f_X(1) + f_X(\sqrt{-1})}{2}, \quad t_{-1} = t_{-3} = \frac{f_X(1) - f_X(\sqrt{-1})}{2}$$

and

$$\begin{pmatrix} s_0 \\ s_{-1} \\ s_{-2} \\ s_{-3} \end{pmatrix} = \frac{1}{4} \begin{pmatrix} 1 & 1 & 2 & 0 \\ 1 & -1 & 0 & -2 \\ 1 & 1 & -2 & 0 \\ 1 & -1 & 0 & 2 \end{pmatrix} \begin{pmatrix} g_X(1) \\ g_X(\sqrt{-1}) \\ \operatorname{Re} g_X\left(\frac{1+\sqrt{-1}}{\sqrt{2}}\right) \\ \operatorname{Im} g_X\left(\frac{1+\sqrt{-1}}{\sqrt{2}}\right) \end{pmatrix}.$$

*Proof.* Since the Atiyah-Hirzebruch spectral sequences for rationalized cohomology theories are trivial, we have

$$t_0 = t_{-2} = \sum_{n \geq 0} \dim_{\mathbb{Q}} H^{4n}(X; \mathbb{Q}) \quad \text{and} \quad t_{-1} = t_{-3} = \sum_{n \geq 0} \dim_{\mathbb{Q}} H^{4n+2}(X; \mathbb{Q}),$$

and then the first two equalities follow. Notice that Lemma 2.2 implies that the extension of  $\bigoplus_{p+q=2n-1} E_{\infty}^{p,q}(X)$  to  $KO^{2n-1}(X)$  is trivial. Then by Bott periodicity and  $E_{\infty}^{p,q}(X) = 0$  for odd  $q$  with  $q \not\equiv -1 \pmod{8}$ , we have

$$KO^{2n-1}(X) \cong \bigoplus_{p+q=2n-1} E_{\infty}^{p,q}(X) \cong \bigoplus_{4k+n \geq 0} E_{\infty}^{8k+2n,-1}(X).$$

On the other hand, we have

$$g_X(t) = \sum_{n=0}^3 \sum_{k \geq 0} \dim_{\mathbb{Z}/2} E_{\infty}^{8k+2n,-1}(X) t^{8k+2n}.$$

Then for  $\omega = \frac{1+\sqrt{-1}}{\sqrt{2}}$ , a primitive 8<sup>th</sup> root of unity, we get

$$g_X(\omega^\ell) = \sum_{n=0}^3 \omega^{2\ell n} s_n = \begin{cases} s_0 + s_{-1} + s_{-2} + s_{-3} & \ell = 0 \\ s_0 - \sqrt{-1}s_{-1} - s_{-2} + \sqrt{-1}s_{-3} & \ell = 1 \\ s_0 - s_{-1} + s_{-2} - s_{-3} & \ell = 2 \end{cases}$$

and thus the last equality follows.  $\square$

**2.2. Self-conjugate  $K$ -theory.** Let us next consider self-conjugate  $K$ -theory. Our basic reference is [A]. We denote the self-conjugate  $K$ -theory of a space  $X$  by  $KSC^*(X)$ . The coefficient of self conjugate  $K$ -theory is periodic by multiplication by a generator of  $KSC^{-4}$ . Moreover, there is an exact sequence

$$\cdots \rightarrow KO^{*+2}(X) \xrightarrow{\eta^2} KO^*(X) \xrightarrow{\mathbf{c}} KSC^*(X) \rightarrow KO^{*+3}(X) \rightarrow \cdots,$$

where  $\mathbf{c}$  is the complexification. Then it follows that

$$KSC^* \cong \begin{cases} \mathbb{Z} & * \equiv 0, -3 \pmod{4} \\ \mathbb{Z}/2 & * \equiv -1 \pmod{4} \\ 0 & * \equiv -2 \pmod{4} \end{cases}$$

and  $\mathbf{c} : KO^* \rightarrow KSC^*$  is an isomorphism for  $* \equiv 0, -1 \pmod{8}$ . Let  $({}'E_r, {}'d_r)$  be the Atiyah-Hirzebruch spectral sequence

$${}'E_2^{p,q} \cong H^p(X; KSC^q) \implies KSC^*(X).$$

**Lemma 2.4.** *Let  $X$  be a CW-complex satisfying (2.2).*

(1) *The complexification*

$$\mathbf{c} : E_3^{p,q}(X) \rightarrow {}'E_3^{p,q}(X)$$

*is an isomorphism for  $q \equiv 0 \pmod{8}$  and a monomorphism for  $q \equiv -1 \pmod{8}$ .*

(2) *If  $r$  is the least integer such that  $'d_r \neq 0$  for  $r \geq 3$ , then*

$$r \equiv 2 \pmod{8} \quad \text{and} \quad {}'d_r^{*,0} \neq 0.$$

*Proof.* (1) This follows from the above observation on  $\mathbf{c} : KO^* \rightarrow KSC^*$ .

(2) Quite similarly to the proof of Lemma 2.1, we see that  $r \equiv 2 \pmod{4}$  and  $'d_r^{*,0} \neq 0$ . By (1), we further see that  $r \equiv 2 \pmod{8}$ , completing the proof.  $\square$

*Remark 2.5.* All results in this section hold if we localized at the prime 2 and will be used in the proof of Theorem 3.7 below.

## 3. KO-THEORY OF A SPACE RELATED WITH A TORUS

In [KI1], the cohomology of  $BT^6$  in connection with the Weyl group action of  $E_6$  is given as

$$H^*(BT^6; \mathbb{Z}) = \mathbb{Z}[t, t_1, \dots, t_6]/(t_1 + \dots + t_6 - 3t), \quad |t| = |t_i| = 2.$$

Generalizing, we may put

$$H^*(BT^N; \mathbb{Z}) = \mathbb{Z}[t, t_1, \dots, t_N]/(t_1 + \dots + t_N - 3t), \quad |t| = |t_i| = 2$$

for  $N \geq 6$ , which respects the above case of  $N = 6$ . Let  $c_i$  be the elementary symmetric function in  $t_1, \dots, t_N$ , and let  $y_4 = c_2 - 4t^2 \in H^4(BT^N; \mathbb{Z})$ . Define  $B\tilde{T}^N$  as the homotopy fiber of

$$y_4 : BT^N \rightarrow K(\mathbb{Z}, 4),$$

where  $B\tilde{T}^6$  is the 4-connective cover of  $BT^6$  in the sense of [KI1]. Let us calculate the mod 2 cohomology of  $B\tilde{T}^N$  following [KI1]. Define  $\bar{c}_{2^i+1} \in \mathbb{Z}/2[t_1, \dots, t_N]$  for  $i \geq 0$  inductively as

$$\bar{c}_2 = c_2 \quad \text{and} \quad \bar{c}_{2^i+1} = \text{Sq}^{2^i} \bar{c}_{2^{i-1}+1}.$$

**Proposition 3.1.** *The mod 2 cohomology of  $B\tilde{T}^N$  is given as*

$$H^*(B\tilde{T}^N; \mathbb{Z}/2) = \mathbb{Z}/2[t_1, \dots, t_N, \gamma_{2^i+1} \mid i \geq 1]/(\bar{c}_{2^i+1} \mid i \geq 0)$$

for  $* \leq 2N$ , where  $|\gamma_{2^i+1}| = 2(2^i + 1)$ .

*Proof.* Let us consider the Serre spectral sequence of a homotopy fiber sequence

$$K(\mathbb{Z}, 3) \rightarrow B\tilde{T}^N \rightarrow BT^N.$$

Recall that the mod 2 cohomology of  $K(\mathbb{Z}, 3)$  is given as

$$H^*(K(\mathbb{Z}, 3); \mathbb{Z}/2) = \mathbb{Z}/2[u_{2^i+1} \mid i \geq 1],$$

where  $u_3$  is the modulo 2 reduction of the fundamental class and  $u_{2^i+1} = \text{Sq}^{2^{i-1}} u_{2^{i-1}+1}$  for  $i \geq 2$ . By definition of  $B\tilde{T}^N$ , the transgression  $\tau$  satisfies  $\tau(u_3) = c_2 (= \bar{c}_2)$  and then  $\tau(u_{2^i+1}) = \bar{c}_{2^i+1}$  for  $i \geq 0$ . Inductively, one sees that  $\bar{c}_{2^i+1}$  includes the term  $c_{2^i+1}$ , implying that  $\{\bar{c}_{2^i+1} \mid 2 \leq 2^i + 1 \leq n\}$  is a regular sequence in  $\mathbb{Z}/2[t_1, \dots, t_N]$ . On the other hand, since  $u_3^2$  is a permanent cycle, there exists  $\gamma_3 \in H^6(B\tilde{T}^N; \mathbb{Z}/2)$  which restricts to  $u_3^2$ . Put

$$\gamma_{2^i+1} = \text{Sq}^{2^i} \gamma_{2^{i-1}+1}$$

for  $i \geq 2$ . By the Cartan formula, we have that  $\gamma_{2^i+1}$  restricts to  $u_{2^i+1}^2$ . Summarizing the above calculation, we obtain the desired result, where we need the condition  $* \leq N$  for regularity of  $\{\bar{c}_{2^i+1} \mid i \geq 0\}$ .  $\square$

There is a sequence of natural maps

$$B\tilde{T}^N \rightarrow B\tilde{T}^{N+1} \rightarrow B\tilde{T}^{N+2} \rightarrow \dots$$

We denote the colimit of this sequence by  $B\tilde{T}^\infty$ . Then by Proposition 3.1, the Milnor exact sequence shows the following. Let  $R$  be a graded algebra over  $\mathbb{Z}/2$  consisting of finite sums of homogeneous formal power series in  $t_1, t_2, \dots$  with  $|t_i| = 2$ .

**Corollary 3.2.** *The mod 2 cohomology  $B\tilde{T}^\infty$  is given as*

$$H^*(B\tilde{T}^\infty; \mathbb{Z}/2) = R \otimes \mathbb{Z}/2[\gamma_{2^i+1} \mid i \geq 1] / (\bar{c}_{2^i+1} \mid i \geq 0).$$

*In particular, for  $n \geq 0$ ,  $H^{2n}(B\tilde{T}^\infty; \mathbb{Z}_{(2)})$  is a free  $\mathbb{Z}_{(2)}$ -module and  $H^{2n+1}(B\tilde{T}^\infty; \mathbb{Z}_{(2)}) = 0$ .*

Let us next calculate the  $\text{Sq}^2$ -cohomology of  $B\tilde{T}^N$  up to a certain dimension. To this end, we recall from [KH1] a special cohomology calculation.

**Lemma 3.3.** *Let  $(A, d)$  be a differential graded algebra over a field.*

- (1) *Suppose that for  $a \in A^n$ ,  $da$  is a non-zero-divisor and  $a^2 = db$  for some  $b \in A^{2n-1}$ . Then it holds that*

$$H^*(A/(da)) \cong \Lambda(a) \otimes H^*(A).$$

- (2) *Suppose that for  $a \in A^n$ ,  $\{a, da\}$  is a regular sequence and  $a^2 = db, b^2 = dc$  for some  $b \in A^{2n-1}, c \in A^{4n-3}$ . Then it holds that*

$$H^*(A/(a, da)) \cong \Lambda(b) \otimes H^*(A).$$

*Proof.* (1) Since  $da$  is a non-zero-divisor, there is a short exact sequence

$$0 \rightarrow A \xrightarrow{\cdot da} A \rightarrow A/(da) \rightarrow 0$$

which induces a long exact sequence

$$\dots \rightarrow H^*(A) \xrightarrow{\cdot H^*(da)} H^{*+n+1}(A) \rightarrow H^{*+n+1}(A/(da)) \xrightarrow{\delta} H^{*+1}(A) \rightarrow \dots,$$

where  $A/(da)$  is, of course, a differential graded algebra. Obviously,  $H^*(da) = 0$  and  $\delta(a) = 1$ . Then it follows that  $H^*(A/(da))$  is a free  $H^*(A)$ -module with a basis  $\{1, a\}$ . Since  $a^2 = db$ , we obtain the desired result.

- (2) Since  $\{a, da\}$  is a regular sequence, there is an exact sequence

$$\dots \rightarrow H^*(A/(da)) \xrightarrow{\cdot H^*(a)} H^{*+n}(A/(da)) \rightarrow H^{*+n}(A/(a, da)) \xrightarrow{\delta} H^{*+1}(A/(da)) \rightarrow \dots$$

as well as above, in which  $\delta(b) = a$ . Since  $H^*(A/(da)) \cong \Lambda(a) \otimes H^*(A)$  by (1), we see that  $H^*(A/(a, da))$  is a free  $H^*(A)$ -module with a basis  $\{1, b\}$ . For  $b^2 = dc$ , the proof is completed.  $\square$

**Proposition 3.4.** *For  $* \leq 2N - 2$ ,*

$$H^*(B\tilde{T}^N; \text{Sq}^2) = \Lambda(x_3, x_7, x_{2^i} \mid i \geq 3), \quad |x_j| = 2j,$$

*where  $N$  can be  $\infty$ .*

*Proof.* Put  $A = \mathbb{Z}/2[t_1, \dots, t_N]$  (or the above  $R$  for  $N = \infty$ ). Notice that since  $A$  is acyclic under  $\text{Sq}^2$ , for any  $x \in A^+$ , there exists  $y \in A$  satisfying  $x^2 = dy$ .

By Lemma 3.3, we have

$$H^*(A/(\bar{c}_2, \bar{c}_3)) = \Lambda(x_3),$$

where  $x_3 = \sum_{i < j} t_i t_j^2$  satisfying  $\text{Sq}^2 x_3 = c_2^2$ . The Adem relation  $\text{Sq}^2 \text{Sq}^{2^i} = \text{Sq}^{2^i+2} + \text{Sq}^{2^i+1} \text{Sq}^1$  implies that

$$(3.1) \quad \text{Sq}^2 \bar{c}_{2^i+1} = \bar{c}_{2^{i-1}+1}^2$$

for  $i \geq 2$ . On the other hand, as is noted in the proof of Proposition 3.1,  $\{\bar{c}_{2^i+1} \mid 2 \leq 2^i+1 \leq N\}$  is a regular sequence in  $A$ . Then, applying Lemma 3.3 repeatedly, one gets

$$H^*(A/(\bar{c}_{2^i+1} \mid i \geq 0)) = \Lambda(x_3, x_{2^i} \mid i \geq 2)$$

for  $* \leq 2N$ , where  $\text{Sq}^2 x_{2^i} \equiv \bar{c}_{2^i+1} \pmod{(\bar{c}_{2^j+1} \mid 0 \leq j \leq i-1)}$ . Notice here that since  $H^{2(2^{i+1}+1)}(A/(\bar{c}_{2^j+1} \mid j \geq 0)) = 0$ , we can apply Lemma 3.3 repeatedly. Since  $\text{Sq}^2 c_4 = \bar{c}_5 \pmod{(\bar{c}_2, \bar{c}_3)}$ , we may take  $x_4 = c_4$ .

Put  $F_0 = A/(\bar{c}_{2^i+1} \mid i \geq 0)$  and  $F_n = A/(\bar{c}_{2^i+1} \mid i \geq 0) \otimes \mathbb{Z}/2[\gamma_{2^i+1} \mid i \leq n-1]$  for  $n \geq 1$ . It is proved in [KI1] that  $\text{Sq}^2 \gamma_3 = c_4$ . Consider the spectral sequence associated with a filtration  $F_0 \subset F_1$ . Then we get

$$H^*(F_1) = \Lambda(x_3, x_7, x_{2^i} \mid i \geq 3) \otimes \mathbb{Z}/2[\gamma_3^2],$$

where  $x_7 = \gamma_3 c_4 + d_7$  for  $d_7 \in A$  with  $\text{Sq}^2 d_7 = c_4^2$ . Similarly to (3.1), we have  $\text{Sq}^2 \gamma_{2^i+1} = \gamma_{2^{i-1}+1}$ . Then by considering the spectral sequence associated with a filtration  $F_n \subset F_{n+1}$  for  $n \geq 1$  inductively, we obtain

$$H^*(F_{n+1}) = \Lambda(x_3, x_7, x_{2^i} \mid i \geq 3) \otimes \mathbb{Z}/2[\gamma_{2^n+1}^2].$$

Thus the proof is completed.  $\square$

Let us next consider the homotopy fiber  $F$  of the cohomology class  $t : B\tilde{T}^\infty \rightarrow K(\mathbb{Z}, 2)$ . Let  $\alpha : F \rightarrow B\tilde{T}^\infty$  be the natural map.

**Proposition 3.5.** *For  $n \geq 0$ ,  $H^{2n}(F; \mathbb{Z}_{(2)})$  is a free  $\mathbb{Z}_{(2)}$ -module and  $H^{2n+1}(F; \mathbb{Z}_{(2)}) = 0$ .*

*Proof.* By Proposition 3.1, for  $* \leq 2N$ , the same claim is true for  $B\tilde{T}^N$ , and then also for  $B\tilde{T}^\infty$  by sending  $N$  to  $\infty$ . Since the map  $t : B\tilde{T}^\infty \rightarrow K(\mathbb{Z}, 2)$  is injective in the  $\mathbb{Z}_{(2)}$ -cohomology,  $\alpha^* : H^*(B\tilde{T}^\infty; \mathbb{Z}_{(2)}) \rightarrow H^*(F; \mathbb{Z}_{(2)})$  is surjective, and thus the proof is completed.  $\square$

Define a map  $\mu : BT^\infty \times BT^\infty \rightarrow BT^\infty$  by the equation

$$\mu^*(t_{2i}) = 1 \otimes t_i \quad \text{and} \quad \mu^*(t_{2i-1}) = t_i \otimes 1$$

for  $i \geq 1$  in cohomology. Then by an easy inspection we see that  $\mu$  lifts to a map  $\tilde{\mu} : F \times F \rightarrow F$ .



**Proposition 3.6.** *The natural map  $\alpha : F \rightarrow B\tilde{T}^\infty$  induces an isomorphism in the  $\text{Sq}^2$ -cohomology. Moreover,  $H^*(F; \text{Sq}^2)$  becomes a Hopf algebra by  $\tilde{\mu}$  in which  $\alpha^*(x_{2^i})$  is not primitive for  $i \geq 4$ , where  $x_j$  is as in Proposition 3.4.*

*Proof.* The first assertion easily follows from a direct calculation.

Computing the  $\text{Sq}^2$ -cohomology of the subring  $\mathbb{Z}/2[c_1, c_2, c_3, \dots]/(c_1, \bar{c}_2, \bar{c}_3, \dots)$  of  $H^*(F; \mathbb{Z}/2)$ , we see that  $\alpha^*(x_{2^i})$  can be chosen as an element of this subring for  $i \geq 3$ . Then for

$$(3.2) \quad \tilde{\mu}^*(\alpha^*(c_n)) = \sum_{i=0}^n \alpha^*(c_i) \otimes \alpha^*(c_{n-i}),$$

we obtain

$$\tilde{\mu}^*(\alpha^*(x_{2^i})) = \alpha^*(x_{2^i}) \otimes 1 + 1 \otimes \alpha^*(x_{2^i}) + \cdots.$$

Choose representatives of  $x_3, x_7$  as in the proof of Proposition 3.4. As in [KKO], it is straightforward to see that  $\tilde{\mu}^*(\alpha^*(x_3)) = x_3 \otimes 1 + 1 \otimes x_3$ . By definition, we have  $\tilde{\mu}^*(\alpha^*(\gamma_3)) = \alpha^*(\gamma_3) \otimes 1 + 1 \otimes \alpha^*(\gamma_3) + \cdots$ . Then by an easy calculation analogous to  $\alpha^*(x_3)$ , we see that  $\tilde{\mu}^*(\alpha^*(x_7)) = \alpha^*(x_7) \otimes 1 + 1 \otimes \alpha^*(x_7)$ . Thus we have obtained that  $H^*(F; \text{Sq}^2)$  is a Hopf algebra by the map  $\tilde{\mu}$ .

Since  $\bar{c}_{2^{i+1}} = c_{2^{i+1}} + \cdots$  as above, we have  $x_{2^i} = c_{2^i} + \cdots$  for  $i \geq 3$ . Then by (3.2), the last assertion follows.  $\square$

We now aim at proving the following.

**Theorem 3.7.** *The Atiyah-Hirzebruch spectral sequence  $E_r(B\tilde{T}^\infty)_{(2)}$  collapses at  $E_3$ -term.*

*Proof.* By Corollary 3.2,  $B\tilde{T}^\infty$  satisfies the condition (2.2) at the prime 2. Let  $\bar{x}_j$  be an element of  $\text{Ker}\{\text{Sq}^2 : H^*(B\tilde{T}^\infty; \mathbb{Z}_{(2)}) \rightarrow H^*(B\tilde{T}^\infty; \mathbb{Z}/2)\} \cong E_3^{*,0}(B\tilde{T}^\infty)_{(2)}$  whose modulo 2 reduction is  $x_j \in H^*(B\tilde{T}^\infty; \text{Sq}^2)$  for  $j = 3, 7, 2^i$  ( $i \geq 3$ ). Then by Lemma 2.1, our aim is to prove that  $\bar{x}_j$  is a permanent cycle for  $j = 3, 7, 2^i$  ( $i \geq 3$ ).

Consider the natural map  $\alpha : F \rightarrow B\tilde{T}^\infty$ . Then it follows from Lemma 2.1, Proposition 3.5 and Proposition 3.6 that it is sufficient to show that  $\alpha^*(\bar{x}_3) \in \text{Ker}\{\text{Sq}^2 : H^*(F; \mathbb{Z}_{(2)}) \rightarrow H^*(F; \mathbb{Z}/2)\} \cong E_3^{*,0}(F)_{(2)}$  is a permanent cycle. We next consider the complexification  $\mathbf{c} : E_r(F)_{(2)} \rightarrow {}'E_r(F)_{(2)}$ . Then by Lemma 2.4, we only have to prove  $\mathbf{c}(\alpha^*(\bar{x}_3)) \in {}'E_3(F)_{(2)}$  is a permanent cycle.

Let  $u$  be a generator of  $K_{(2)}^{-2}$  satisfying  $(1-\mathbf{t})(u) = 0$  for the complex conjugation  $\mathbf{t}$ , and let  $H_i$  be the pullback of the Hopf bundle on  $BT^1$  by the composite  $F \rightarrow BT^\infty \rightarrow BT^1$  in which the first arrow is the natural map and the second arrow corresponds to the cohomology class  $t_i$ . Put  $\xi_3 = u^{-3} \sum_{i < j} H_i H_j^2 \in K^6(B\tilde{T}^\infty)_{(2)}$ . Then for  $(1-\mathbf{t})(\xi_3) = 0$ ,  $\xi_3$  lies in  $KSC^6(F)_{(2)}$ . Obviously,  $\xi_3$  corresponds to  $\mathbf{c}(\alpha^*(\bar{x}_3))$ , and thus  $\mathbf{c}(\alpha^*(\bar{x}_3))$  is a permanent cycle as is desired.  $\square$

4. KO-THEORY OF  $G_2/T$ 

The mod 2 cohomology of  $G_2/T$  including the action of the Steenrod operations is calculated as

$$H^*(G_2/T; \mathbb{Z}/2) = \mathbb{Z}/2[t_1, t_2, \gamma_3]/(\rho_2, \rho_3, \gamma_3^2), \quad |t_i| = 2, |\gamma_3| = 6, \quad \text{Sq}^2 \gamma_3 = 0,$$

where

$$\rho_2 = t_1^2 + t_1 t_2 + t_2^2 \quad \text{and} \quad \rho_3 = t_1^2 t_2 + t_1 t_2^2.$$

**Proposition 4.1.** *The  $\text{Sq}^2$ -cohomology of  $G_2/T$  is given as*

$$H^*(G_2/T; \text{Sq}^2) = \Lambda(x_3, \gamma_3),$$

where  $x_3 = t_1^3 + t_1 t_2^2 + t_2^3$ .

*Proof.* Since  $\text{Sq}^2 \rho_2 = \rho_3$ , we obtain the desired result by Lemma 3.3. □

**Corollary 4.2.** *The Atiyah-Hirzebruch spectral sequence  $E_r(G_2/T)$  collapses at  $E_3$ -term. In particular, we have*


$$g_{G_2/T}(t) = (1 + t^6)^2.$$

*Proof.* The result follows from Lemma 2.1 and Proposition 4.1. □

*Proof of Theorem 1.1 for  $G_2$ .* The result follows from (2.5), Lemma 2.2 (1) and Corollary 4.2. □

 5. KO-THEORY OF  $F_4/T$ 

Recall that the Dynkin diagram of  $F_4$  is given as follows.



It is shown in [IT] that the centralizer of the circle in  $F_4$  defined by  $\alpha_2 = \alpha_3 = \alpha_4 = 0$  is isomorphic to  $T^1 \cdot \text{Sp}(3)$ . Let  $U$  be the centralizer of the torus defined by  $\alpha_2 = 0$ . Then  $U \cong T^3 \times \text{Sp}(1)$  as a space, implying that the homology of  $U$  is torsion free. Note that  $F_4/U$  satisfies the condition (2.2). Then we calculate the Atiyah-Hirzebruch spectral sequence converging to  $KO^*(F_4/U)$  from which we deduce the one converging to  $KO^*(F_4/T)$ .

**5.1. KO-theory of  $F_4/U$ .** We first calculate the mod 2 cohomology of  $F_4/U$ . Let  $\omega_i$  ( $i = 1, 2, 3, 4$ ) be the fundamental weight of  $F_4$  as in [TW], and put

$$t = \omega_1, \quad y_1 = \omega_2 - \omega_3, \quad y_2 = \omega_3 - \omega_4, \quad y_4 = \omega_4.$$

Then it is clear that

$$H^*(BT; \mathbb{Z}) = \mathbb{Z}[t, y_1, y_2, y_3].$$

As in [IT], the Weyl group of  $U$  is generated by a single element  $R$  satisfying

$$R(t) = t, \quad R(y_1) = t - y_1, \quad R(y_2) = y_2, \quad R(y_3) = y_3.$$

Since  $H^*(BU; \mathbb{Z})$  is torsion free as noted above,  $H^*(BU; \mathbb{Z})$  is the invariant ring of  $H^*(BT; \mathbb{Z})$  under the action of the Weyl group of  $U$ . Then one gets

$$H^*(BU; \mathbb{Z}) = \mathbb{Z}[t, y_2, y_3, q], \quad q = y_1(t - y_1).$$

On the other hand, the mod 2 cohomology of  $F_4$  is given as

$$H^*(F_4; \mathbb{Z}/2) = \mathbb{Z}/2[a_3]/(a_3^4) \otimes \Lambda(a_5, a_{15}, a_{23}), \quad |a_i| = i, \quad \beta a_5 = a_3^2.$$

Then by a result of Toda [T], we can calculate the  $\mathbb{Z}_{(2)}$ -coefficient cohomology of  $F_4/U$  as follows.

**Proposition 5.1.** *There is a regular sequence  $\bar{\rho}_2, \bar{\rho}_6, \bar{\rho}_8, \bar{\rho}_{12}$  in  $\mathbb{Z}_{(2)}[t, y_2, y_3, q]$  with  $|\bar{\rho}_i| = 2i$  such that*

$$H^*(F_4/U; \mathbb{Z}_{(2)}) = \mathbb{Z}_{(2)}[t, y_2, y_3, q, \gamma_3]/(\bar{\rho}_2, \bar{\rho}_6, \bar{\rho}_8, \bar{\rho}_{12}, 2\gamma_3 + \bar{\rho}_3),$$

where  $\bar{\rho}_3$  is defined by the equation  $\text{Sq}^2 \bar{\rho}_2 = \bar{\rho}_3$ .

We now determine the mod 2 cohomology of  $F_4/U$ . Define  $q_i \in \mathbb{Z}[t, y_2, y_3, q]$  ( $|q_i| = 4i$ ) as

$$1 + q_1 + q_2 + q_3 = (1 + q)(1 + y_2(t - y_2))(1 + y_3(t - y_3)).$$

By definition, one has

$$(5.1) \quad \text{Sq}^2 q_1 = tq_1, \quad \text{Sq}^2 q_2 = 0, \quad \text{Sq}^2 q_3 = tq_3.$$

A calculation in [IT] implies that the rational cohomology of  $F_4/U$  is given as

$$(5.2) \quad H^*(F_4/U; \mathbb{Q}) = \mathbb{Q}[t, y_2, y_3, q]/(\sigma_2, \sigma_6, \sigma_8, \sigma_{12}),$$

where

$$(5.3) \quad \sigma_2 = -t^2 + q_1, \quad \sigma_6 = -t^6 + 4t^2 q_2 - 8q_3, \quad \sigma_8 = 3t^2 q_3 - q_2^2, \quad \sigma_{12} = -q_2^3 + 27q_3^2.$$

Let  $\bar{\rho}_i$  ( $i = 2, 6, 8, 12$ ) be as in Proposition 5.1. Then by (5.1) and (5.3), we may put

$$\bar{\rho}_2 = -t^2 + q_1 \quad \text{and} \quad \bar{\rho}_3 = tq_1.$$

Put

$$R = \mathbb{Z}_{(2)}[t, y_2, y_3, q, \gamma_3]/(\bar{\rho}_2, \bar{\rho}_3, -\gamma_3^2 + t^2 q_2 - 2q_3, \sigma_8, \sigma_{12}).$$

Since  $\sigma_6 \equiv 4(-\gamma_3^2 + t^2 q_2 - 2q_3) \pmod{(\bar{\rho}_2, \bar{\rho}_3)}$  and the natural map  $H^*(F_4/U; \mathbb{Z}_{(2)}) \rightarrow H^*(F_4/U; \mathbb{Q})$  is injective, there is a surjection  $R \rightarrow H^*(F_4/U; \mathbb{Z}_{(2)})$  which induces a surjection

$$\phi : R/2 \rightarrow H^*(F_4/U; \mathbb{Z}/2).$$

We now put

$$(5.4) \quad \rho_2 = t^2 + q_1, \quad \rho_3 = tq_1, \quad \rho_6 = \gamma_3^2 + t^2 q_2, \quad \rho_8 = t^2 q_3 + q_2^2, \quad \rho_{12} = q_2^3 + q_3^2.$$

Then since the Poincaré series of  $F_4/U$  over  $\mathbb{Q}$  and  $\mathbb{Z}/2$  are the same, we have

$$R/2 = \mathbb{Z}/2[t, y_2, y_3, q, \gamma_3]/(\rho_2, \rho_3, \rho_6, \rho_8, \rho_{12}),$$

here in the Poincaré series,  $\gamma_3$  is cancelled by  $\rho_3$ . One can easily verify that  $\rho_2, \rho_3, \rho_6, \rho_8, \rho_{12}$  is a regular sequence in  $\mathbb{Z}/2[t, y_2, y_3, q, \gamma_3]$ , implying that the Poincaré series of  $R/2$  is  $\frac{(1-t^{12})(1-t^{16})(1-t^{24})}{(1-t^2)^3}$ . On the other hand, the Poincaré series of  $H^*(F_4/U; \mathbb{Z}/2)$  is equal to that of  $H^*(F_4/U; \mathbb{Q})$  which is  $\frac{(1-t^{12})(1-t^{16})(1-t^{24})}{(1-t^2)^3}$  by (5.2). Then we conclude that Poincaré series of  $R/2$  and  $H^*(F_4/U; \mathbb{Z}/2)$  are the same, and thus the map  $\phi$  is an isomorphism. Summarizing, we obtain the following.

**Proposition 5.2.** *The mod 2 cohomology of  $F_4/U$  is given as*

$$H^*(F_4/U; \mathbb{Z}/2) = \mathbb{Z}/2[t, y_2, y_3, q, \gamma_3]/(\rho_2, \rho_3, \rho_6, \rho_8, \rho_{12}),$$

where  $|t| = |y_2| = |y_3| = 2, |q| = 4, |\gamma_3| = 6$  and  $\rho_i$  is as in (5.4).

**Corollary 5.3.** *The  $\text{Sq}^2$ -cohomology of  $F_4/U$  is given as*

$$H^*(F_4/U; \text{Sq}^2) = \Lambda(x_7, x_{11}, \bar{\gamma}_3), \quad |x_i| = 2i, |\bar{\gamma}_3| = 6,$$

where  $\text{Sq}^2 x_7 \equiv \rho_8 \pmod{(\rho_2, \rho_3)}$ ,  $\text{Sq}^2 x_{11} = \rho_{12}$ ,  $\bar{\gamma}_3 = \gamma_3 + \delta_3$  and  $\text{Sq}^2 \delta_3 = q_2$  for  $\delta_3 \in \mathbb{Z}/2[t, y_2, y_3, q]$ .

*Proof.* Considering the projection  $F_4/T \rightarrow F_4/U$ , one sees from [KI2] that

$$\text{Sq}^2 \gamma_3 = q_2.$$

Let  $A$  be a differential graded algebra  $\mathbb{Z}/2[t, y_2, y_3, q]$  with  $|t| = |y_i| = 2, |q| = 4$  and  $dt = t^2, dy_i = y_i^2, dq = tq$ , where the degree of the differential is 2. Then by Proposition 5.2, our aim is to determine the cohomology of a differential graded algebra

$$A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_6, \rho_8, \rho_{12}),$$

where  $|\gamma_3| = 6, d\gamma_3 = q_2$  and  $\rho_i$  is as in (5.4). By definition, we have

$$A/(\rho_2, \rho_3) = \mathbb{Z}/2[y_2, y_3] \otimes \langle 1, t, t^2 \rangle$$

as a  $\mathbb{Z}/2[y_2, y_3]$ -module, and then  $H^*(A/(\rho_2, \rho_3)) = 0$ . Hence for  $d\rho_8 \equiv 0 \pmod{(\rho_2, \rho_3)}$  and  $d\rho_{12} = 0$ , it follows from (3.3) that

$$H^*(A/(\rho_2, \rho_3, \rho_8, \rho_{12})) = \Lambda(x_7, x_{11}), \quad |x_i| = 2i.$$

Since  $dq_2 = 0$  and  $H^*(A) = 0$ , there exists  $\delta_3 \in H^6(A)$  satisfying  $d\delta_3 = q_2$ . Put  $\bar{\gamma}_3 = \gamma_3 + \delta_3$ . Then one has

$$A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_8, \rho_{12}) = A \otimes \mathbb{Z}/2[\bar{\gamma}_3]/(\rho_2, \rho_3, \rho_8, \rho_{12})$$

and  $\rho_6 \equiv \bar{\gamma}_3^2 + d(t^2 \delta_3 + \delta_5) \pmod{(\rho_2, \rho_3)}$ , where  $\delta_5 \in H^{10}(A)$  is given by  $d\delta_5 = \delta_3^2$ . Thus for  $d\bar{\gamma}_3 = 0$ , we obtain

$$H^*(A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_6, \rho_8, \rho_{12})) = \Lambda(x_7, x_{11}, \bar{\gamma}_3),$$

completing the proof.  $\square$

**Theorem 5.4.** *The Atiyah-Hirzebruch spectral sequence  $E_r(F_4/U)$  collapses at  $E_3$ -term. In particular, we have*

$$g_{F_4/U}(t) = (1 + t^6)(1 + t^{14})(1 + t^{22}).$$

*Proof.* The result follows from Lemma 2.1 (1), (2) and Corollary 5.3.  $\square$

**Theorem 5.5.** *The  $KO$ -theory of  $F_4/U$  is given as*

$$KO^{2n-1}(F_4/U) \cong (\mathbb{Z}/2)^{s_n} \quad \text{and} \quad KO^{2n}(F_4/U) \cong (\mathbb{Z}/2)^{s_{n+1}} \oplus \mathbb{Z}^t$$

for  $n \in \mathbb{Z}/4$ , where

$$t = 144, \quad s_0 = s_{-3} = 1, \quad s_{-1} = s_{-2} = 3.$$

*Proof.* As is noted above, we have  $f_{F_4/U}(t) = \frac{(1-t^{12})(1-t^{16})(1-t^{24})}{(1-t^2)^3}$ . Then the proof is completed by Lemma 2.2, 2.3 and Theorem 5.4.  $\square$

**5.2.  $KO$ -theory of  $F_4/T$ .** Let  $\rho_i \in \mathbb{Z}/2[t, y_1, y_2, y_3, \gamma_3]$  be as in (5.4), where  $q = y_1(t - y_1)$ . In [KI2], the mod 2 cohomology of  $F_4/T$  is calculated as

$$H^*(F_4/T; \mathbb{Z}/2) = \mathbb{Z}/2[t, y_1, y_2, y_3, \gamma_3]/(\rho_2, \rho_3, \rho_6, \rho_8, \rho_{12})$$

and  $\text{Sq}^2 \gamma_3 = q_2$ . Then the induced map from the projection  $\pi : F_4/T \rightarrow F_4/U$  in the mod 2 cohomology satisfies

$$(5.5) \quad \pi^*(t) = t \quad \text{and} \quad \pi^*(y_i) = y_i \quad (i = 1, 2, 3).$$

Define a map  $\lambda : F_4/T \rightarrow BT^6$  by  $\lambda^*(t_i) = t - y_{4-i}$  and  $\lambda^*(t_{i+3}) = y_i$  for  $i = 1, 2, 3$ . Then  $\lambda^*(c_2 - 4t^2) = -t^2 + q_1 = 0$ , implying that there is a lift  $\tilde{\lambda} : F_4/T \rightarrow B\tilde{T}^6$  satisfying

$$(5.6) \quad \tilde{\lambda}^*(t_i) = t - y_{4-i}, \quad \tilde{\lambda}^*(t_{i+3}) = y_i \quad (i = 1, 2, 3) \quad \text{and} \quad \tilde{\lambda}^*(\gamma_3) = \gamma_3,$$

where the last equality is shown in [KI2].

**Proposition 5.6.** *The  $\text{Sq}^2$ -cohomology of  $F_4/T$  is given as*

$$H^*(F_4/T; \text{Sq}^2) = \Lambda(x_3, x_7, x_{11}, \bar{\gamma}_3), \quad |x_i| = 2i, |\bar{\gamma}_3| = 6,$$

where  $\tilde{\lambda}^*(x_3) = x_3$ ,  $\pi^*(x_7) = x_7$ ,  $\pi^*(x_{11}) = x_{11}$  and  $\pi^*(\bar{\gamma}_3) = \bar{\gamma}_3$ .

*Proof.* Let  $A$  be a differential graded algebra  $\mathbb{Z}/2[t, y_1, y_2, y_3]$  with  $|t| = |y_i| = 2$  and  $dt = t^2, dy_i = y_i^2$ . Then the desired  $\text{Sq}^2$ -cohomology is equal to the cohomology of

$$A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_6, \rho_8, \rho_{12}),$$

where  $d\gamma_3 = q_2$ . Since  $H^*(A) = 0$ ,  $d\rho_2 = \rho_3$ ,  $d\rho_8 \equiv 0 \pmod{(\rho_2, \rho_3)}$  and  $d\rho_{12} = 0$ , it follows from Lemma 3.3 that

$$H^*(A/(\rho_2, \rho_3, \rho_8, \rho_{12})) = \Lambda(x_3, x_7, x_{11}),$$

where  $dx_3 = q_2$  and  $x_7, x_{11}$  are as in Proposition 5.3. Then by defining  $\bar{\gamma}_3$  as in the proof of Proposition 5.3, the first assertion follows. The second assertion follows from (5.5) and (5.6).  $\square$

*Remark 5.7.* Since  $H^*(F_4/T; \text{Sq}^2)$  is an exterior algebra generated by four generators of degree  $-2 \pmod 8$  as in Proposition 5.6, we cannot directly see that  $E_r(F_4/T)$  collapses at  $E_3$ -term by Lemma 2.1. On the other hand,  $H^*(F_4/U; \text{Sq}^2)$  can be thought as a subalgebra of  $H^*(F_4/T; \text{Sq}^2)$  generated by three of its four generators, and then we can apply Lemma 2.1 to see that  $E_r(F_4/U)$  collapses at  $E_3$ -term as above.

**Theorem 5.8.** *The Atiyah-Hirzebruch spectral sequence  $E_r(F_4/T)$  collapses at  $E_3$ -term. In particular, we have*

$$g_{F_4/T}(t) = (1 + t^6)^2(1 + t^{14})(1 + t^{22}).$$


*Proof.* By Theorem 3.7 and Proposition 5.6,  $\iota^{-1}(x_3)$  in the 2-localized spectral sequence  $E_3^{6,-1}(F_4/T)_{(2)}$  is a permanent cycle. Then since the 2-localization  $E_3^{p,q}(F_4/T) \rightarrow E_3^{p,q}(F_4/T)_{(2)}$  is injective,  $\iota^{-1}(x_3)$  in the integral spectral sequence  $E_3^{6,-1}(F_4/T)$  is also a permanent cycle. By Theorem 5.4 and Proposition 5.6,  $\iota^{-1}(x_7), \iota^{-1}(x_{11}), \iota^{-1}(\bar{\gamma}_3) \in E_3^{*, -1}(F_4/T)$  are also permanent cycles. Thus the proof is completed by Lemma 2.1 (2).  $\square$

*Proof of Theorem 1.1 for  $F_4$ .* The result follows from (2.5), Lemma 2.2 and Corollary 5.8.  $\square$

## 6. KO-THEORY OF $E_6/T$

Our method of computing the Atiyah-Hirzebruch spectral sequence  $E_r(E_6/T)$  is similar to the case of  $F_4/T$ . Namely, we first calculate the Atiyah-Hirzebruch spectral sequence converging to  $KO^*(E_6/U)$  for an appropriate maximal rank subgroup  $U$  and then deduce that of  $KO^*(E_6/T)$ .

We know that the Dynkin diagram of  $E_6$  is given as follows.



In [IT], it is proved that the centralizer of the circle in  $E_6$  defined by  $\alpha_1 = \alpha_3 = \alpha_4 = \alpha_5 = \alpha_6 = 0$  is isomorphic to  $T^1 \cdot \text{SU}(6)$ . Then the identity component of the centralizer of the torus defined by  $\alpha_5 = \alpha_6 = 0$  is isomorphic to  $T^1 \cdot (T^2 \times \text{U}(3))$  which we denote by  $U$ . It is clear that the homology of  $U$  is torsion free and  $E_6/U$  satisfies the condition (2.2).

**6.1. KO-theory of  $E_6/U$ .** Let us calculate the  $\mathbb{Z}_{(2)}$ -coefficient cohomology of  $F_4/U$ . We set some notation. Let  $\omega_i$  ( $i = 1, \dots, 6$ ) be the fundamental weight of  $E_6$  as in [TW]. Put

$$t_1 = -\omega_1 + \omega_2, \quad t_2 = \omega_1 + \omega_2 - \omega_3, \quad t_3 = \omega_2 + \omega_3 - \omega_4, \quad t_4 = \omega_4 - \omega_5, \quad t_5 = \omega_5 - \omega_6, \quad t_6 = \omega_6.$$

Then as in §2, we have

$$H^*(BT; \mathbb{Z}) = \mathbb{Z}_{(2)}[t, t_1, \dots, t_6]/(c_1 - 3t).$$

As in [TW], the Weyl group of  $U$  is generated by two elements  $R_1, R_2$  satisfying

$$\begin{aligned} R_1(t_i) &= t_i & (i = 1, 2, 3, 6), & & R_1(t_4) &= t_5, & & R_1(t_5) &= t_4, \\ R_2(t_i) &= t_i & (i = 1, 2, 3, 4), & & R_2(t_5) &= t_6, & & R_2(t_6) &= t_5. \end{aligned}$$

Then it follows that

$$H^*(BU; \mathbb{Z}_{(2)}) = \mathbb{Z}_{(2)}[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3],$$

where  $\hat{c}_1 = t_4 + t_5 + t_6$ ,  $\hat{c}_2 = t_4t_5 + t_5t_6 + t_6t_4$  and  $\hat{c}_3 = t_4t_5t_6$ .

As in [MT], the mod 2 cohomology of  $E_6$  is given as

$$H^*(E_6; \mathbb{Z}/2) = \mathbb{Z}/2[a_3]/(a_3^4) \otimes \Lambda(a_5, a_9, a_{15}, a_{17}, a_{23}), \quad |a_i| = i, \quad \beta a_5 = a_3^2.$$

Then by [T], we obtain the following.

**Proposition 6.1.** *There is a regular sequence  $\bar{\rho}_2, \bar{\rho}_5, \bar{\rho}_6, \bar{\rho}_8, \bar{\rho}_9, \bar{\rho}_{12}$  in  $\mathbb{Z}_{(2)}[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3]$  with  $|\bar{\rho}_i| = 2i$  satisfying*

$$H^*(E_6/U; \mathbb{Z}_{(2)}) = \mathbb{Z}_{(2)}[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3, \gamma_3]/(\bar{\rho}_2, \bar{\rho}_5, \bar{\rho}_6, \bar{\rho}_8, \bar{\rho}_9, \bar{\rho}_{12}, 2\gamma_3 + \bar{\rho}_3),$$

where  $\bar{\rho}_3$  is defined by the equation  $\text{Sq}^2 \bar{\rho}_2 = \bar{\rho}_3$ .

Let us compute the mod 2 cohomology of  $E_6/U$ . Let  $c_i$  be the  $i^{\text{th}}$  symmetric function in  $t_1, \dots, t_6$  for  $i = 1, \dots, 6$ . Obviously,  $c_i$  is a polynomial in  $t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3$ . A calculation in [TW] implies that the rational cohomology of  $E_6/U$  is given as

$$(6.1) \quad H^*(E_6/U; \mathbb{Q}) = \mathbb{Q}[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3]/(\sigma_2, \sigma_5, \sigma_6, \sigma_8, \sigma_9, \sigma_{12}),$$

where

$$\begin{aligned} \sigma_2 &= c_2 - \frac{4}{3^2}c_1^2, & \sigma_5 &= c_5 - \frac{1}{3}c_4c_1 + \frac{1}{3^2}c_3c_1^2 - \frac{2}{3^5}c_1^5, \\ \sigma_6 &= 8c_6 + c_3^2 - \frac{4}{3^2}c_4c_1^2 - \frac{4}{3^6}c_1^6, & \sigma_8 &= -3c_6c_1^2 + c_4^2 - c_4c_3c_1 + \frac{19}{3^4}c_4c_1^4 - \frac{5}{3^4}c_3c_1^5 + \frac{31}{3^8}c_1^8. \end{aligned}$$

By Proposition 6.1, we may put

$$\bar{\rho}_2 = c_2 - \frac{4}{3^2}c_1^2 \quad \text{and} \quad \bar{\rho}_3 = c_3 + c_2c_1.$$

Put

$$R_1 = \mathbb{Z}_{(2)}[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3, \gamma_3]/(\bar{\rho}_2, \bar{\rho}_5, 2\gamma_3 + \bar{\rho}_3)$$

Then since the natural map  $H^*(E_6/U; \mathbb{Z}_{(2)}) \rightarrow H^*(E_6/U; \mathbb{Q})$  is injective, there is a surjection  $R_1 \rightarrow H^*(E_6/U; \mathbb{Z}_{(2)})$  which reduces to a surjection

$$\phi_1 : R_1/2 \rightarrow H^*(E_6/U; \mathbb{Z}/2).$$

Put

$$(6.2) \quad \rho_2 = c_2, \quad \rho_3 = c_3 + c_2c_1, \quad \rho_5 = c_5 + c_4c_1.$$

Then  $\rho_2, \rho_3, \rho_5$  is a regular sequence in  $\mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3]$  and

$$R_1/2 = \mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3, \gamma_3]/(\rho_2, \rho_3, \rho_5),$$

implying that the Poincaré series of  $R_1/2$  is  $\frac{1-t^{10}}{(1-t^2)^4(1-t^6)}$ . On the other hand, the Poincaré series of  $H^*(E_6/U; \mathbb{Z}/2)$  and  $H^*(E_6/U; \mathbb{Q})$  are the same, which is  $\frac{(1-t^{10})(1-t^{12})(1-t^{16})(1-t^{18})(1-t^{24})}{(1-t^2)^4(1-t^6)}$  by (6.1). Then  $\phi_1$  is an isomorphism in dimension  $\leq 11$ .

Note that  $\sigma_6 \equiv 4(2c_6 + \gamma_3^2 + \frac{4}{3^2}\gamma_3c_1^3 - \frac{1}{3^2}c_4c_1^2 + \frac{35}{3^6}c_1^6) \pmod{(\bar{\rho}_2, 2\gamma_3 + \bar{\rho}_3)}$ . Then since  $H^*(E_6/U; \mathbb{Z}/2) \rightarrow H^*(E_6/U; \mathbb{Q})$  is injective, if we put

$$R_2 = R_1/(2c_6 + \gamma_3^2 + \frac{4}{3^2}\gamma_3c_1^3 - \frac{1}{3^2}c_4c_1^2 + \frac{35}{3^6}c_1^6, \sigma_8),$$

$\phi_1$  induces a surjection

$$\phi_2 : R_2/2 \rightarrow H^*(E_6/U; \mathbb{Z}/2).$$

Put

$$(6.3) \quad \rho_6 = \gamma_3^2 + c_4c_1^2 + c_1^6, \quad \rho_8 = c_6c_1^2 + c_4^2 + c_4c_1^4 + c_1^8.$$

Then one sees that

$$R_2/2 = \mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3, \gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_6, \rho_8).$$

Since  $\rho_2, \rho_3, \rho_5, \rho_6, \rho_8$  is a regular sequence in  $\mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3, \gamma_3]$ , one can calculate the Poincaré series of  $R_2/2$ . Then comparing the Poincaré series as above, we obtain that  $\phi_2$  is an isomorphism in dimension  $\leq 35$ .

Put

$$(6.4) \quad \rho_9 = c_6c_1^3, \quad \rho_{12} = c_6^2 + c_6c_4c_1^2 + c_4^2c_1^4 + c_4c_1^8.$$

Since  $\text{Sq}^2\phi_2(\rho_8) = \phi_2(\rho_9)$  and  $\text{Sq}^8\phi_2(\rho_8) = \phi_2(\rho_{12})$ , there is also a surjection

$$\phi_3 : R_3 \rightarrow H^*(E_6/U; \mathbb{Z}/2),$$

where

$$R_3 = R_2/(2, \rho_8, \rho_{12}).$$

Since  $\rho_2, \rho_3, \rho_5, \rho_6, \rho_8, \rho_9, \rho_{12}$  is a regular sequence in  $\mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3, \gamma_3]$ , one can calculate the Poincaré series of  $R_3$ . Comparing it with the Poincaré series of  $H^*(E_6/U; \mathbb{Z}/2)$ , we conclude that  $\phi_3$  is an isomorphism. Summarizing, we obtain the following.

**Proposition 6.2.** *The mod 2 cohomology of  $E_6/U$  is given as*

$$H^*(E_6/U; \mathbb{Z}/2) = \mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3, \gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_6, \rho_8, \rho_9, \rho_{12}),$$

where  $|t_i| = 2, |\hat{c}_i| = 2i, |\gamma_3| = 6$  and  $\rho_i$  is as in (6.2), (6.3) and (6.4).

**Corollary 6.3.** *The  $\text{Sq}^2$ -cohomology of  $E_6/U$  is given as*

$$H^*(E_6/U; \text{Sq}^2) = \Lambda(x_7, x_{11}, x_{15}), \quad |x_i| = 2i,$$

where  $\text{Sq}^2x_{11} \equiv \rho_{12} \pmod{(\rho_2, \rho_3, \rho_5, \rho_9)}$ ,  $\text{Sq}^2x_{15} = \rho_8^2$ ,  $x_7 = \gamma_3c_4 + \delta_7$  and  $\text{Sq}^2\delta_7 = c_4^2$  for  $\delta_7 \in \mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3]$ .



*Proof.* As in the proof of Corollary 5.3, we see that  $\text{Sq}^2\gamma_3 = c_4$ . Put  $A = \mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3]$ . Then our aim is to calculate the cohomology of a differential graded algebra

$$A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_6, \rho_8, \rho_9, \rho_{12}).$$

Obviously,  $A/(\rho_2, \rho_3) \cong \mathbb{Z}/2[t_1, t_2, t_3] \otimes \langle 1, \hat{c}_1, \hat{c}_1^2 \rangle$  as a  $\mathbb{Z}/2[t_1, t_2, t_3]$ -module, implying  $H^*(A/(\rho_2, \rho_3)) = 0$ . Then since  $dc_4 = \rho_5$  and  $d\rho_8 = \rho_9$ , it follows from Lemma 3.3 that

$$H^*(A/(\rho_2, \rho_3, \rho_5, \rho_8, \rho_9)) = \Lambda(c_4, x_{15}), \quad |x_i| = 2i,$$

where  $\text{Sq}^2x_{15} = \rho_8^2$ . For  $d\rho_{12} \equiv 0 \pmod{(\rho_5, \rho_9)}$  and  $H^{24}(A/(\rho_2, \rho_3, \rho_5, \rho_8, \rho_9)) = 0$ , we get

$$H^*(A/(\rho_2, \rho_3, \rho_5, \rho_8, \rho_9, \rho_{12})) = \Lambda(c_4, x_{11}, x_{15}), \quad |x_i| = 2i,$$

where  $\text{Sq}^2x_{11} \equiv \rho_{12} \pmod{(\rho_2, \rho_3, \rho_5, \rho_9)}$ . By the spectral sequence associated with a filtration

$$A/(\rho_2, \rho_3, \rho_5, \rho_8, \rho_{12}) \subset A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_8, \rho_9, \rho_{12}),$$

we get

$$H^*(A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_8, \rho_9, \rho_{12})) = \Lambda(x_7, x_{11}, x_{15}) \otimes \mathbb{Z}/2[\gamma_3^2],$$

where  $x_7 = \gamma_3c_4 + \delta_7$  and  $\delta_7 \in \mathbb{Z}/2[t_1, t_2, t_3, \hat{c}_1, \hat{c}_2, \hat{c}_3]$  is given by  $d\delta_7 = c_4^2$ . Since  $\rho_6 = \gamma_3^2 + d(\gamma_3c_1^2 + c_1^5)$ , we obtain

$$H^*(A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_6, \rho_8, \rho_9, \rho_{12})) = \Lambda(x_7, x_{11}, x_{15}),$$

completing the proof. □

**Theorem 6.4.** *The Atiyah-Hirzebruch spectral sequence  $E_r(E_6/U)$  collapses at  $E_3$ -term. In particular, we have*

$$g_{E_6/U}(t) = (1 + t^{14})(1 + t^{22})(1 + t^{30}).$$

*Proof.* From Lemma 2.1 and Proposition 6.3, the result follows. □

**Theorem 6.5.** *The KO-theory of  $E_6/U$  is given as*

$$KO^{2n-1}(E_6/U) \cong (\mathbb{Z}/2)^{s_n} \quad \text{and} \quad KO^{2n}(E_6/U) \cong (\mathbb{Z}/2)^{s_{n+1}} \oplus \mathbb{Z}^t$$

for  $n \in \mathbb{Z}/4$ , where

$$t = 4320, \quad s_0 = s_{-3} = 1, \quad s_{-1} = s_{-2} = 3.$$

*Proof.* By (6.1), we have  $f_{E_6/U}(t) = \frac{(1-t^{10})(1-t^{12})(1-t^{16})(1-t^{18})(1-t^{24})}{(1-t^2)^4(1-t^6)}$ . Then the proof is completed by Lemma 2.2 and Theorem 6.4. □

6.2. **KO-theory of  $E_6/T$ .** Let  $\rho_i \in \mathbb{Z}/2[t_1, \dots, t_6, \gamma_3]$  be as in (6.2), (6.3) and (6.4). The mod 2 cohomology of  $E_6/T$  is calculated in [KI2] as

$$H^*(E_6/T; \mathbb{Z}/2) = \mathbb{Z}/2[t_1, \dots, t_6, \gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_6, \rho_8, \rho_9, \rho_{12}),$$

where  $\text{Sq}^2 \gamma_3 = c_4$ . For the projection  $\pi : E_6/T \rightarrow E_6/U$ , we have

(6.5)

$$\pi^*(t_i) = t_i \quad (i = 1, 2, 3), \quad \pi^*(\hat{c}_1) = t_4 + t_5 + t_6, \quad \pi^*(\hat{c}_2) = t_4 t_5 + t_5 t_6 + t_6 t_4, \quad \pi^*(\hat{c}_3) = t_4 t_5 t_6.$$

Define a map  $\lambda : (E_6/T)_{(2)} \rightarrow BT_{(2)}^6$  by  $\lambda^*(t_i) = t_i$  for  $i = 1, \dots, 6$ . Then there is a lift  $\tilde{\lambda} : (E_6/T)_{(2)} \rightarrow B\tilde{T}_{(2)}^6$  satisfying

$$(6.6) \quad \tilde{\lambda}^*(t_i) = t_i \quad (i = 1, \dots, 6), \quad \tilde{\lambda}^*(\gamma_3) = \gamma_3,$$

where the second equality is shown in [KI1].

**Proposition 6.6.** *The  $\text{Sq}^2$ -cohomology of  $E_6/T$  is given as*

$$H^*(E_6/T; \text{Sq}^2) = \Lambda(x_3, x_7, x_{11}, x_{15}), \quad |x_i| = 2i,$$

where  $\tilde{\lambda}^*(x_3) = x_3$ ,  $\pi^*(x_7) = x_7$ ,  $\pi^*(x_{11}) = x_{11}$  and  $\pi^*(x_{15}) = x_{15}$ .

*Proof.* Define a differential graded algebra  $A$  as  $A = \mathbb{Z}/2[t_1, \dots, t_6]$  with  $|t_i| = 2$  and  $dt_i = t_i^2$ . Then we calculate the cohomology of a differential graded algebra  $A \otimes \mathbb{Z}/2[\gamma_3]/(\rho_2, \rho_3, \rho_5, \rho_6, \rho_8, \rho_9, \rho_{12})$ , where  $d\gamma_3 = c_4$ . This is done quite similarly to the proof of Proposition 6.3. The second assertion follows from (6.5) and (6.6).  $\square$

**Theorem 6.7.** *The spectral sequence  $E_r(E_6/T)$  collapses at  $E_3$ -term. In particular, we have*

$$g_{E_6/T}(t) = (1 + t^6)(1 + t^{14})(1 + t^{22})(1 + t^{30}).$$

*Proof.* By Theorem 3.7 and Proposition 6.6,  $\iota^{-1}(x_3)$  in the 2-localized spectral sequence  $E_3^{6,-1}(E_6/T)_{(2)}$  is a permanent cycle, implying that  $\iota^{-1}(x_3)$  in the integral spectral sequence  $E_3^{6,-1}(E_6/T)$  is also a permanent cycle since the 2-localization  $E_3^{p,q}(E_6/T) \rightarrow E_3^{p,q}(E_6/T)$  is injective. By Theorem 6.4 and Proposition 6.6,  $\iota^{-1}(x_i) \in E_3^{*,-1}(E_6/T)$  is also a permanent cycle for  $i = 7, 11, 15$ . Thus the result follows from Lemma 2.1.  $\square$

*Proof of Theorem 1.1 for  $E_6$ .* The result follows from (2.5), Lemma 2.2 and Corollary 6.7.  $\square$

*Remark 6.8.* We can not apply the same calculation method to  $E_7/T$  and  $E_8/T$  for which there is no control on elements  $\gamma_5, \gamma_9$  in their mod 2 cohomology [KI2].

## REFERENCES

- [A] M.F. Atiyah, *K-theory and reality*, Quart. J. Math. Oxford Ser. **17** (1966), 367-386.
- [F] M. Fujii, *KO-groups of projective spaces*, Osaka J. Math. **4** (1967), 141-149.
- [IT] K. Ishitoya and H. Toda, *On the cohomology of irreducible symmetric spaces of exceptional type*, J. Math. Kyoto Univ. **17** (1977), no. 2, 225-243.
- [K] D. Kishimoto, *KO-theory of complex Stiefel manifolds*, J. Math. Kyoto Univ. **44** (2004), no. 3, 669-674.

- [KKO] D. Kishimoto, A. Kono and A. Ohsita, *KO-theory of flag manifolds*, J. Math. Kyoto Univ. **44** (2004), no. 1, 217-227.
- [KH1] A. Kono and S. Hara, *KO-theory of complex Grassmannians*, J. Math. Kyoto Univ. **31** (1991), no. 3, 827-833.
- [KH2] A. Kono and S. Hara, *KO-theory of Hermitian symmetric spaces*, Hokkaido Math. J. **21** (1992), no. 1, 103-116.
- [KI1] A. Kono and K. Ishitoya, *Squaring operations in the 4-connective fibre spaces over the classifying spaces of the exceptional Lie groups*, Publ. Res. Inst. Math. Sci. **21** (1985), no. 6, 1299-1310.
- [KI2] A. Kono and K. Ishitoya, *Squaring operations in mod 2 cohomology of quotients of compact Lie groups by maximal tori*, Algebraic topology, Barcelona, 1986, 192-206, Lecture Notes in Math. **1298**, Springer, Berlin, 1987.
- [MT] M. Mimura and H. Toda, *Topology of Lie groups. I, II*, Translations of Mathematical Monographs **91**, American Mathematical Society, Providence, RI, 1991.
- [T] H. Toda, *On the cohomology ring of some homogeneous spaces*, J. Math. Kyoto Univ. **15** (1975), 185-199.
- [TW] H. Toda and T. Watanabe, *The integral cohomology ring of  $F_4/T$  and  $E_6/T$* , J. Math. Kyoto Univ. **14** (1974), 257-286.
- [Y1] N. Yagita, *A note on the Witt group and the KO-theory of complex Grassmannians*, J. K-theory **9** (2012), 161-175.
- [Y2] N. Yagita, *Witt groups of algebraic groups*, preprint.
- [Z] M. Zibrowius, *Witt groups of complex cellular varieties*, Doc. Math. **16** (2011), 465-511.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO, 606-8502, JAPAN  
*E-mail address:* kishi@math.kyoto-u.ac.jp

FACULTY OF ECONOMICS, OSAKA UNIVERSITY OF ECONOMICS, OSAKA 533-8533, JAPAN  
*E-mail address:* ohsita@osaka-ue.ac.jp