Homotopy-commutativity in rotation groups

Hiroaki Hamanaka
Department of Mathematics,
Kyoto University.

October 17, 1995

1 Introduction

Assume G is a topological group and S, S' are subspaces of G, each of which contains the unit as its base point. There is the commutator map c from $S \wedge S'$ to G which maps $(x, y) \in S \wedge S'$ to $xyx^{-1}y^{-1} \in G$. We say S and S' homotopy-commute in G if c is null homotopic.

In this paper, we describe the homotopy-commutativity of the case $G = SO(n+m-1)$, $S = SO(n)$ and $S' = SO(m)$ where $n, m > 1$. Here we use the usual embeddings

$$SO(1) \subset SO(2) \subset SO(3) \subset \cdots.$$

Trivially $SO(n)$ and $SO(m)$ homotopy-commute in $SO(n+m)$. And it is known that if $n + m > 4$, $SO(n)$ and $SO(m)$ do not homotopy-commute in $SO(n+m-2)$. (See [1] and [2].) But the homotopy-commutativity in $SO(n+m-1)$ has not been solved exactly.

We shall say a pair (n, m) is irregular if $SO(n)$ and $SO(m)$ homotopy-commute in $SO(n+m-1)$, and regular if they do not. In [1] the following problem is proposed; "when is (n, m) irregular?", and the next theorem is showed.

Theorem 1.1 *(James and Thomas) Let $n + m \neq 4, 8$. If n or m is even or if $d(n) = d(m)$ then (n, m) is regular, where $d(q)$, for $q \geq 2$, denotes the greatest power of 2 which devides $q - 1$.*

In this paper we shall prove the more strict result as showed in the next theorem.

Theorem 1.2 *If n or m is even or if $\binom{n+m-2}{n-1} \equiv 0$ mod 2 then (n, m) is regular.*

We identify $\mathbf{RP}^{k-1} \overset{i_k}{\hookrightarrow} SO(k)$ by the following way. Let $i'_k : \mathbf{RP}^{k-1} \to O(k)$ be the map which attatches a line $l \in \mathbf{RP}^{k-1}$ with $i'_k(l) \in O(k)$ defined by

$$i'_k(l)(v) = v - 2(v, e)e,$$

where e is a unit vector of l and $v \in \mathbf{R}^k$. And let $i_k(l) = i'_k(l_0)^{-1} \cdot i'_k(l)$ where l_0 is the base point of $\mathbf{RP}^{k-1}$. Then i_k preserves the base points.

Theorem 1.2 follows from the next theorem.

Theorem 1.3 *Let n and m be odd. $\mathbf{RP}^{n-1} \subset SO(n)$ and $\mathbf{RP}^{m-1} \subset SO(m)$ homotopy-commute in $SO(n+m-1)$ if and only if*

$$\binom{n+m-2}{n-1} \equiv 1 \text{ mod } 2.$$

Let $\mathbf{SO}$ be $\lim_{\to}(SO(1) \subset SO(2) \subset SO(3) \subset \cdots)$ and consider the fibration $SO(n+m-1) \to \mathbf{SO} \to \mathbf{SO}/SO(n+m-1)$. Then we have a sequence of spaces

$$\cdots \to \Omega\mathbf{SO} \overset{\Omega p}{\to} \Omega(\mathbf{SO}/SO(n+m-1)) \overset{\delta}{\to} SO(n+m-1) \overset{i}{\to} \mathbf{SO} \overset{p}{\to} \mathbf{SO}/SO(n+m-1).$$

We can see $i \circ c|_{\mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1}} \simeq * : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \mathbf{SO}$. This means there exists $\lambda : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega\mathbf{SO}/SO(n+m-1)$ such that $\delta \circ \lambda = c|_{\mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1}}$. The construction of λ and the cohomology map λ^* are argued in §2. We describe about lifts of λ in §3 and finally, in §4, we determine when a lift of λ exists, which means when $c|_{\mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1}} \simeq *$.

2 Lift λ of c

Definition A sequence of spaces X_i and continuous maps f_i

$$\cdots \to X_{i+1} \overset{f_i}{\to} X_i \to \cdots \overset{f_0}{\to} X_0$$

is called a fibration sequence if, for any $i \geq 0$, there exists a fibration $Y_i^{(2)} \xrightarrow{j_i} Y_i^{(1)} \xrightarrow{\pi_i} Y_i^{(0)}$, homotopy equivalence maps $\psi_i^{(k)} : X_{i+k} \to Y_i^{(k)}$ (k =0,1,2), and the following diagram commutes upto homotopy.

$$\begin{array}{ccccc} X_{i+2} & \xrightarrow{f_{i+1}} & X_{i+1} & \xrightarrow{f_i} & X_i \\ \simeq\downarrow \psi_i^{(2)} & & \simeq\downarrow \psi_i^{(1)} & & \simeq\downarrow \psi_i^{(0)} \\ Y_i^{(2)} & \xrightarrow{j_i} & Y_i^{(1)} & \xrightarrow{\pi_i} & Y_i^{(0)} \end{array}$$

For example, given a fibration $F \to E \to B$, there is a fibration sequence

$$\cdots \to \Omega F \to \Omega E \to \Omega B \to F \to E \to B.$$

Consider the fibration $\mathbf{SO} \to \mathbf{SO}/SO(n+m-1)$ with the fibre $SO(n+m-1)$. Then we have a fibration sequence.

$$\cdots \to \Omega\mathbf{SO} \xrightarrow{\Omega p} \Omega(\mathbf{SO}/SO(n+m-1)) \xrightarrow{\delta} SO(n+m-1) \xrightarrow{i} \mathbf{SO} \xrightarrow{p} \mathbf{SO}/SO(n+m-1)$$

Obviouly $i \circ c : SO(n) \wedge SO(m) \to \mathbf{SO}$ is null homotopic. This means there exists a lift of c, that is, a map $\lambda : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega(\mathbf{SO}/SO(n+m-1))$ such that $\delta \circ \lambda \simeq c$.

In R.Bott[3] it is showed that the following map $\lambda_0 : SO(n) \wedge SO(m) \to \Omega(\mathbf{SO}/SO(n+m-1))$ is a lift of c.

Recall the fibration $SO(k-1) \to SO(k) \xrightarrow{p_k} S^{k-1}$. Define h as $h = \Sigma(p_n \wedge p_m) : \Sigma(SO(n) \wedge SO(m)) \to \Sigma(S^{n-1} \wedge S^{m-1}) \simeq S^{n+m-1}$. Then ad$h$ is a lift of c in the following fibration sequence. (See [5].)

$$\begin{array}{ccccccc} \cdots \to \Omega SO(n+m) \to & \Omega S^{n+m-1} & \to & SO(n+m-1) & \to SO(n+m) \to S^{n+m-1} \\ & & \nwarrow \mathrm{ad}h & \uparrow c & & \\ & & & SO(n) \wedge SO(m) & & \end{array}$$

The fibration $SO(n+m) \to S^{n+m-1}$ is the restriction of $\mathbf{SO} \to \mathbf{SO}/SO(n+m-1)$ to $S^{n+m-1} = SO(n+m)/SO(n+m-1) \xrightarrow{j} \mathbf{SO}/SO(n+m-1)$. Therefore we define λ_0 as $\Omega j \circ \mathrm{ad}h$. Refer to the commutative diagram below.

The rest of this section is devoted to the computation of the cohomology map of λ. And throughout this paper we use $\mathbf{Z}/2\mathbf{Z}$ as the coefficient ring of cohomology unless mentioned.


$\Omega\mathbf{SO}$
Ωp
Ωj
ΩS^{n+m-1}
$\Omega(\mathbf{SO}/SO(n+m-1))$
$\mathrm{ad}h$
δ
$SO(n)\wedge SO(m)$
$SO(n+m-1)$
$SO(n+m-1)$
i
$\mathbf{RP}^{n-1}\wedge\mathbf{RP}^{m-1}$
$SO(n+m)$
$\mathbf{SO}$
p
j
S^{n+m-1}
$\mathbf{SO}/SO(n+m-1)$

First we refer to the cohomology rings of spaces which are used in this paper, that is,

$$\begin{aligned}
\mathrm{H}^*(\Omega_0\mathbf{SO}) &= \mathbf{Z}/2\mathbf{Z}[\alpha_2, \alpha_4, \alpha_6, \cdots]/(\alpha_{4k} - \alpha_{2k}^2), \\
\mathrm{H}^*(\Omega(\mathbf{SO}/SO(n+m-1))) &= \mathbf{Z}/2\mathbf{Z}[\alpha'_{n+m-2}, \alpha'_{n+m}, \cdots]/(\alpha'_{4k} - \alpha'^2_{2k}), \\
\mathrm{H}^*(SO(k)) &= \Delta(x_1, \cdots, x_{k-1}), \\
\mathrm{H}^*(SO(k)/SO(k-l)) &= \Delta(x'_{k-l}, \cdots, x'_{k-1}),
\end{aligned}$$

where $\deg(\alpha_{2i}) = 2i$, $\deg(\alpha'_{2i}) = 2i$, $\deg(x_i) = i$. And also

$$\Omega p^*(\alpha'_k) = \alpha_k.$$

Lemma 2.4 $\lambda_0^*(\alpha'_{n+m-2}) = x_{n-1} \otimes x_{m-1}$.

Proof. Consider the fibration $p_k : SO(k) \to S^{k-1}$ with the fibre $SO(k-1)$. Let c_i be the generator of $\mathrm{H}^i(S^i)$. Then $p_k^*(c_{k-1}) = x_{k-1}$. Thus we have

$$\begin{aligned}
h^*(c_{n+m-1}) &= \Sigma(p_n \wedge p_m)^*(\Sigma c_{n-1} \otimes c_{m-1}) \\
&= \Sigma(x_{n-1} \otimes x_{m-1}).
\end{aligned}$$

Hence $(\mathrm{ad}h)^*(\sigma c_{n+m-1}) = x_{n-1} \otimes x_{m-1}$, where σ is the cohomology suspension $\sigma : \mathrm{H}^{*+1}(X) \to \mathrm{H}^*(\Omega X)$.

On the other hand, $j^*(x_{n+m-1}) = c_{n+m-1}$ means

$$\begin{aligned}
(\Omega j)^*(\alpha'_{n+m-2}) &= (\Omega j)^*(\sigma x'_{n+m-1}) \\
&= \sigma c_{n+m-1}.
\end{aligned}$$

Therefore it follows that

$$\begin{aligned}
\lambda_0^*(\alpha'_{n+m-2}) &= (\mathrm{ad}h)^*(\Omega j)^*(\alpha'_{n+m-2}) \\
&= x_{n-1} \otimes x_{m-1}.
\end{aligned}$$

Q.E.D.

Now let $\lambda = \lambda_0 \circ (i_m \wedge i_n) : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega(\mathbf{SO}/SO(n+m-1))$ and in the following we use c as the commutator map from $\mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1}$ to $SO(n+m-1)$. Easily we have $i_k^*(x_{k-1}) = \tau^{k-1}$ where τmeans the generator of $\mathrm{H}^1(\mathbf{RP}^{k-1})$. (See Whitehead [4].) Thus

$$\begin{aligned}
\lambda^*(\alpha'_{n+m-2}) &= (i_m \wedge i_n)^* \circ \lambda_0^*(\alpha'_{n+m-2}) \\
&= \tau^{n-1} \otimes \tau^{m-1}.
\end{aligned}$$

3 Lift of λ and homotopy commutativity

In this section we prove the next theorem.

Theorem 3.5 *Let n, m be odd.*

1. *$c \simeq *$ if and only if there exists a lift of λ, that is, a map $x : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega_0(\mathbf{SO})$ such that $\lambda = \Omega p \circ x$.*
2. *$c \simeq *$ if and only if there exists $x : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega_0(\mathbf{SO})$ such that $x^*(\alpha_{n+m-2}) \simeq \tau^{n-1} \otimes \tau^{m-1}$.*

Proof. 1. The sequence

$$\cdots \to \Omega_0(\mathbf{SO}) \stackrel{\Omega p}{\to} \Omega(\mathbf{SO}/SO(n+m-1)) \stackrel{\delta}{\to} SO(n+m-1)$$

is a fibration sequence and λ is a lift of c. Therefore the statement follows.

2. By the first statement it is sufficient to prove that x is a lift of λ if and only if $x^*(\alpha_{n+m-2}) = \tau^{n-1} \otimes \tau^{m-1}$. We need the following lemma.

Lemma 3.6 *Let n and m be odd. Then*

$$\pi_i(\mathbf{SO}/SO(n+m-1)) = \begin{cases} 0 & i \leq n+m-2 \\ \mathbf{Z}/2\mathbf{Z} & i = n+m-1 \end{cases}$$

Proof. Consider the fibration

$$SO(n+m+1)/SO(n+m-1) \to \mathbf{SO}/SO(n+m-1) \to \mathbf{SO}/SO(n+m+1)$$

and see the homotopy exact sequence. Remark that $\pi_i(\mathbf{SO}/SO(2k+1)) = 0$ for $i \leq 2k$ and we obtain

$$\pi_{n+m-1}(\mathbf{SO}/SO(n+m-1)) = \pi_{n+m-1}(SO(n+m+1)/SO(n+m-1)).$$

It is known that $\pi_{n+m-1}(SO(n+m+1)/SO(n+m-1)) = \mathbf{Z}/2\mathbf{Z}$ provided $n+m-1$ is odd. Hence we obtained the statement.

Q.E.D.

By Lemma 3.6 it follows that

$$\pi_i(\Omega(\mathbf{SO}/SO(n+m-1))) = \begin{cases} 0 & i \leq n+m-3 \\ \mathbf{Z}/2\mathbf{Z} & i = n+m-2. \end{cases}$$

Now add cells $e_i (i \geq 1)$ to $\Omega(\mathbf{SO}/SO(n+m-1))$ so that $\pi_k(\Omega(\mathbf{SO}/SO(n+m-1))$ vanishes for $k \geq n+m-1$, where $\dim e_i \geq n+m$. We shall call the obtained space K, that is,

$$\Omega(\mathbf{SO}/SO(n+m-1)) \hookrightarrow \Omega(\mathbf{SO}/SO(n+m-1)) \cup e_1 \cup e_2 \cup \cdots = K \quad (1)$$

and

$$\pi_i(K) = \begin{cases} \mathbf{Z}/2\mathbf{Z} & i = n+m-2 \\ 0 & \text{otherwise.} \end{cases} \quad (2)$$

Thus K is an Eilenberg-Maclane space $K(\mathbf{Z}/2\mathbf{Z}; n+m-2)$. Let γ denote the inclusion map from $\Omega(\mathbf{SO}/SO(n+m-1))$ to K. Here

$$\gamma_* : \pi_{n+m-2}(\Omega(\mathbf{SO}/SO(n+m-1))) \to \pi_{n+m-2}(K)$$

is not a 0-map. This means that by the isomorphism

$$[\Omega(\mathbf{SO}/SO(n+m-1)), K] \cong \mathrm{H}^{n+m-2}(\Omega(\mathbf{SO}/SO(n+m-1)))$$

γ corresponds to α'_{n+m-2}, that is, $\gamma^* u = \alpha'_{n+m-2}$ where u is the generator of $\mathrm{H}^{n+m-2}(K)$.

On the other hand, (1) and (2) imply that $\gamma_* : \pi_i(\Omega(\mathbf{SO}/SO(n+m-2))) \to \pi_i(K)$ is isomorphic for $i \leq n+m-2$ and epic for $i \geq n+m-1$. Then by Whitehead's theorem

$$\begin{aligned} [\mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1}, \Omega(\mathbf{SO}/SO(n+m-1))] &\cong [\mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1}, K] \\ &\cong \mathrm{H}^{n+m-2}(\mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1}). \end{aligned}$$

Thus maps f and $g : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega(\mathbf{SO}/SO(n+m-2))$ are homotopic if and only if $f^*(\alpha'_{n+m-2}) = g^*(\alpha'_{n+m-2})$.

Now we assume $x : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega(\mathbf{SO}/SO(n+m-1))$ satisfies that $x^*(\alpha_{n+m-2}) = \tau^{n-1} \otimes \tau^{m-1}$. Then

$$(\Omega p \circ x)^*(\alpha'_{n+m-2}) = \tau^{n-1} \otimes \tau^{m-1}.$$

By §2 $\lambda^*(\alpha'_{n+m-2}) = \tau^{n-1} \otimes \tau^{m-1}$. Thus we obtain $\Omega p \circ x \simeq \lambda$ and x is a lift of λ.

The inverse is trivial and the proof of theorem 3.5 is finished.

4 Existence of lift of λ

In this section we prove the next theorem which completes the proof of Theorem 1.3.

Theorem 4.7 *Let n and m be odd. There exists a map $x : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega_0(\mathbf{SO})$ such that $x^*(\alpha_{n+m-2}) = \tau^{n-1} \otimes \tau^{m-1}$ if and only if*

$$\binom{n+m-2}{n-1} \equiv 1 \ mod\ 2.$$

Proof. First consider

$$\theta := (r_1 - 1)\hat{\otimes}(r_1 - 1)\hat{\otimes}(r_\infty - 1)\hat{\otimes}(r_\infty - 1) \in \widetilde{KO}(\Sigma^2(\mathbf{RP}^\infty \wedge \mathbf{RP}^\infty)).$$

Here r_1 is the Möbius line bundle over S^1 and r_∞ is the canonical line bundle over $\mathbf{RP}^\infty$. Now we compute the total Stiefel Whitney class of θ. We start from the next lemma.

Lemma 4.8 *Let $A = 1 + a_1 + a_2 + \cdots \in \mathrm{H}^{**}(\mathbf{RP}^\infty \times \mathbf{RP}^\infty)$ where $a_i \in \mathrm{H}^i(\mathbf{RP}^\infty \times \mathbf{RP}^\infty)$ and let $s_i \in \mathrm{H}^*(S^1 \times S^1 \times \mathbf{RP}^\infty \times \mathbf{RP}^\infty)$ $(i = 1, 2)$ be the pull back of the generator of $\mathrm{H}^1(S^1)$ by the canonical projection from $S^1 \times S^1 \times \mathbf{RP}^\infty \times \mathbf{RP}^\infty$ to the ith factor S^1. Then we have*

$$\frac{(A + s_1 + s_2)A}{(A + s_1)(A + s_2)} = \frac{A^2 + s_1 s_2}{A^2} \in \mathrm{H}^{**}(S^1 \times S^1 \times \mathbf{RP}^\infty \times \mathbf{RP}^\infty).$$

Proof. By direct computation, we see

$$\begin{aligned}
\frac{(A + s_1 + s_2)A}{(A + s_1)(A + s_2)} &= \frac{\{(A + s_1)^2 + (A + s_1)s_2\}A}{(A + s_1)^2(A + s_2)} \\
&= \frac{(A^2 + s_2 A + s_1 s_2)A}{A^2(A + s_2)} \\
&= \frac{A(A + s_2)^2 + (A + s_2)s_1 s_2}{A(A + s_2)^2} \\
&= \frac{A^2 + s_1 s_2}{A^2}.
\end{aligned}$$

Q.E.D.

Let $\pi : S^1 \times S^1 \times \mathbf{RP}^\infty \times \mathbf{RP}^\infty \to \Sigma^2(\mathbf{RP}^\infty \wedge \mathbf{RP}^\infty)$ be the canonical projection and decompose $\pi^*\theta$ as

$$\begin{aligned}\pi^*\theta &= r_1 \times r_1 \times r_\infty \times r_\infty + 1 \times 1 \times r_\infty \times r_\infty - 1 \times r_1 \times r_\infty \times r_\infty - r_1 \times 1 \times r_\infty \times r_\infty \\ &\quad - r_1 \times r_1 \times 1 \times r_\infty - 1 \times 1 \times 1 \times r_\infty + 1 \times r_1 \times 1 \times r_\infty + r_1 \times 1 \times 1 \times r_\infty \\ &\quad - r_1 \times r_1 \times r_\infty \times 1 - 1 \times 1 \times r_\infty \times 1 + 1 \times r_1 \times r_\infty \times 1 + r_1 \times 1 \times r_\infty \times 1 \\ &\quad + r_1 \times r_1 \times 1 \times 1 + 1 \times 1 \times 1 \times 1 - 1 \times r_1 \times 1 \times 1 - r_1 \times 1 \times 1 \times 1.\end{aligned}$$

Then the total Stiefel Whitney class $w(\pi^*\theta)$ of $\pi^*\theta$ is given by

$$\begin{aligned}w(\pi^*\theta) &= \frac{(1+\tau_1+\tau_2+s_1+s_2)(1+\tau_1+\tau_2)}{(1+\tau_1+\tau_2+s_1)(1+\tau_1+\tau_2+s_2)} \cdot \frac{(1+s_1+s_2)}{(1+s_1)(1+s_2)} \\ &\quad \cdot \Big\{ \frac{(1+\tau_1+s_1+s_2)(1+\tau_1)}{(1+\tau_1+s_1)(1+\tau_1+s_2)} \cdot \frac{(1+\tau_2+s_1+s_2)(1+\tau_2)}{(1+\tau_2+s_1)(1+\tau_2+s_2)} \Big\}^{-1}.\end{aligned}$$

Here τ_i $(i = 1, 2)$ is the pull back of the generator of the cohomology ring of the ith factor of $\mathbf{RP}^\infty \times \mathbf{RP}^\infty$. By the previous lemma, we obtain

$$\begin{aligned}w(\pi^*\theta) &= \frac{1+\tau_1^2+\tau_2^2+s_1s_2}{1+\tau_1^2+\tau_2^2} \cdot (1+s_1s_2) \cdot (\frac{1+\tau_1^2+s_1s_2}{1+\tau_1^2})^{-1} \cdot (\frac{1+\tau_2^2+s_1s_2}{1+\tau_2^2})^{-1} \\ &= \{1+(1+\tau_1^2+\tau_2^2)^{-1}s_1s_2\}(1+s_1s_2)\{1+(1+\tau_1^2)^{-1}s_1s_2\}\{1+(1+\tau_2^2)^{-1}s_1s_2\} \\ &= 1+s_1s_2\{(1+\tau_1^2+\tau_2^2)^{-1}+1+(1+\tau_1^2)^{-1}+(1+\tau_2^2)^{-1}\} \\ &= 1+s_1s_2\Big\{\sum_{i=0}^{\infty}(\tau_1^2+\tau_2^2)^i+1+\sum_{i=0}^{\infty}\tau_1^{2i}+\sum_{i=0}^{\infty}\tau_2^{2i}\Big\} \\ &= 1+s_1s_2\Big\{\sum_{i=2}^{\infty}\sum_{j=1}^{i-1}\binom{i}{j}\tau_1^{2j}\tau_2^{2i-2j}\Big\}.\end{aligned}$$

Therefore we see

$$w(\theta) = 1+\Sigma^2\Big\{\sum_{i=2}^{\infty}\sum_{j=1}^{i-1}\binom{i}{j}\tau^{2j}\otimes\tau^{2i-2j}\Big\}.$$

Let f be the classifing map of θ, that is, the map

$$f : \Sigma^2(\mathbf{RP}^\infty \wedge \mathbf{RP}^\infty) \to \mathbf{BSO}$$

such that $f^*(\xi) = \theta$ where $\xi = \lim_{n\to\infty}(\xi_n - n)$ and ξ_n is the universal $SO(n)$ vector bundle over $\mathbf{BSO}(n)$.

It is known that $\mathrm{H}^*(\mathbf{BSO}) = \mathbf{Z}/2\mathbf{Z}[w_1, w_2, \cdots]$ where w_i is the ith Stiefel Whitney class. Let $\iota_k : \mathbf{RP}^k \to \mathbf{RP}^\infty$ be the inclusion map and let

$$x_0 := (\mathrm{ad}^2 f) \circ (\iota_{n-1} \wedge \iota_{m-1}) : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega\mathbf{SO}.$$

Then it follows that for $N \geq 1$

$$\begin{aligned} x_0^*(\alpha_{2N}) &= (\iota_{n-1} \wedge \iota_{m-1})^* (\mathrm{ad}^2 f)^* \sigma^2 w_{2N+2} \\ &= (\iota_{n-1} \wedge \iota_{m-1})^* \Big(\sum_{j=1}^{N-1} \tbinom{2N}{2j} \tau^{2j} \otimes \tau^{2N-2j} \Big). \end{aligned}$$

Particularly $x_0^*(\alpha_{n+m-2}) = \binom{n+m-2}{n-1} \tau^{n-1} \otimes \tau^{m-1}$. Thus if $\binom{n+m-2}{n-1} \equiv 1$ then there exists $x_0 : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega\mathbf{SO}$ such that $x_0^*(\alpha_{n+m-2}) = \tau^{n-1} \otimes \tau^{m-1}$.

Now we shall prove the inverse, that is, prove that if $\binom{n+m-2}{n-1} \equiv 0 \bmod 2$ then $x^*(\alpha_{n+m-2}) = 0$ for any $x : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega\mathbf{SO}$. Let $n = 2a+1$, $m = 2b+1$ where $a, b \in \mathbf{Z}$, $a, b \geq 1$. Moreover we set $a \leq b$.

Here we use the Steenrod's square operators Sq^i. In $\mathrm{H}^*(\Omega_0\mathbf{SO})$, Sq^i acts as follows

$$\mathrm{Sq}^i(\alpha_{2j}) = \begin{cases} \binom{2j+1}{i} \alpha_{2j+i} & i \text{ is even} \\ 0 & i \text{ is odd.} \end{cases}$$

Let $x : \mathbf{RP}^{2a} \wedge \mathbf{RP}^{2b} \to \Omega_0\mathbf{SO}$ be an arbitrary map.

Lemma 4.9 *We set a, b, x as above then*

$$x^*(\alpha_2) = 0 \text{ and } x^*(\alpha_6) = \tau^2 \otimes \tau^4 + \tau^4 \otimes \tau^2 \text{ or } 0.$$

Proof. Since $x^*(\alpha_2) \in \mathrm{H}^*(\mathbf{RP}^{2a} \wedge \mathbf{RP}^{2b})$, $x^*(\alpha_2) = \tau \otimes \tau$ or 0. If $x^*(\alpha_2) = \tau \otimes \tau$, then we have

$$\mathrm{Sq}^1 x^*(\alpha_2) = \tau^2 \otimes \tau + \tau \otimes \tau^2.$$

On the other hand,

$$\mathrm{Sq}^1 x^*(\alpha_2) = x^*(\mathrm{Sq}^1 \alpha_2) = 0.$$

Therefore $x^*(\alpha_2) = 0$.

Next we consider $x^*(\alpha_6)$. If $(a,b) = (1,1)$ then $x^*(\alpha_6) = 0$, and if $(a,b) = (1,2)$ we can see $x^*(\alpha_6) = \tau^2 \otimes \tau^4$ or 0 as asserted. And otherwise, set

$$x^*(\alpha_6) = \rho_1\tau \otimes \tau^5 + \rho_2\tau^2 \otimes \tau^4 + \rho_3\tau^3 \otimes \tau^3 + \rho_4\tau^4 \otimes \tau^2 + \rho_5\tau^5 \otimes \tau^1,$$

where $\rho_i \in \mathbf{Z}/2\mathbf{Z}$ and the statement follows the next two equations.

$$\mathrm{Sq}^1 x^*(\alpha_6) = x^*(\mathrm{Sq}^1\alpha_6) = 0$$

$$\mathrm{Sq}^2 x^*(\alpha_6) = x^*(\alpha_8) = x^*(\alpha_2)^4 = 0$$

Q.E.D.

Remark that if $2(a+b) = 2^d - 2$ for some $d \in \mathbf{N}$, then $\binom{2(a+b)}{2i} \equiv 1 \bmod 2$ for any $i \in \mathbf{Z}$ such that $0 \leq i \leq a+b$. And also when $2(a+b) = 2^d$ for some $d \in \mathbf{N}$,

$$\binom{2(a+b)}{2i} \equiv \begin{cases} 1 \bmod 2 & i = 0 \text{ or } a+b \\ 0 \bmod 2 & \text{otherwise.} \end{cases}$$

In this case

$$\begin{aligned} x^*(\alpha_{2(a+b)}) &= x^*(\alpha_{2^d}) \\ &= x^*(\text{a power of } \alpha_2) \\ &= 0 \end{aligned}$$

as asserted. Hence we can assume that $2(a+b) \neq 2^k$ or $2^k - 2$ for any $k \in \mathbf{N}$.

Next we shall prove the next theorem.

Theorem 4.10 *Let a, b and x be as above. If $x^*(\alpha_6) = 0$ then $x^*(\alpha_{2(a+b)}) = 0$.*

Proof. Let d be the number which satisfies

$$2^d < 2(a+b) < 2^{d+1} - 2 \quad d \in \mathbf{N}.\ (d \geq 3)$$

We distinguish between the following two cases.

I)
$$2^d < 2(a+b) < 3 \cdot 2^{d-1} - 2 \tag{3}$$

II)
$$3 \cdot 2^{d-1} - 2 \leq 2(a+b) < 2^{d+1} - 2 \tag{4}$$

Lemma 4.11 *Let a, b and x be as above. In any of the case* I) *and* II), *if $x^*(\alpha_6) = 0$ then one of the following holds.*

i) $x^*(\alpha_{2^k-2}) = 0$ *for* $3 \leq k \leq d-1$.

ii) $2a = 2^r - 2$ *for some* $r \in \mathbf{N}$, $r \leq d-1$ *and*

$$x^*(\alpha_{2^k-2}) = \begin{cases} 0 & 3 \leq k \leq r \\ \tau^{2^r-2} \otimes \tau^{2^k-2^r} & r+1 \leq k \leq d-1. \end{cases}$$

Proof. We use induction, that is, we prove the next two propositions.

a) If $x^*(\alpha_{2^{k-1}-2}) = 0$ and $4 \leq k \leq d-1$, then one of the followings holds.

- $x^*(\alpha_{2^k-2}) = 0$.
- $2a = 2^{k-1} - 2$ and $x^*(\alpha_{2^k-2}) = \tau^{2^{k-1}-2} \otimes \tau^{2^{k-1}}$.

b) If $2a = 2^r - 2$ and $x^*(\alpha_{2^{k-1}-2}) = \tau^{2^r-2} \otimes \tau^{2^{k-1}-2^r}$ and $r+2 \leq k \leq d-1$, then

$$x^*(\alpha_{2^k-2}) = \tau^{2^r-2} \otimes \tau^{2^k-2^r}.$$

First we assume $4 \leq k \leq d-1$ and $x^*(\alpha_{2^{k-1}-2}) = 0$ and prove a). Let

$$x^*(\alpha_{2^k-2}) = \sum_{i=s}^{t} \rho_i \tau^i \otimes \tau^{(2^k-2)-i},$$

where

$$\begin{aligned} s &= \max\{1, (2^k-2) - 2b\}, \\ t &= \min\{2^k - 3, 2a\}, \\ \rho_i &\in \mathbf{Z}/2\mathbf{Z}. \end{aligned}$$

Since $\mathrm{Sq}^1(x^*(\alpha_{2^k-2})) = x^*(\mathrm{Sq}^1\alpha_{2^k-2}) = 0$, we have that

$$\begin{aligned} & \mathrm{Sq}^1\Big(\sum_{i=s}^{t} \rho_i \tau^i \otimes \tau^{(2^k-2)-i}\Big) \\ = & \sum_{s \leq i \leq t,\ i:\ \text{odd}} \rho_i(\tau^{i+1} \otimes \tau^{(2^k-2)-i} + \tau^i \otimes \tau^{(2^k-2)-i+1}) \\ = & \sum_{s \leq i \leq t,\ i:\ \text{odd}} \rho_i(\tau^i \otimes \tau^{(2^k-2)-i+1}) + \sum_{s+1 \leq i \leq t+1,\ i:\ \text{even}} \rho_{i-1}(\tau^i \otimes \tau^{(2^k-2)-i+1}) \\ = & \ 0. \end{aligned}$$

Here, $\tau^i \otimes \tau^{(2^k-2)-i+1} \neq 0$ for $s+1 \leq i \leq t$. Therefore

$$\rho_i = 0 \text{ for } i: \text{ odd}, \ s \leq i \leq t. \tag{5}$$

Next we use Sq^2. By (5) we can set

$$x^*(\alpha_{2^k-2}) = \sum_{i=s'}^{t'} \rho_{2i}\tau^{2i} \otimes \tau^{(2^k-2)-2i},$$

$$\begin{aligned} \text{where } s' &= \max\{1, \frac{2^k-2}{2} - b\} \\ t' &= \min\{\frac{2^k-4}{2}, a\}. \end{aligned}$$

Since

$$\begin{aligned} \mathrm{Sq}^2 x^*(\alpha_{2^k-2}) &= x^*(\mathrm{Sq}^2\alpha_{2^k-2}) \\ &= x^*(\alpha_{2^k}) \\ &= x^*(\alpha_2^{2^{k-1}}) \\ &= 0, \end{aligned}$$

we have

$$\begin{aligned} & \mathrm{Sq}^2\Big(\sum_{i=s'}^{t'} \rho_{2i}\tau^{2i} \otimes \tau^{(2^k-2)-2i}\Big) \\ = & \sum_{s' \leq 2j \leq t'} \rho_{4j}\mathrm{Sq}^2(\tau^{4j} \otimes \tau^{(2^k-2)-4j}) + \sum_{s' \leq 2j-1 \leq t'} \rho_{4j-2}\mathrm{Sq}^2(\tau^{4j-2} \otimes \tau^{(2^k-2)-4j+2}) \\ = & \sum_{s' \leq 2j \leq t'} \rho_{4j}\tau^{4j} \otimes \tau^{2^k-4j} + \sum_{s' \leq 2j-1 \leq t'} \rho_{4j-2}\tau^{4j} \otimes \tau^{2^k-4j} = 0 \end{aligned} \tag{6}$$

Here $\tau^{4j} \otimes \tau^{2^k-4j} \neq 0$ for $s'+1 \leq 2j \leq t'$. Thus

$$\rho_{4j} = \rho_{4j-2} \text{ for } s'+1 \leq 2j \leq t'. \tag{7}$$

Next we consider Sq^4. Since

$$\begin{aligned} \mathrm{Sq}^4 x^*(\alpha_{2^k-2}) &= x^*(\mathrm{Sq}^4\alpha_{2^k-2}) \\ &= x^*(\alpha_{2^k+2}) \\ &= x^*(\mathrm{Sq}^{2^{k-1}}\mathrm{Sq}^4\alpha_{2^{k-1}-2}) \\ &= \mathrm{Sq}^{2^{k-1}}\mathrm{Sq}^4 x^*(\alpha_{2^{k-1}-2}) \\ &= 0, \end{aligned}$$

we have that

$$\begin{aligned}
& \mathrm{Sq}^4\Big(\sum_{i=s'}^{t'} \rho_{2i}\tau^{2i} \otimes \tau^{(2^k-2)-2i}\Big) \\
= \ & \mathrm{Sq}^4\Big(\sum_{s'\le 4j\le t'} \rho_{8j}\tau^{8j} \otimes \tau^{(2^k-2)-8j} + \sum_{s'\le 4j-1\le t'} \rho_{8j-2}\tau^{8j-2} \otimes \tau^{(2^k-2)-8j+2} \\
& + \sum_{s'\le 4j-2\le t'} \rho_{8j-4}\tau^{8j-4} \otimes \tau^{(2^k-2)-8j+4} + \sum_{s'\le 4j+1\le t'} \rho_{8j+2}\tau^{8j+2} \otimes \tau^{(2^k-2)-8j-2}\Big) \\
= \ & \sum_{s'\le 4j\le t'} \rho_{8j}\tau^{8j} \otimes \tau^{2^k+2-8j} + \sum_{s'\le 4j-1\le t'} \rho_{8j-2}\tau^{8j+2} \otimes \tau^{2^k-8j} \\
& + \sum_{s'\le 4j-2\le t'} \rho_{8j-4}\tau^{8j} \otimes \tau^{2^k+2-8j} + \sum_{s'\le 4j+1\le t'} \rho_{8j+2}\tau^{8j+2} \otimes \tau^{2^k-8j} \\
= \ & 0.
\end{aligned} \tag{8}$$

Thus

$$\begin{cases} \rho_{8j} = \rho_{8j-4} \text{ for } s'+2 \le 4j \le t' \\ \rho_{8j-2} = \rho_{8j+2} \text{ for } s'+1 \le 4j \le t'-1 \end{cases} \tag{9}$$

We set A as the set $\{i \in \mathbf{N} \mid s' \le i \le t'\}$. (7) and (9) mean that

$$2i, 2i-1 \in A \quad \text{then} \quad \rho_{4i-2} = \rho_{4i}, \tag{10}$$

$$4i, 4i-2 \in A \quad \text{then} \quad \rho_{8i} = \rho_{8i-4}, \tag{11}$$

$$4i-1, 4i+1 \in A \quad \text{then} \quad \rho_{8i-2} = \rho_{8i+2}. \tag{12}$$

Therefore, for $i \in A - \{s', t'-1, t'\}$, $\rho_{2i} = \rho_{2i+2}$. The reason is this: if i is odd, it is trivial from (10); if $i = 4j$ for some j, $\rho_{8j} = \rho_{8j-2} = \rho_{8j+2}$; if $i = 4j-2$ for some j, $\rho_{8j-4} = \rho_{8j} = \rho_{8j-2}$.

We obtain that

$$\rho_{2s'+2} = \rho_{2s'+4} = \cdots = \rho_{2t'-2}.$$

Also, we see

$$2b \ge a+b > 2^{d-1} \text{ and } \frac{2^k-2}{2} - b \le \frac{2^{d-1}-2}{2} - 2^{d-2} < 1 \tag{13}$$

and we have

$$s' = \max\{1, \frac{2^k-2}{2} - b\} = 1.$$

We see again (8) and look into the term of $\tau^2 \otimes \tau^{2^k}$, then we have that $\rho_2 = 0$ and from (10) $\rho_2 = \rho_4$. Hence we have

$$0 = \rho_2 = \rho_4 = \cdots = \rho_{2t'-2},$$

that is,

$$x^*(\alpha_{2^k-2}) = \rho_{2t'}\tau^{2t'} \otimes \tau^{(2^k-2)-2t'}. \tag{14}$$

If $2a \geq 2^k - 4$ then we have

$$t' = \min\{\frac{2^k-4}{2}, a\} = 2^{k-1} - 2$$

and from (10)

$$\rho_{2t'-2} = \rho_{2t'},$$

that is ,

$$x^*(\alpha_{2^k-2}) = 0.$$

Therefore we can assume

$$2a < 2^k - 4, \tag{15}$$

that is, $t' = a$. Here if $2a = 2^{k-1}-2$, then by (14) $x^*(\alpha_{2^k-2}) = \tau^{2^{k-1}-2} \otimes \tau^{2^{k-1}}$ or 0 as asserted. Hence what we have to prove prove is that if $2a \neq 2^{k-1} - 2$ then $\rho_{2t'} = 0$.

We set $p(2a)$ so that $2^{p(2a)}$ is the greatest power of 2 which devides $2a+2$.

Let $p := p(2a)$. We remark that $p \leq k - 2$ since, if it were not, by (15) $2a = 2^{k-1} - 2$. Using Sq^{2^p}, we see

$$\begin{aligned} \mathrm{Sq}^{2^p} x^*(\alpha_{2^k-2}) &= x^*(\alpha_{2^k+2^p-2}) \\ &= \mathrm{Sq}^{2^{k-1}} \mathrm{Sq}^{2^p} x^*(\alpha_{2^{k-1}-2}) \\ &= 0. \end{aligned}$$

Thus it follows that

$$\begin{aligned} \mathrm{Sq}^{2^p}(\rho_{2t'}\tau^{2a} \otimes \tau^{(2^k-2)-2a}) &= \rho_{2t'}\tau^{2a} \otimes \mathrm{Sq}^{2^p}\tau^{2^k-2-2a} \\ &= \rho_{2t'}\tau^{2a} \otimes \tau^{2^k+2^p-2-2a} \\ &= 0 \end{aligned}$$

Here $\tau^{2a} \otimes \tau^{2^k+2^p-2-2a} \neq 0$ since by (3) and (4)

$$\begin{aligned} 2b &> 2^d - 2a \\ &\geq 2 \cdot 2^k - 2a \\ &> 2^k + 2^p - 2 - 2a. \end{aligned} \tag{16}$$

Thus $\rho_{2t'} = 0$, that is, $x^*(\alpha_{2^k-2}) = 0$ as asserted.

Next we shall prove b). Let $x^*(\alpha_{2^{k-1}-2}) = \tau^{2^r-2} \otimes \tau^{2^{k-1}-2^r}$, $r+2 \leq k \leq d-1$ and $2a = 2^r - 2$. Then

$$\begin{aligned} \mathrm{Sq}^i x^*(\alpha_{2^{k-1}-2}) &= \tau^{2^r-2} \otimes \mathrm{Sq}^i(\tau^{2^{k-1}-2^r}) \\ &= \binom{2^{k-1}-2^r}{i} \tau^{2^r-2} \otimes \tau^{2^{k-1}-2^r+i}. \end{aligned}$$

Here we remark that $r \geq 2$. For, if $r = 2$, by a) $x^*(\alpha_{2^i-2}) = 0$ for $3 \leq i \leq d-1$. Thus $\mathrm{Sq}^4 x^*(\alpha_{2^{k-1}-2}) = 0$ and we obtain

$$\begin{aligned} \mathrm{Sq}^1(x^*(\alpha_{2^k-2})) &= x^*(\mathrm{Sq}^1 \alpha_{2^k-2}) = 0, \\ \mathrm{Sq}^2(x^*(\alpha_{2^k-2})) &= x^*(\alpha_2^{2^{k-1}}) = 0, \\ \mathrm{Sq}^4(x^*(\alpha_{2^k-2})) &= \mathrm{Sq}^{2^{k-1}} \mathrm{Sq}^4 x^*(\alpha_{2^{k-1}-2}) = 0. \end{aligned}$$

Then it follows from the previous argument in a) that

$$x^*(\alpha_{2^k-2}) = \rho \tau^{2^r-2} \otimes \tau^{2^k-2^r},$$

where $\rho \in \mathbf{Z}/2\mathbf{Z}$.

Next using Sq^{2^r}, we have

$$\begin{aligned} \mathrm{Sq}^{2^r} x^*(\alpha_{2^k-2}) &= \rho \mathrm{Sq}^{2^r}(\tau^{2^r-2} \otimes \tau^{2^k-2^r}) \\ &= \rho \tau^{2^r-2} \otimes \tau^{2^k}, \end{aligned}$$

while

$$\begin{aligned} \mathrm{Sq}^{2^r} x^*(\alpha_{2^k-2}) &= x^*(\alpha_{2^k+2^r-2}) \\ &= x^*(\mathrm{Sq}^{2^{k-1}} \mathrm{Sq}^{2^r} \alpha_{2^{k-1}-2}) \\ &= \mathrm{Sq}^{2^{k-1}} \mathrm{Sq}^{2^r} x^*(\alpha_{2^{k-1}-2}) \\ &= \tau^{2^r-2} \otimes \tau^{2^k}. \end{aligned}$$

Here $\tau^{2^r-2} \otimes \tau^{2^k} \neq 0$ since

$$\begin{aligned} 2a &= 2^r - 2 \\ 2b &= 2(a+b) - 2a \\ &> 2^d - 2^r + 2 \\ &\geq 2^{d-1} \\ &\geq 2^k. \end{aligned} \tag{17}$$

Therefore $\rho = 1$ and

$$x^*(\alpha_{2^k-2}) = \tau^{2^r-2} \otimes \tau^{2^k-2^r}.$$

Thus lemma 4.11 is proved.

Lemma 4.12 *In the case* I) *if* $x^*(\alpha_6) = 0$ *then* $x^*(\alpha_{2(a+b)}) = 0$.

Proof. By Lemma 4.11

$$x^*(\alpha_{2^{d-1}-2}) = 0$$

or

$$2a = 2^r - 2 \text{ and } x^*(\alpha_{2^{d-1}-2}) = \tau^{2^r-2} \otimes \tau^{2^{d-1}-2^r}.$$

Since

$$x^*(\alpha_{2(a+b)}) = \mathrm{Sq}^{2^{d-1}} \mathrm{Sq}^{2(a+b)-(2^d-2)} x^*(\alpha_{2^{d-1}-2}),$$

if $x^*(\alpha_{2(a+b)}) \neq 0$ then $x^*(\alpha_{2^{d-1}-2}) \neq 0$ and $2(a+b) \equiv -2 \bmod 2^r$. But if $2(a+b) \equiv -2 \bmod 2^r$ then

$$\binom{2(a+b)}{2a} = \binom{2(a+b)}{2^r-2} \equiv 1 \bmod 2.$$

Thus if $\binom{2(a+b)}{2a} \equiv 0 \bmod 2$ and $x^*(\alpha_6) = 0$ then

$$x^*(\alpha_{2(a+b)}) = 0.$$

Q.E.D.

Now we consider the case II) we start from the next lemma.

Lemma 4.13 *Assume $i+j = 2^d - 2$ for some $d \in \mathbf{N}$, $d > 3$, i and j are even, $i, j \geq 2$ and*

$$i = \sum_{k=1}^{d-1} \epsilon_k 2^k,$$

where $\epsilon_k = 0$ or 1. Then

$$\mathrm{Sq}^{2^p} \tau^i \otimes \tau^j = \begin{cases} \tau^{i+2^p} \otimes \tau^j & \epsilon_p = 1 \\ \tau^i \otimes \tau^{j+2^p} & \epsilon_p = 0 \end{cases}$$

for $1 \leq p \leq d-1$ where $\tau^i \otimes \tau^j \in \mathrm{H}^{2^d-2}(\mathbf{RP}^\infty \wedge \mathbf{RP}^\infty)$.

Proof. We use induction. Let $\overline{\epsilon_k} = 1 - \epsilon_k$. Then $j = \sum_{k=1}^{d-1} \overline{\epsilon_k} 2^k$.

The statement is true for $p = 1$. Let we assume that the statement is true for $\mathrm{Sq}^{2^{p-1}}$ and also $\epsilon_{p-1} = 1$. Then

$$\begin{aligned} \mathrm{Sq}^{2^{p-1}} \tau^i \otimes \tau^j &= \sum_{l=0}^{2^{p-1}} (\mathrm{Sq}^l \tau^i) \otimes (\mathrm{Sq}^{2^{p-1}-l} \tau^j) \\ &= \sum_{l=0}^{2^{p-1}} \binom{i}{l} \binom{j}{2^{p-1}-l} \tau^{i+l} \otimes \tau^{j+2^{p-1}-l} \\ &= \tau^{i+2^{p-1}} \otimes \tau^j, \end{aligned}$$

that is,

$$\binom{i}{l}\binom{j}{2^{p-1}-l} = \begin{cases} 0 & 0 \leq l \leq 2^{p-1}-1 \\ 1 & l = 2^{p-1}. \end{cases}$$

Hence

$$\begin{aligned} \mathrm{Sq}^{2^p} \tau^i \otimes \tau^j &= \sum_{l=0}^{2^p} \binom{i}{l}\binom{j}{2^p-l} \tau^{i+l} \otimes \tau^{j+2^p-l} \\ &= \sum_{l=0}^{2^{p-1}} \binom{i}{l}\binom{j}{2^p-l} \tau^{i+l} \otimes \tau^{j+2^p-l} + \sum_{l=0}^{2^{p-1}} \binom{i}{2^{p-1}+l}\binom{j}{2^{p-1}-l} \tau^{i+2^{p-1}+l} \otimes \tau^{j+2^{p-1}-l} \\ &= \sum_{l=1}^{2^{p-1}} \binom{i}{l}\binom{\overline{\epsilon_{p-1}}}{1}\binom{j}{2^{p-1}-l} \tau^{i+l} \otimes \tau^{j+2^p-l} \\ &\quad + \sum_{l=0}^{2^{p-1}-1} \binom{\epsilon_{p-1}}{1}\binom{i}{l}\binom{j}{2^{p-1}-l} \tau^{i+2^{p-1}+l} \otimes \tau^{j+2^{p-1}-l} \\ &\quad + \binom{j}{2^p} \tau^i \otimes \tau^{j+2^p} + \binom{i}{2^p} \tau^{i+2^p} \otimes \tau^j \\ &= \binom{\overline{\epsilon_p}}{1} \tau^i \otimes \tau^{j+2^p} + \binom{\epsilon_p}{1} \tau^{i+2^p} \otimes \tau^j \end{aligned}$$

as asserted. And even if $\epsilon_{p-1} = 1$, it can be proved in the same manner.

Q.E.D.

Lemma 4.14 *Let $b \geq a$. In the case* II), *if $x^*(\alpha_6) = 0$, then*

$$x^*(\alpha_{2^d-2}) = \begin{cases} \rho \sum_{i=1}^{(2^d-4)/2} \tau^{2i} \otimes \tau^{(2^d-2)-2i} + \rho' \tau^{2^{d-1}-2} \otimes \tau^{2^{d-1}} \\ \qquad \textit{where } 2a = 2^{d-1} - 2 \textit{ if } \rho' = 1 \\ \textit{or} \\ \tau^{2^r-2} \otimes \tau^{2^d-2^r} \textit{ and } 2a = 2^r - 2,\ 3 \leq r \leq d-2. \end{cases}$$

Proof. We start from the computation of $x^*(\alpha_{2^{d-1}-2})$. By lemma 4.11

$$x^*(\alpha_{2^{d-1}-2}) = \begin{cases} 0 \\ \text{or} \\ \tau^{2^r-2} \otimes \tau^{2^{d-1}-2^r} \text{ in this case } 2a = 2^r - 2,\ 3 \leq r \leq d-1. \end{cases}$$

Next we consider $x^*(\alpha_{2^d-2})$. Since

$$\begin{aligned} \mathrm{Sq}^1(x^*(\alpha_{2^d-2})) &= x^*(\mathrm{Sq}^1 \alpha_{2^d-2}) = 0, && (18) \\ \mathrm{Sq}^2(x^*(\alpha_{2^d-2})) &= x^*(\alpha_2^{2^{d-1}}) = 0, && (19) \\ \mathrm{Sq}^4(x^*(\alpha_{2^d-2})) &= \mathrm{Sq}^{2^{d-1}} \mathrm{Sq}^4 x^*(\alpha_{2^{d-1}-2}) = 0, && (20) \end{aligned}$$

as in the proof of Lemma 4.11, we have

$$x^*(\alpha_{2^d-2}) = \rho \tau^{2s} \otimes \tau^{(2^d-2)-2s} + \rho' \sum_{i=s+1}^{t-1} \tau^{2i} \otimes \tau^{(2^d-2)-2i} + \rho'' \tau^{2t} \otimes \tau^{(2^d-2)-2t},$$

where

$$\begin{aligned} s &= \max\{1, \frac{2^d-2}{2} - b\}, \\ t &= \min\{\frac{2^d-4}{2}, a\}. \end{aligned}$$

Firstly we assume $x^*(\alpha_{2^{d-1}-2}) = 0$. And we shall prove $\rho = \rho'$. If $s = 1$ then the equation $\mathrm{Sq}^2 x^*(\alpha_{2^d-2}) = 0$ means $\rho = \rho'$. Thus we assume $s = \frac{2^d-2}{2} - b$, that is,

$$2b \leq 2^d - 4. \qquad (21)$$

Here we remark that by (4),

$$\begin{aligned} 2b &\geq a+b && (22)\\ &> 2^{d-1}-2 && (23)\end{aligned}$$

Let $q := p(2b)$ then (21) and (23) mean $q \leq d-2$. Also

$$\mathrm{Sq}^{2^q} x^*(\alpha_{2^d-2}) = \mathrm{Sq}^{2^{d-1}} \mathrm{Sq}^{2^q} x^*(\alpha_{2^{d-1}-2}) = 0.$$

Thus, by Lemma 4.13, compare the term of $\tau^{(2^d-2)-2b+2^q} \otimes \tau^{2b}$ in $\mathrm{Sq}^{2^q} x^*(\alpha_{2^d-2})$ and we obtain

$$(\rho+\rho')\tau^{(2^d-2)-2b+2^q} \otimes \tau^{2b} = 0. \quad (24)$$

Here we remark that $(2^d-2)-2b+2^q \leq 2a$ by (4). Thus (24) means $\rho' = \rho''$. Therefore

$$x^*(\alpha_{2^d-2}) = \rho' \sum_{i=s}^{t-1} \tau^{2i} \otimes \tau^{(2^d-2)-2i} + \rho'' \tau^{2t} \otimes \tau^{(2^d-2)-2t}.$$

Next we consider the term $\rho''\tau^{2t} \otimes \tau^{(2^d-2)-2t}$. If $2t = 2^d-4$, then by the computation of $\mathrm{Sq}^2 x^*(\alpha_{2^d-2})$ we have $\rho' = \rho''$ and $x^*(\alpha_{2^d-2}) = \sum_{i=s}^{t} \tau^{2i} \otimes \tau^{(2^d-2)-2i}$ or 0 as asserted. Thus we assume $2t = 2a$, that is,

$$2a < 2^d - 4 \quad (25)$$

Let $p := p(2a)$. Here from (25) $p \leq d-1$. And $p = d-1$ if and only if $2a = 2^{d-1}-2$.

If $2a = 2^{d-1}-2$ then

$$x^*(\alpha_{2^d-2}) = \rho' \sum_{i=1}^{(2^d-4)/2} \tau^{2i} \otimes \tau^{(2^d-2)-2i} + (\rho''+\rho')\tau^{2^{d-1}-2} \otimes \tau^{2^{d-1}}.$$

If $p \leq d-2$ then

$$\mathrm{Sq}^{2^p} x^*(\alpha_{2^d-2}) = \mathrm{Sq}^{2^{d-1}} \mathrm{Sq}^{2^p} x^*(\alpha_{2^{d-1}-2}) = 0. \quad (26)$$

By Lemma 4.13 look into the term of $\tau^{2a} \otimes \tau^{(2^d-2)-2a+2^p}$ of (26) and we obtain

$$(\rho'+\rho'')\tau^{2a} \otimes \tau^{(2^d-2)-2a+2^p} = 0. \quad (27)$$

Remark that by (4)

$$(2^d-2)-2a+2^p \leq 2b.$$

Therefore $\rho' = \rho''$ and

$$x^*(\alpha_{2^d-2}) = \rho' \sum_{i=s}^{t} \tau^{2i} \otimes \tau^{(2^d-2)-2i}.$$

Secondly we assume $x^*(\alpha_{2^{d-1}-2}) = \tau^{2^r-2} \otimes \tau^{2^{d-1}-2^r}$ and $2a = 2^r - 2$ and observe $x^*(\alpha_{2^d-2})$ again. We reset

$$x^*(\alpha_{2^d-2}) = \rho\tau^{2s} \otimes \tau^{(2^d-2)-2s} + \rho' \sum_{i=s+1}^{t-1} \tau^{2i} \otimes \tau^{(2^d-2)-2i} + \rho''\tau^{2t} \otimes \tau^{(2^d-2)-2t},$$

where

$$\begin{aligned} s &= \max\{1, \frac{2^d-2}{2} - b\}, \\ t &= \min\{\frac{2^d-4}{2}, a\}. \end{aligned}$$

Then

$$\begin{aligned} 2b &= 2(a+b) - 2a && (28) \\ &\geq (3 \cdot 2^{d-1} - 2) - (2^{d-1} - 2) && (29) \\ &= 2^d && (30) \end{aligned}$$

This means $s = 1$. Thus by the computation of $\mathrm{Sq}^2 x^*(\alpha_{2^d-2})$ we have

$$\rho = \rho'$$

and also by the computation of $\mathrm{Sq}^4 x^*(\alpha_{2^d-2})$ and by (30) we have

$$\rho = 0.$$

Therefore we obtain

$$x^*(\alpha_{2^d-2}) = \rho''\tau^{2^r-2} \otimes \tau^{2^d-2^r}.$$

Finally we have obtained the following result

$$x^*(\alpha_{2^d-2}) = \begin{cases} \rho \sum_{i=1}^{(2^d-4)/2} \tau^{2i} \otimes \tau^{(2^d-2)-2i} + \rho'\tau^{2^{d-1}-2} \otimes \tau^{2^{d-1}} \\ \qquad \text{where } 2a = 2^{d-1} - 2 \text{ if } \rho' = 1 \\ \text{or} \\ \tau^{2^r-2} \otimes \tau^{2^d-2^r} \text{ and } 2a = 2^r - 2,\ 3 \leq r \leq d-2. \end{cases}$$

Q.E.D.

Lemma 4.15 *In the case* II) *if* $x^*(\alpha_6) = 0$ *then* $x^*(\alpha_{2(a+b)}) = 0$.

Proof. By (4)

$$x^*(\alpha_{2(a+b)}) = \mathrm{Sq}^{2(a+b)-(2^d-2)} x^*(\alpha_{2^d-2}).$$

And by Lemma 4.14 we shall prove that

$$\begin{cases} \mathrm{Sq}^{2(a+b)-(2^d-2)}\left(\sum_{i=1}^{(2^d-4)/2} \tau^{2i} \otimes \tau^{(2^d-2)-2i}\right) = 0 \\ \mathrm{Sq}^{2(a+b)-(2^d-2)}(\tau^{2^r-2} \otimes \tau^{2^d-2^r}) = 0 \text{ in case } a = 2^r - 2,\ 3 \le r \le d-1. \end{cases}$$

Since

$$\sum_{i=1}^{(2^d-4)/2} \tau^{2i} \otimes \tau^{(2^d-2)-2i} = x_0^*(\alpha_{2^d-2}),$$

it follows that

$$\begin{aligned} \mathrm{Sq}^{2(a+b)-(2^d-2)}\Big(\sum_{i=1}^{(2^d-4)/2} \tau^{2i} \otimes \tau^{(2^d-2)-2i}\Big) &= \mathrm{Sq}^{2(a+b)-(2^d-2)} x_0^*(\alpha_{2^d-2}) \\ &= x_0^*(\alpha_{2(a+b)}) \\ &= \tbinom{2(a+b)}{2a} \tau^{2a} \otimes \tau^{2b} \\ &= 0. \end{aligned}$$

Also

$$\begin{aligned} & \mathrm{Sq}^{2(a+b)-(2^d-2)}(\tau^{2^r-2} \otimes \tau^{2^d-2^r}) \\ = \ & \tau^{2^r-2} \otimes \tbinom{2^d-2^r}{2(a+b)-(2^d-2)} \tau^{2(a+b)-(2^d-2)} \\ = \ & \begin{cases} \tau^{2^r-2} \otimes \tau^{2(a+b)-(2^r-2)} & \text{if } 2(a+b) \equiv -2 \bmod 2^r \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

But if $2(a+b) \equiv -2 \bmod 2^r$ then

$$\tbinom{2(a+b)}{2a} = \tbinom{2(a+b)}{2^r-2} \equiv 1 \bmod 2.$$

Thus if $\binom{2(a+b)}{2a} \equiv 0 \bmod 2$ then $x^*(\alpha_{2(a+b)}) = 0$.

Q.E.D.

Now we shall finish the proof of Theorem 4.7. Let $x : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega_0\mathbf{SO}$ be an arbitrary map, $n > 1$, $m > 1$ and $\binom{n+m-2}{n-1} \equiv 0 \bmod 2$. If $x^*(\alpha_6) = 0$ then by Lemma 4.12, Lemma 4.15 we obtain $x^*(\alpha_{n+m-2}) = 0$. Therefore we assume $x^*(\alpha_6) \neq 0$. Then from Lemma 4.9

$$x^*(\alpha_6) = \tau^2 \otimes \tau^4 + \tau^4 \otimes \tau^2.$$

Let $x + x_0 : \mathbf{RP}^{n-1} \wedge \mathbf{RP}^{m-1} \to \Omega_0\mathbf{SO}$ be a map which is contained in the homotopy class $[x] + [x_0]$. Since $\Omega_0\mathbf{SO}$ is an H-space and it is known that $\alpha_{2i} \in \mathrm{H}^*(\Omega_0\mathbf{SO})$ are primitive elements,

$$(x + x_0)^*(\alpha_6) = 2(\tau^2 \otimes \tau^4 + \tau^4 \otimes \tau^2) = 0.$$

Therefore

$$(x + x_0)^*(\alpha_{n+m-2}) = 0,$$

while

$$\begin{aligned} (x + x_0)^*(\alpha_{n+m-2}) &= x^*(\alpha_{n+m-2}) + x_0^*(\alpha_{n+m-2}) \\ &= x^*(\alpha_{n+m-2}) + \tbinom{n+m-2}{n-1}\tau^{n-1} \otimes \tau^{m-1} \\ &= x^*(\alpha_{n+m-2}). \end{aligned}$$

Finally we obtained that $x^*(\alpha_{n+m-2}) = 0$ and Theorem 4.7 is proved.

References

[1] I.M.James & E.Thomas, *Homotopy-commutativity in rotation groups,* Topology **1**(1962),121-124.

[2] I.M.James, *The topology of Stiefel manifolds,* London Math. Soc. Lecture Notes **24**, Cambridge University Press(1976).

[3] R.Bott, *A note on the Samelson product in the classical groups,* Comment. Math. Helv. **34**(1960),249-256.

[4] J.H.C.Whitehead, *On the groups $\pi_r(V_{n,m})$ and sphere-bundles,* Proc. Lond. Math. Soc. **48**(1944),243-291.

[5] S.Y.Husseini, *A note on the intrinsic join of Stiefel manifolds,* Comment. Math. Helv. **38**(1963),26-30.