On the other hand, for a rational map f of degree d such that $\infty \in \Omega_f$ the following inequality is known to hold and be best possible [2]:

$$\text{H-dim}J_f \geq \frac{\log d}{\log K_0},$$

where $K_0 = \max_{z \in J_f} |f'(z)|$.

In our present case, $d = 2$ and $K_0 = 2\alpha = 1 + \sqrt{1 - 4c}$, thus we have

$$\text{H-dim}J_c \geq \frac{\log 2}{\log(1 + \sqrt{1 - 4c})}.$$

Therefore, it seems that our estimate is sufficiently good at least in case $c < -2$.

References

1. Ahlfors, L. V. *Conformal Invariants*, McGraw Hill, New York (1973).
2. Beardon, A. F. *Iteration of Rational Functions*, No. 132 in Grad. Texts Math., Springer-Verlag (1991).
3. Beardon, A. F. and Pommerenke, Ch. The Poincaré metric of plane domains, *J. London Math. Soc. (2)*, **18** (1978), 475–483.
4. Carleson, L. and Gamelin, T. W. *Complex Dyamics*, Springer-Verlag (1993).
5. Hempel, J. A. The Poincaré metric on the twice punctured plane and the theorems of Landau and Schottky, *J. London Math. Soc. (2)*, **20** (1979), 435–445.
6. Hinkkanen, A. Julia sets of rational functions are uniformly perfect, *Math. Proc. Cambridge Philos. Soc.*, **113** (1993), 543–559.
7. Järvi, P. and Vuorinen, M. Uniformly perfect sets and quasiregular mappings, *J. London Math. Soc.*, **54** (1996), 515–529.
8. Mañé, R. and da Rocha, L. F. Julia sets are uniformly perfect, *Proc. Amer. Math. Soc.*, **116** (1992), 251–257.
9. Pommerenke, Ch. Uniformly perfect sets and the Poincaré metric, *Ark. Math.*, **32** (1979), 192–199.
10. Pommerenke, Ch. On uniformly perfect sets and Fuchsian groups, *Analysis*, **4** (1984), 299–321.
11. Sugawa, T. Uniform perfectness of the limit sets of Kleinian groups, Preprint (1997).
12. Sugawa, T. Various domain constants related to uniform perfectness, Preprint (1997).
13. Tsuji, M. *Potential Theory*, Maruzen, Tokyo (1959).

Department of Mathematics, Kyoto University, 606-01 Sakyo-ku, Kyoto City, Japan

E-mail address: sugawa@kusm.kyoto-u.ac.jp

that $|dz| \geq (|dr| + r|d\theta|)/\sqrt{2}$, where $z = re^{i\theta}$. Put $a_0 = \min|\beta|$, then by (6.1) we have

$$\begin{aligned} 2h(a) &\geq \int_\beta \frac{|dz|}{2|z|(-\log|z| + K)} \geq \int_\beta \frac{|dr| + r|d\theta|}{2r\sqrt{2}(-\log r + K)} \\ &\geq \frac{2}{2\sqrt{2}} \log\left(\frac{-\log a_0 + K}{-\log a + K}\right) + \frac{1}{2\sqrt{2}} \frac{2\pi}{-\log a_0 + K} \\ &\geq \frac{1}{2\sqrt{2}} \frac{\pi/\sqrt{2}}{-\log a + K}, \end{aligned}$$

because $K > \pi$.

Next, we consider the case $a > 1$. Since the Möbius transformation $z \mapsto 1/z$ preserves D_0, we have $h(a) = h(1/a)$, thus we have $h(a) \geq \pi/2\sqrt{2}(\log a + K)$. Therefore, for any $a > 0$, we have

$$\iota_{D_0}(-a) \geq \frac{\pi}{2\sqrt{2}(|\log a| + K)}.$$

Letting $D = \mathbb{C} \setminus [0, \infty)$, we can estimate the injectivity radius of Ω_c at $T^{-1}(-a)$ as follows:

$$\iota_{\Omega_c}(T^{-1}(-a)) = \iota_{T(\Omega_c)}(-a) \geq \inf_{w \in \partial D} d_0(-a, w) = \iota_{D_0}(-a) \geq \frac{\pi}{2\sqrt{2}(|\log a| + K)}.$$

Hence,

$$\begin{aligned} C &:= \min\{2d_{\Omega_c}(c, \infty), 4\iota_{\Omega_c}(c), 4\iota_{\Omega_c}(\infty)\} \\ &\geq \min\left\{\log\frac{\log(3+t) + K}{K} + \left|\log\frac{\log\frac{4+t}{t(3+t)} + K}{K}\right|, \frac{\sqrt{2}\pi}{\log(3+t) + K}, \frac{\sqrt{2}\pi}{\log\frac{4+t}{t(3+t)} + K}\right\} \\ &= \frac{\sqrt{2}\pi}{\log m + K}, \end{aligned}$$

where $m = \max\{3 + t, \frac{4+t}{t(3+t)}\}$.

By Corollary 2.3, we have proved the following: *For $c < -2$ it holds that*

$$L_{\Omega_c} \geq \frac{\sqrt{2}\pi}{\log m + K},$$

where $m = \max\{\sqrt{1-4c}, \frac{\sqrt{1-4c}+1}{\sqrt{1-4c}(\sqrt{1-4c}-3)}\}$ and K is the Hempel's constant in (6.1).

In view of Theorem 5.2, we can see that for $c < -2$,

$$\text{H-dim}J_c \geq \frac{\sqrt{2}\log 2}{\pi(\log m + K) + \sqrt{2}\log 3}.$$

Hempel [5]: The hyperbolic metric $\rho_0(z)|dz|$ of D_0 satisfies

$$\rho_0(z) \geq \frac{1}{2|z|(|\log|z|| + K)}, \tag{6.1}$$

where $K = \Gamma(\frac{1}{4})^4/4\pi^2 = 4.3768796\cdots$, and the equality occurs if $z = -1$. Note that this estimate is efficient only on the half plane $\mathrm{Re} z \leq \frac{1}{2}$, otherwise we have only to use the functional equation $\rho_0(1-z) = \rho_0(z)$.

In order to find out a thrice punctured sphere containing Ω, we have only to specify three points a_1, a_2, a_3 in the Julia set, for example, repelling periodic points. In our present case, any periodic point is repelling since $c \notin \mathcal{M}$. For example, fixed points of f_c are solutions of the equation: $z^2 + c = z$, thus $(1 \pm \sqrt{1-4c})/2$. We note that if α is in J_c, so is $-\alpha$. If we selected the three points a_1, a_2, a_3 in the Julia set, let T be the Möbius transformation mapping a_1, a_2 and a_3 to $0, 1$ and ∞, respectively. Then, $d_{\Omega_c}(c, \infty) \geq d_{\widehat{\mathbb{C}}\setminus\{a_1,a_2,a_3\}}(c, \infty) = d_0(T(c), T(\infty))$, where d_0 denotes the hyperbolic distance in D_0, however it seems impossible to estimate $\iota_{\Omega_c}(c)$ and $\iota_{\Omega_c}(\infty)$ by only the data a_j.

For simplicity, we further assume that $c < -2$ in the following. Set $\alpha = (1 + \sqrt{1-4c})/2$ and $\beta = 1 - \alpha$, then these are fixed points of $f = f_c$. Then we see that $S = \cup_{n=1}^{\infty} f^{-n}(\alpha) \subset [-\alpha, \alpha]$, hence $J_c \subset [-\alpha, \alpha]$ since $\overline{S} = J_c$.

For later convenience, we set $t = \sqrt{1-4c} - 3 > 0$. Let $T(z) = \frac{(\beta-\alpha)(z+\alpha)}{(\beta+\alpha)(z-\alpha)} = (3 + t)\frac{\alpha+z}{\alpha-z}$. Then $T(\infty) = -(3+t)$ and $T(c) = -\frac{t(3+t)}{4+t}$. We also note that $T(J_c) \subset [0, \infty]$. Using (6.1), we can calculate as

$$d_{\Omega_c}(c, \infty) \geq d_0(T(c), T(\infty)) \geq \frac{1}{2}\log\frac{\log(3+t) + K}{K} + \frac{1}{2}\left|\log\frac{\log\frac{4+t}{t(3+t)} + K}{K}\right|.$$

Next, we shall estimate the injectivity radii of Ω_c at c and ∞. As a preparation, we consider the quantity $h(a) = \iota_{D_0}(-a)$ for $a > 0$. First we assume that $0 < a \leq 1$. Let Δ be the domain defined by $\{\tau \in \mathbb{H}; 0 < \mathrm{Im}\tau < 1, |\tau - \frac{1}{2}| > \frac{1}{2}\}$, and $\lambda : \Delta \to \mathbb{H}$ the conformal homeomorphism from Δ onto the upper half plane $\mathbb{H}$ which maps $0, 1, \infty$ to $1, \infty, 0$, respectively. We denote by $g : \mathbb{H} \to \Delta$ the inverse map of λ. Then, as is well-known, λ is analytically continued to the universal covering map of D_0 from $\mathbb{H}$ by the reflection principle, in particular, $1/2\mathrm{Im}\tau = \rho_0(\lambda(\tau))|\lambda'(\tau)|$. The map λ is nothing but the classical elliptic modular function. For a point $\tau = (e^{i\theta} + 1)/2 \in g((1,2])$ $(\pi/2 \leq \theta < \pi)$ we can see that $d_{\mathbb{H}}(\tau, g((0,1))) \leq d_{\mathbb{H}}(\tau, g((-\infty, 0)))$ and that the shortest hyperbolic segment γ connecting τ and $g((0,1)) = \{ti; t > 0\}$ is contained in $\{\tau \in \Delta; \mathrm{Re}\tau \leq \frac{1}{2}\}$. Noting that $\lambda(\{\tau \in \Delta; \mathrm{Re}\tau = \frac{1}{2}\}) = \{z \in \mathbb{H}; |z-1| = 1\}$, we have $h(a) = \iota_{D_0}(-a) = \iota_{D_0}(1+a) = \int_{\lambda_*\gamma} \rho_0(z)|dz|$ and $\lambda_*\gamma$ is contained in $\{z \in \overline{\mathbb{H}}; |z-1| \leq 1\}$. Denote by β the closed curve obtained as the union of $1 - \lambda_*\gamma$ and its complex conjugate, then $|\beta| \leq 1$ and $2h(a) = \int_\beta \rho_0(z)|dz|$. Note

is true in our situation, indeed the validity of this inequality characterize the uniform perfectness.

Theorem 5.3 (cf. [12]**).** *For a rational map f of degree at least two, we set $L = L_{\Omega_f}$. If $\infty \in J_f$ then we have*

$$\frac{1}{4}\tanh L/2 \leq \inf_{z\in\Omega_f} \rho_{\Omega_f}(z)\delta_{\Omega_f}(z) \leq \frac{\sqrt{3}L}{\sqrt{\pi^2+4L^2}}.$$

For another applications and characterizations of uniform perfectness, see [12] and its reference.

6. Quadratic polynomials

In this section, as the simplest example, we shall consider the quadratic polynomials $f(z) = f_c(z) = z^2 + c$ and attempt to give a concrete lower bound for the uniform perfectness constant L_{Ω_f} (abbreviated by L_c) of the Jula set J_c of f_c in terms of the parameter c. (For a general rational map f, we may estimate L_{Ω_f} in the similar way as below, in principle.) For general result of the dynamics of quadratic polynomials, the reader will find a good account in the book [4] by Carleson and Gamelin.

Since f_c is a polynomial, the infinity ∞ is a superattracting fixed point of f_c. And 0 is a unique finite critical point of f_c and c is the corresponding critical value. Let $\mathcal{M}$ denote the Mandelbrot set $\{c \in \mathbb{C};\, (f_c^n(0))_{n=1,2,\cdots} \text{ is a bounded sequence}\}$. As is well-known, $c \in \mathcal{M}$ if and only if the Julia set J_c is connected, in which case $L_c = +\infty$ since $\Omega = \Omega_c := \widehat{\mathbb{C}}\setminus J_c$ is simply connected, thus we have nothing to do. So we assume that $c \notin \mathcal{M}$ in the sequel. In this case, the Julia set J_c is a Cantor set, therefore the Fatou set Ω_c is connected. By Corollary 2.3, it is sufficient to estimate $d_\Omega(c,\infty)$, $\iota_\Omega(c)$ and $\iota_\Omega(\infty)$ from below. To accomplish it, we may utilize the monotonicity property of the hyperbolic metrics. If we find a hyperbolic domain $\widetilde{\Omega}$ containing Ω which is easier to estimate its hyperbolic metric, then $\rho_\Omega \geq \rho_{\widetilde{\Omega}}$ by the Schwarz-Pick lemma. Therefore, it holds that $d_\Omega(a,b) \geq d_{\widetilde{\Omega}}(a,b)$ for any $a, b \in \Omega$. On the other hand, it is not always true that $\iota_\Omega(a) \geq \iota_{\widetilde{\Omega}}(a)$, but we can avoid this difficulty as follows. Fix $a \in \Omega$. Let D be an arbitrary simply connected subdomain of Ω containing a, then we have

$$\iota_\Omega(a) \geq \inf_{w\in\partial D} d_\Omega(a,w) \geq \inf_{w\in\partial D} d_{\widetilde{\Omega}}(a,w).$$

The most useful (but not necessarily sufficient) domain D is thought to be a thrice punctured sphere, since it has been studied for a long time and its hyperbolic metric can be expressed almost explicitly (see, for example, [1] and [3]). Any thrice punctured sphere is conformally (indeed, Möbius) equivalent to the canonical one: $D_0 = \widehat{\mathbb{C}} \setminus \{0, 1, \infty\}$. The following is the precise version of Landau's theorem due to

In particular, we have $\ell_\Omega(\alpha) \geq L_{A_j} \geq C$.

Otherwise, there exists an integer $n \geq 0$ such that α_n is non-trivial while α_{n+1} is trivial in Ω. Since $\ell_\Omega(\alpha) \geq \ell_\Omega(\alpha_n)$, we may assume that $n = 0$, in other words, α is non-trivial but $\beta = f_*(\alpha)$ is contractible in Ω. If $f : U \to W := f(U)$ is unbranched covering, a homotopy joining β and a constant curve in W can be lifted to a homotopy joining α and a constant curve in U via f, thus α is contractible in U, this is a contradiction. Therefore, $f : U \to W$ must be branched, i.e., $U = U_j$ for some $j = 1, \cdots, s$. Now we can apply the key lemma in the previous section! By Lemma 3.2, we have $\ell_\Omega(\alpha) \geq \min\{a_j, b_j\} \geq C$, thus the proof is now completed.

5. Applications

We now present some applications of the main result. For a rational map $f : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ of degree $d \geq 2$, we have seen that $L_{\Omega_f} \geq C$, where Ω_f denotes the Fatou set $\widehat{\mathbb{C}} \setminus J_f$ and $C > 0$ is the constant which appears in Theorem 2.2 or Corollary 2.3.

First of all, we state a result concerning the Hausdorff dimension. The following theorem is essentially due to Järvi-Vuorinen [7], while a quantitative version as in the following is found in [12].

Theorem 5.1. *The Hausdorff dimension of the Julia set J_f of a rational map f can be estimated as*

$$\text{H-dim}(J_f) \geq \frac{\log 2}{\log(2e^{M^\circ_{\Omega_f}} + 1)} \geq \frac{\log 2}{M^\circ_{\Omega_f} + \log 3} \geq \frac{\log 2}{\pi^2/L_{\Omega_f} + \log 3}.$$

In particular, any rational map of degree≥ 2 has always the Julia set of positive Hausdorff dimension, it is a well-known fact and is also shown in [4] by the uniform perfectness in another method.

The next theorem ensures the regurality of the Julia set in the sense of Dirichlet (cf. [13]) by Wiener's criterion.

Theorem 5.2 (Pommerenke [9]. See also [12]). *Let f be a rational map of degree at least two. Then, for each point $a \in J_f$ and $0 < r < \text{diam} J_f$, it holds that* $\text{Cap}(J_f \cap B(a, r)) \geq cr$, *where $c \leq 1$ is a constant satisfying* $\log 1/c \leq M^\circ_{\Omega_f} + 7 \log 2$, diam *stands for the Euclidean diameter,* Cap *the logarithmic capacity and $B(a, r)$ is the closed disk centered at a with radius r.*

In fact, the above property characterizes the uniform perfectness of the set J_f (see [9]). Similarly, we can state a characterization of uniform perfectness of the closed set in terms of the Hausdorff contents [12].

Finally, we mention the esimate of the hyperbolic (or Poincaré) metric $\rho(z)|dz| = \rho_\Omega(z)|dz|$ of $\Omega = \Omega_f$ by the distance function $\delta(z) = \delta_\Omega(z) = \text{dist}(z, J_f) = \inf_{a \in J_f} |z - a|$, provided that $\infty \in J_f$. It is always true that $\rho(z) \leq 1/\delta(z)$. On the other hand, if Ω is simply connected, it is well-known that $\rho(z) \geq 1/4\delta(z)$, while this kind of inequality need not be valid in general, even in the case $\partial\Omega$ is a perfect set. But this

Let $u_n^{\pm}$ $(n = 1, 2, \cdots)$ be an increasing sequence of positive numbers which converges to $\theta_{\pm}$ for each signature, and $\tilde{\beta}_n$ be the curve obtained from $\tilde{\beta}$ by replacing its subarcs $\tilde{\beta}|_{I_n^+}, \tilde{\beta}|_{I_n^-}$ by the hyperboic segments $S_{u_n^+}, S_{-u_n^-}$, respectively, where $I_n^{\pm}$ denotes the subinterval $\{e^{i\theta}; \pm\theta \in [0, u_n^{\pm}]\}$ of S^1. And set $\beta_n = p_*\tilde{\beta}_n$ for each $n = 1, 2, \cdots$. By construction, β_n is freely homotopic to β in $W \setminus f(\mathrm{Crit}(f))$, and it holds that $\ell_W(\beta_n) = \ell_\Delta(\tilde{\beta}_n) \leq \ell_\Delta(\tilde{\beta}) = \ell_W(\beta)$. Let α_n be the lift of β_n via f with initial point $\alpha_n(1) = \alpha(1)$, then α_n is closed and homotopic to α.

So, we can take a unique closed lift α_n of β_n via f so that α_n is freely homotopic to α. Let $\alpha' = \lim \alpha_n$ and $\beta' = f_*\alpha'$. Then, we note that $\ell_W(\beta') = \lim \ell_W(\beta_n) \leq \ell_W(\beta)$. Further, we can see that $\beta' \in \mathcal{S}(v_+, v_-)$ or $\beta' \in \mathcal{T}(v_+)$ according to that $v_+ \neq v_-$ or not. Therefore, we can compute as follows.

$$\ell_U(\alpha) \geq \ell_W(\beta) \geq \ell_W(\beta') \geq \min\{a(v_+, v_-), b(v_+)\} \geq \min\{a, b\}.$$

□

Corollary 3.3. *Let $f : U \to W$ be a holomorphic branched covering between hyperbolic Riemann surfaces U and W. Then it follows that*

$$L_U \geq \min\{L_W, a, b\},$$

where the constants a, b are as in the above.

4. Proof of the Main Theorem

Let α be a non-trivial closed curve in $\Omega = \Omega_f$. In order to prove our main theorem, we should show that $\ell_\Omega(\alpha) \geq C$, where $C = \min\{a_1, \cdots, a_s, b_1, \cdots, c_s, L_{A_1}, \cdots, L_{A_t}\}$. We denote by α_n the image $f^n \circ \alpha = (f^n)_*(\alpha)$ of α under the n-th iterate of f. We note here that $\ell_\alpha \geq \ell_\Omega(\alpha_1) \geq \ell_\Omega(\alpha_2) \cdots$ by the Schwarz-Pick lemma. Let U be the component of Ω containing α. Then, by Sullivan's No Wandering Domains Theorem, U is eventually periodic, i.e., $D = f^k(U)$ is a periodic component for some integer k. As is well-known, a periodic component D is one of the following:

(1) a (super)attracting immedeate basin. In this case, the sequence of curves α_n is attracted to a (super)attracting cycle (in Ω), in particular, α_n is contractible in Ω for sufficiently large n.
(2) a parabolic immedeate basin. In this case, a subsequence of α_n is absorbed by a simply connected attracting petal (in Ω), therefore α_n is contractible in Ω, too, for sufficiently large n.
(3) a Siegel disk. In this time, D is simply connected itself, thus α_k is of course contractible in D.
(4) a Herman ring.

Hence, we can conclude that if α_n is non-trivial for any n, then α_n is contained in a Herman ring A_j for large n. In this case, α_n is freely homotopic to a non-zero multiple of the core curve β of A_j, thus $\ell_\Omega(\alpha_n) \geq \ell_{A_j}(\beta) = L_{A_j}$.

Lemma 3.2. *For a non-trivial loop α in U such that $\beta = f_*\alpha$ is contractible in W, it follows that*

$$\ell_U(\alpha) \geq \min\{a, b\}.$$

Proof. First we show that $\ell_U[\alpha] > 0$. In fact, if $\ell_U[\alpha] = 0$ then α surrounds a puncture of U, in other words, there exists a holomorphic injection $g : \Delta^* = \Delta \setminus \{0\} \to U$ such that α is freely homotopic to ε^n for some integer $n \neq 0$, where $\varepsilon = g(\{|\zeta| = 1/2\})$. As is easily seen, $f(g(\Delta^*))$ is a neighborhood of a puncture of W and thus β is freely homotopic to non-zero multiple of a simple loop around the puncture in W. On the other hand, β is contractible in W, therefore W must be conformally equivalent to the complex plane $\mathbb{C}$, but this is impossible because W is hyperbolic.

Since $\ell_U(\alpha) \geq \ell_U[\alpha] = \ell_U(\alpha_0)$ for the closed geodesic α_0 freely homotopic to α, it suffices to show the claim in the case α is a smooth curve. Approximating α by another smooth curve if necessary, we may further assume that α does not pass through any critical point. Here, we should observe $\ell_U(\alpha) \geq \ell_W(\beta)$ by the Schwarz-Pick lemma: $f^*\rho_W \leq \rho_U$.

Let $p : \Delta \to W$ be a holomorphic universal covering map of W from the unit disk Δ and set $C = p^{-1}(f(\mathrm{Crit}(f)))$. Since β is contractible, a lift $\tilde{\beta} : \mathrm{S}^1 \to \Delta$ of β via p is closed. Let K be the holomorphically convex hull of $\tilde{\beta}(\mathrm{S}^1)$ in Δ. In other words, $K = \Delta \setminus D_0$, where D_0 is the relatively non-compact component of $\Delta \setminus \tilde{\beta}(\mathrm{S}^1)$.

Now we will show that $\#(K \cap C) \geq 2$. If $K \cap C$ is an empty set, then it is clear that β is homotopic to a point with a homotopy in $W \setminus f(\mathrm{Crit}(f))$. Since $f : U \setminus f^{-1}(f(\mathrm{Crit}(f))) \to W \setminus f(\mathrm{Crit}(f))$ is an unbranched covering map, this homotopy can be lifted via f to a homotopy from α to a point, but this contradicts to the assumption that α is non-trivial. Next, suppose that $K \cap C$ consists of one point ζ_0. By assumption, we note that $\zeta_0 \in K \setminus \tilde{\beta}(\mathrm{S}^1)$. Then it is not difficult to see that the loop $\tilde{\beta}$ is freely homotopic to ε^n in $\Delta \setminus C$, where ε is a sufficiently small simple loop around ζ_0 in $\Delta \setminus C$ and n is the winding number of $\tilde{\beta}$ around ζ_0. This implies that β is freely homotopic to $p_*(\varepsilon^n)$ in $\Delta \setminus f(\mathrm{Crit}(f))$, therefore α is freely homotopic to a loop δ with $f_*\delta = p_*(\varepsilon^n)$. In particular, $\ell_U[\alpha] \leq \ell_U(\delta)$ and the length of δ can be arbitrarily small, therefore $\ell_U[\alpha] = 0$, this is not the case. Now we have proved that $\#(K \cap C) \geq 2$.

Here we recall that $\tilde{\beta}$ is parametrized by $\mathrm{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$. For each $\theta \in \mathbb{R}$, we denote by S_θ the hyperbolic segment joining $\tilde{\beta}(1)$ and $\tilde{\beta}(e^{i\theta})$ in Δ. Now we define positive numbers θ_+ and θ_- by

$$\theta_\pm = \max\{\theta \geq 0; S_{\pm u} \cap C = \emptyset \quad \text{for all } u \in [0, \theta)\}.$$

Then, by the above observation, we see that $\theta_+ + \theta_- \leq 2\pi$ and if the equality occurs we have $\#(S_{\theta_+} \cap C) \geq 2$ since $K \subset \bigcup_{t \in \mathbb{R}} S_t$. In any case, there exist distinct two points $\tilde{v}_+$ and $\tilde{v}_-$ such that $\tilde{v}_\pm \in S_{\pm\theta_\pm} \cap C$. We put $v_\pm = p(\tilde{v}_\pm)$.

to be analytically conjugate to an irrational rotation of a round annulus, where m is the period of A_j. Hence, L_{A_j} depends only on the cycle of Herman rings to which A_j belongs.

Remark 3. A pair v_1, v_2 of critical values can accidentally be very close to each other in Ω, i.e., $d_\Omega(v_1, v_2)$ is very small, while $a_j(v_1, v_2)$ is not so small. (The phenomenon $\mathcal{T}(v) \neq \emptyset$ can be considered as a limiting case of the above situation.) So, the formulation in the above corollary does not always provide a good estimate for the uniform perfectness.

3. Branched coverings and uniform perfectness

In this section, we shall investigate the connection between branched coverings and uniform perfectness. Let $f : U \to W$ be a holomorphic (possibly branched) covering map from a (connected) hyperbolic Riemann surface U onto another W. Precisely speaking, for each point $w \in W$ there exists an open neighborhood V of w satisfying the condtion: For each component $\tilde{V}$ of $f^{-1}(V)$ there exist a natural number $n \geq 1$ and conformal homeomorphisms $\varphi : \tilde{V} \to \Delta_r$ and $\psi : V \to \Delta_{r^n}$ with $\psi(w) = 0$ such that $\psi \circ f \circ \varphi^{-1}(\zeta) = \zeta^n$, where Δ_r denotes the disk $\{|\zeta| < r\}$.

When a loop α is freely homotopic to another α' in U, $f_*\alpha := f \circ \alpha$ is freely homotopic to $f_*\alpha'$. Therefore, the natural homomorphism $f_* : \mathcal{C}_U \to \mathcal{C}_W$ is induced by f so that $[\alpha]$ sends to $[f_*\alpha]$.

First, suppose that f is unbranched, then by the homotopy lifting property we can see that the induced map f_* is injective. And moreover $\ell_U[\alpha] = \ell_W[f_*\alpha]$ because f is a local isometry, therefore we have the next

Proposition 3.1. *If $f : U \to W$ is an unbranched holomorphic covering map, then it is valid that $L_U \geq L_W$.*

In the case when f is branched, we need more efforts to estimate L_U from below. In fact, for any finitely connected planar Jordan domain U, it is known that there exists a branched holomorphic covering map from U onto the unit disk (so-called the Ahlfors map), thus L_U cannot be estimated by only the data of W (in this case, $L_W = +\infty$).

Let $\mathrm{Crit}(f)$ be the set of critical points of f and for $v_1, v_2, v \in f(\mathrm{Crit}(f))$ with $v_1 \neq v_2$, define the curve families $\mathcal{S}(v_1, v_2)$ and $\mathcal{T}(v)$ by the same way as in the previous section. And we set

$$a(v_1, v_2) = \inf_{\beta \in \mathcal{S}(v_1, v_2)} \ell_W(\beta), \quad b(v) = \inf_{\beta \in \mathcal{T}(v)} \ell_W(\beta), \text{ and}$$

$$a = \inf_{v_1 \neq v_2 \in f(\mathrm{Crit}(f))} a(v_1, v_2), \quad b = \inf_{v \in f(\mathrm{Crit}(f))} b(v).$$

Then, the following lemma is a key step to our proof of the Main Theorem.

(b') β passes through v essentially two times, at least.

(c) There exists a non-trivial loop α in U_j such that $f_*(\alpha) = \beta$.

More precisely, the condition (b') says that there exist distinct points ζ_0 and ζ_1 in S^1 with $\beta(\zeta_0) = \beta(\zeta_1) = v$ such that the restrictions $\beta|_{\overline{I_1}}$ and $\beta|_{\overline{I_2}}$ of the loop β are both non-trivial closed curves in W_j, where I_1 and I_2 are the connected component of $\mathrm{S}^1 \setminus \{\zeta_0, \zeta_1\}$. And we set

$$a_j(v_1, v_2) = \inf_{\beta \in \mathcal{S}(v_1, v_2)} \ell_\Omega(\beta), \quad b_j(v) = \inf_{\beta \in \mathcal{T}(v)} \ell_\Omega(\beta) \text{ and}$$

$$a_j = \min_{v_1, v_2 \in f(C_j), v_1 \neq v_2} a_j(v_1, v_2), \quad b_j = \min_{v \in f(C_j)} b_j(v),$$

where we set $a_j = +\infty$ if $\#f(C_j) = 1$.

Finally, let $A_1, \cdots, A_t$ be the complete system of representatives of the cycles of Herman rings of f. We note here that, by Shishikura's theorem, $0 \leq t \leq d-2$, in particular, if $d = 2$ there are no Herman rings. And, since the Julia set has no isolated points, the Herman rings have finite moduli, so $L_{A_k} > 0$ for all k.

Now we are ready to state our main theorem.

Theorem 2.2 (Main Thoerem). *For an arbitrary rational map $f : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ of degree $d \geq 2$, the following holds.*

$$L_{\Omega_f} \geq \min\{a_1, \cdots, a_s, b_1, \cdots, b_s, L_{A_1}, \cdots, L_{A_t}\}.$$

The proof of this theorem will be given in Section 4.

For any $\beta \in \mathcal{S}(v_1, v_2)$, it is clear by definition that $\ell_\Omega(\beta) \geq 2d_\Omega(v_1, v_2)$. Similarly, for $\beta \in \mathcal{T}(v)$, we have $\ell_\Omega(\beta) \geq 4\iota_\Omega(v)$. Thus, we conclude that $a_j(v_1, v_2) \geq 2d_\Omega(v_1, v_2)$ and $b_j(v) \geq 4\iota_\Omega(v)$ and hence have the following

Corollary 2.3. *Under the same situation as the Main Theorem, it follows that*

$$L_{\Omega_f} \geq \min\{C_1, C_2, C_3\} (> 0),$$

where $C_1 = \min_{v_1 \neq v_2 \in f(\mathrm{Crit}(f))} 2d_{\Omega_f}(v_1, v_2)$, $C_2 = \min_{v \in f(\mathrm{Crit}(f))} 4\iota_{\Omega_f}(v)$, and $C_3 = \min_{k=1,\cdots,t} L_{A_k}$. In particular, the Julia set J_f is uniformly perfect.

Remark 1. As is well-known, any polynomial has no Herman rings. If there exsits a Herman ring A, it is known that the boundary of A is contained in the closure of forward orbits of the critical set of f. Therefore, if each critical point of f is periodic or contained in a (super)attracting or parabolic basin, then we can conclude that f has no Herman rings. We also note that a cycle of (super)attracting or parabolic components always contains a critical point, thus a component of it appears as a member of the list $U_1, \cdots, U_s$.

Remark 2. If $A_k = f^l(A_j)$ for some $1 \leq j, k \leq t$ and $1 \leq l$, then A_j and A_k belong to the same cylce of Herman rings, thus $L_{A_j} = L_{A_k}$ since $f^m : A_j \to A_j$ is known

is, $\iota_\Omega(z)$ is the maximal radius r so that the hyperbolic disk $\{w \in \Omega; d_\Omega(z,w) < r\}$ is simply connected.

Let $\mathcal{C}_\Omega$ denote the set of free homotopy classes of non-trivial loops in Ω, where a loop (=closed curve) is called non-trivial if this is not null-homotopic (=contractible) in Ω. For a loop α in Ω, we define the length of it by

$$\ell_\Omega(\alpha) = \int_\alpha \rho_\Omega(z)|dz|,$$

and for the free homotopy class $[\alpha]$ represented by α, we define

$$\ell_\Omega[\alpha] = \inf_{\alpha' \in [\alpha]} \ell_\Omega(\alpha').$$

Finally, we set

$$L_\Omega = \inf_{[\alpha] \in \mathcal{C}_\Omega} \ell_\Omega[\alpha].$$

(If $\mathcal{C}_\Omega$ is an empty set, we set $L_\Omega = +\infty$.) We remark that the injectivity radius $\iota_\Omega(z)$ is equal to the half of the infimum of lengths of non-trivial loops in Ω passing through z. In particular, L_Ω is nothing other than twice the (global) injectivity radius $\inf_{z\in\Omega} \iota_\Omega(z)$ of Ω.

Concerning the constant L_Ω, the following estimate is fundamental.

Proposition 2.1 ([12])**.**

$$L_\Omega \le \frac{\pi^2}{M_\Omega} \le \min\{L_\Omega e^{L_\Omega}, \frac{L_\Omega^2}{2}\coth^2(L_\Omega/2)\}.$$

In particular, $M_\Omega < \infty$ if and only if $L_\Omega > 0$.

By this proposition, we have only to estimate L_Ω from below. Now let us state the main theorem. For basic definitions and results about the complex dyamics of the rational maps, we refer to the textbook [2] by Beardon as a general reference.

Let $f : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ be a rational map of degree $d \ge 2$. We denote by $J = J_f$ and $\Omega = \Omega_f$ the Julia set and the Fatou set of f, respectively. (In other words, Ω_f is the domain of normality of the iteration family $\{f^n\}_{n=1,2,\dots}$ of f and $J_f = \widehat{\mathbb{C}} \setminus \Omega_f$.) Note that Ω_f is completely invariant under f, precisely, $f(\Omega_f) = \Omega_f = f^{-1}(\Omega_f)$.

We denote by $\mathrm{Crit}(f)$ the set of critical points of f in the Fatou set Ω_f and let $U_1, \cdots, U_s$ be the complete list of the components of Ω_f which contains at least one critical point of f and is not simply connected. And we set $W_j = f(U_j)$ and $C_j = \mathrm{Crit}(f) \cap U_j$ for $j = 1, \cdots, s$. Note here that $\#\mathrm{Crit}(f) \le 2d-2$, so $s \le 2d-2$. Now we introduce two kinds of curve family: $\mathcal{S}(v_1, v_2)$ and $\mathcal{T}(v)$, for $v_1, v_2, v \in f(C_j)$ with $v_1 \ne v_2$. Let $\mathcal{S}(v_1, v_2)$ and $\mathcal{T}(v)$ consist of the loops $\beta : \mathrm{S}^1 \to W_j$ satisfying the conditions (a), (b), (c) and (a), (b'), (c), respectively, in the following:

(a) β is contractible in W_j.
(b) β passes through v_1 and v_2.

We will state the main result in the next section as well as fundamental definitions and notations. In Section 3, we shall discuss the connection between branched coverings and the uniform perfectness, which will be a key to prove our main theorem. Section 4 is devoted to prove our main theorem, and we make essential use of Sullivan's No Wandering Domains Theorem. We shall give applications of the main result to estimate the Hausdorff dimension, the (logarithmic) capacity density of the Julia set and the Poincaré metric of the Fatou set in Section 5. In the last section, as a special case, we investigate the quadratic family of polynomials. In particular, we present a concrete estimate for a subclass of this family.

Finally, the author would like to express his hearty gratitude to Professors M. Taniguchi and K. Matsuzaki for helpful comments.

2. Main result

Let C be a closed set in the Riemann sphere containing at least three points and Ω its complement. We denote by $\mathcal{A}_\Omega$ and $\mathcal{A}_\Omega^\circ$ the sets of annuli and round annuli, respectively, in Ω separating C, where annuli mean doubly connected domains and round annuli do special annuli of the form $\{z \in \mathbb{C}; r_1 < |z-a| < r_2\}$ for some $a \in \mathbb{C}$ and $0 \le r_1 < r_2 \le \infty$. The modulus $m(A)$ of an annulus A is the number m such that A is conformally equivalent to the round annulus $\{z; 1 < |z| < e^m\}$. We set

$$M_\Omega = \sup_{A \in \mathcal{A}_\Omega} m(A), \quad M_\Omega^\circ = \sup_{A \in \mathcal{A}_\Omega^\circ} m(A),$$

and call them the modulus and the round modulus of Ω. (If $\mathcal{A}_\Omega$ or $\mathcal{A}_\Omega^\circ$ is empty, then we define $M_\Omega = 0$ or $M_\Omega^\circ = 0$, respectively.) For these constants, it is known that $\frac{1}{2}M_\Omega - 1.7332\cdots \le M_\Omega^\circ \le M_\Omega$, and that if $\Omega \subset \mathbb{C}$, $\quad M_\Omega - 2.8911\cdots \le M_\Omega^\circ \le M_\Omega$ (cf. [12]).

It is easily verified that C is uniformly perfect if and only if $M_\Omega^\circ < \infty$, equivalently $M_\Omega < \infty$.

Next, for later use, we consider quantities determined by the hyperbolic geometry of Ω. Since $\#C \ge 3$, each component D of Ω is hyperbolic, i.e., there exists a holomorphic universal covering map $p : \mathbb{H} \to D$ from the upper half plane onto D. Thus D can be regarded as the quotient space $\mathbb{H}/\Gamma$ of $\mathbb{H}$ over the covering transformation group $\Gamma = \{\gamma \in \mathrm{PSL}(2,\mathbb{R}); p \circ \gamma = p\}$. Since the hyperbolic (Poincaré) metric $\rho_{\mathbb{H}}(z)|dz| = \frac{|dz|}{2\mathrm{Im}z}$ is invariant under the action of $\mathrm{PSL}(2,\mathbb{R})$, $\quad D$ inherits the hyperbolic metric $\rho_D(z)|dz|$ so that $p : \mathbb{H} \to D$ is a local isometry with respect to the hyperbolic metric, i.e., $\rho_{\mathbb{H}} = p^*\rho_D$. Therefore, we can define the hyperbolic metric ρ_Ω of Ω componentwise.

The hyperbolic distance $d_\Omega(z_0, z_1)$ of a pair of points z_0, z_1 in the same component of Ω can be defined by $\inf_\alpha \int_\alpha \rho_\Omega(z)|dz|$, where the infimum is taken over all paths joining z_0 to z_1 in Ω. We also difine $d_\Omega(z_0, z_1) = +\infty$ if z_0 and z_1 do not belong to the same component of Ω. For $z \in \Omega$ we denote by $\iota_\Omega(z)$ the injectivity radius of Ω at z, that

AN EXPLICIT BOUND FOR UNIFORM PERFECTNESS OF THE JULIA SETS OF RATIONAL MAPS

TOSHIYUKI SUGAWA

August 4, 1997

Abstract. A compact set C in the Riemann sphere is called uniformly perfect if the moduli of annuli separating C are bounded. Mañé-da Rocha and Hinkkanen showed independently the uniform perfectness of the Julia sets of rational maps of degree ≥ 2, but they presented no explicit bounds for uniform perfectness. In this note, we shall provide such an explicit bound and, as a result, we give another proof of uniform perfectness of the Julia sets. As an application, we refer to a lower estimate of the Hausdorff dimension of the Julia sets.

1. Introduction

Let C be a closed set in the Riemann sphere $\widehat{\mathbb{C}}$ and Ω its complement. We say C is *uniformly perfect* if there exists a constant $0 < c < 1$ such that $C \cap \{z \in \mathbb{C}; cr < |z - a| < r\} \neq \emptyset$ for any $a \in C$ and $0 < r < \mathrm{diam}(C)$, where diam denotes the Euclidean diameter.

The notion of uniform perfectness first appeared in Beardon-Pommerenke [3], and was investigated more deeply by Pommerenke [9] and [10], and afterwards by many authors (see [12] and its reference). By definition, the sets with self-similarities in some sense are expected to have uniform perfectness. In fact, the limit sets are known to be uniformly perfect for a wide class of Kleinian groups (cf. Sugawa [11]). On the other hand, Pommerenke [10] first showed the uniform perfectness of the Julia sets of hyperbolic rational maps. Later, Mañé-da Rocha [8] and Hinkkanen [6] proved in the case of general rational maps of degree ≥ 2, independently. For a simpler proof, see the textbook [4] by Carleson and Gamelin. But, their proofs are done by contradiction, thus no explicit bounds for uniform perfectness are given. In this note, we shall present such an explicit bound and also exhibit some applications of this result. Our proof emploies the hyperbolic geometry, and hence is different from ones of the above authors. As Hinkkanen remarked in [6], we should note that there exists an entire function whose limit set is not uniformly perfect.

1991 *Mathematics Subject Classification.* Primary 30D05, Secondary 30F45.

Key words and phrases. uniformly perfect, Julia set.