RELATIVE BOGOMOLOV'S INEQUALITY

ATSUSHI MORIWAKI

ABSTRACT. Let $f : X \to Y$ be a projective morphism of smooth quasi-projective varieties over an algebraically closed field of characteristic zero such that the generic fiber of f is a geometrically irreducible smooth curve. Let E be a vector bundle of rank r on X. In this paper, we would like to show that if X_y is smooth and E_y is semistable for $y \in Y$, then $f_*\left(2rc_2(E) - (r-1)c_1(E)^2\right)$ is weakly positive at y. We apply this result to a cone of weakly positive $\mathbb{Q}$-Cartier divisors on the moduli space of stable curves, namely, we have the following. Let $\overline{\mathcal{M}}_g$ (resp. $\mathcal{M}_g$) be the moduli space of stable (resp. smooth) curves of genus $g \geq 2$. Let λ be the Hodge class and δ_i's ($i = 0, \ldots, [g/2]$) boundary classes. Then, a $\mathbb{Q}$-Cartier divisor $x\lambda + \sum_{i=0}^{[g/2]} y_i\delta_i$ on $\overline{\mathcal{M}}_g$ is weakly positive over $\mathcal{M}_g$ if and only if $x \geq 0$, $gx + (8g+4)y_0 \geq 0$, and $i(g-i)x + (2g+1)y_i \geq 0$ for all $1 \leq i \leq [g/2]$.

CONTENTS

Introduction 1
1. Pseudo-effectivity, semi-ampleness and weak positivity of divisors 4
2. Strong semistability of vector bundles 8
3. Proof of relative Bogomolov's inequality 10
4. A weakly positive divisor on the moduli space of stable curves 15
5. Cones of positive divisors on the moduli space of stable curves 16
6. Effective Bogomolov's conjecture over function fields 19
7. Generalization to higher dimensional fibrations 24
Appendix A. A construction of a certain fibration of hyperelliptic curves 25
References 31

INTRODUCTION

Throughout this paper, we fix an algebraically closed field k.

Let $f : X \to Y$ be a projective morphism of quasi-projective varieties over k. We assume that the generic fiber of f is a geometrically irreducible smooth curve. Let E be a vector bundle of rank r on X. Then, $(2rc_2(E) - (r-1)c_1(E)^2) \cap [X]$ is a cycle of dimension $\dim Y - 1$ on X. Thus, we get a divisor $f_*\left((2rc_2(E) - (r-1)c_1(E)^2) \cap [X]\right)$ on Y. We denote this divisor by $\mathrm{dis}_{X/Y}(E)$ and it is called a *discriminant divisor* of E with respect to $f : X \to Y$. In this paper, we would like to show the following theorem (cf. Corollary 3.5) and give its applications.

Date: 25/May/1997, (Version 4.0), LAST MODIFIED ON 08/April/1997.
1991 *Mathematics Subject Classification.* Primary 14H10, 14C20; Secondary 14G40.

Theorem A ($\mathrm{char}(k) \geq 0$)**.** *We assume that Y is smooth over k. Let y be a point of Y and $\overline{\kappa(y)}$ the algebraic closure of the residue field $\kappa(y)$ at y. If f is smooth over y and E is strongly semistable on the geometric fiber $X_{\bar{y}} = X \times_Y \mathrm{Spec}\left(\overline{\kappa(y)}\right)$ over y, then $\mathrm{dis}_{X/Y}(E)$ is weakly positive at y, namely, for any ample divisors A on Y and any positive integers n, there is a positive integer m such that*

$$H^0(Y, \mathcal{O}_Y(m(n\,\mathrm{dis}_{X/Y}(E) + A))) \otimes \mathcal{O}_Y \to \mathcal{O}_Y(m(n\,\mathrm{dis}_{X/Y}(E) + A))$$

is surjective at y. (For the definition of strong semistability, see §2. If the characteristic of k is zero, the strong semistability is nothing more than the usual semistability of vector bundles.)

An interesting point of the above theorem is that even if the weak positivity of $\mathrm{dis}_{X/Y}(E)$ at y is a global property on Y, it can be derived from the local assumption "smoothness of $X_{\bar{y}}$ and strong semistability of $E_{\bar{y}}$". This gives a great advantage to our applications.

An immediate application of our inequality is a problem concerning the positivity of divisors on the moduli space of stable curves. Let $g \geq 2$ be an integer, and $\overline{\mathcal{M}}_g$ (resp. $\mathcal{M}_g$) the moduli space of stable (resp. smooth) curves of genus g over k. The boundary $\overline{\mathcal{M}}_g \setminus \mathcal{M}_g$ is of codimension one and has $[g/2]+1$ irreducible components, say, $\Delta_0, \Delta_1, \ldots, \Delta_{[g/2]}$. The geometrical meaning of indexes is as follows. A general point of Δ_0 represents an irreducible stable curve with one node, and a general point of Δ_i ($i > 0$) represents a stable curve consisting of a curve of genus i and a curve of genus $g-i$ joined at one point. Let δ_i be the class of Δ_i in $\mathrm{Pic}(\overline{\mathcal{M}}_g) \otimes \mathbb{Q}$ (exactly speaking, $\delta_i = c_1(\mathcal{O}(\Delta_i))$ for $i \neq 1$, and $\delta_1 = \frac{1}{2}c_1(\mathcal{O}(\Delta_1))$), and λ the Hodge class on $\overline{\mathcal{M}}_g$. The fundamental question due to Mumford is whether a $\mathbb{Q}$-divisor

$$a\lambda - b_0\delta_0 - b_1\delta_1 - \cdots - b_{[g/2]}\delta_{[g/2]}$$

has a kind of positivity for some rational numbers $a, b_0, \cdots, b_{[g/2]}$. For this problem, we can use a lot of types of positivity, namely, ampleness, numerical effectivity, effectivity, pseudo-effectivity, and so on. Besides them, we would like to introduce a new sort of positivity for our purposes. Let V be a projective variety over k and U a non-empty Zariski open set of V. A $\mathbb{Q}$-Cartier divisor D on V is said to be *numerically effective over U* if $(D \cdot C) \geq 0$ for all irreducible curves C on V with $C \cap U \neq \emptyset$. A first general result in this direction was found by Cornalba-Harris [2], Xiao [18] and Bost [1]. They proved that a $\mathbb{Q}$-divisor

$$(8g+4)\lambda - g(\delta_0 + \delta_1 + \cdots + \delta_{[g/2]})$$

is numerically effective over $\mathcal{M}_g$. As we observed in [14] and [15], it is not sharp in coefficients of δ_i ($i > 0$). Actually, the existence of a certain refinement of the above result was predicted at the end of the paper [2]. Our answer for this problem is the following (cf. Theorem 4.2 and Proposition 1.9).

Theorem B ($\mathrm{char}(k) = 0$)**.** *A divisor*

$$(8g+4)\lambda - g\delta_0 - \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i$$

is weakly positive over $\mathcal{M}_g$, i.e., if we denote $(8g+4)\lambda - g\delta_0 - \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i$ by D, then for any ample $\mathbb{Q}$-Cartier divisors A on $\overline{\mathcal{M}}_g$, there is a positive integer n such that $n(D+A)$ is a Cartier divisor and

$$H^0(\overline{\mathcal{M}}_g, \mathcal{O}_{\overline{\mathcal{M}}_g}(n(D+A))) \otimes \mathcal{O}_{\overline{\mathcal{M}}_g} \to \mathcal{O}_{\overline{\mathcal{M}}_g}(n(D+A))$$

is surjective on $\mathcal{M}_g$. In particular, it is pseudo-effective, and numerically effective over $\mathcal{M}_g$.

As an application of this theorem, we can decide a cone of weakly positive divisors over $\mathcal{M}_g$ (cf. Corollary 5.4).

Theorem C ($\mathrm{char}(k) = 0$)**.** *If we denote by $\mathrm{WP}(\overline{\mathcal{M}}_g; \mathcal{M}_g)$ the cone in $\mathrm{Pic}(\overline{\mathcal{M}}_g) \otimes \mathbb{Q}$ consisting of weakly positive $\mathbb{Q}$-Cartier divisors over $\mathcal{M}_g$, then*

$$\mathrm{WP}(\overline{\mathcal{M}}_g; \mathcal{M}_g) = \left\{ x\lambda + \sum_{i=0}^{[g/2]} y_i\delta_i \;\middle|\; \begin{array}{l} x \geq 0, \\ gx + (8g+4)y_0 \geq 0, \\ i(g-i)x + (2g+1)y_i \geq 0 \quad (1 \leq i \leq [g/2]). \end{array} \right\}.$$

Moreover, using Theorem B, we can deduce a certain kind of inequality on an algebraic surface. In order to give an exact statement, we will introduce types of nodes of semistable curves. Let Z be a semistable curve over k, and P a node of Z. We can assign a number i to the node P in the following way. Let $\iota_P : Z_P \to Z$ be the partial normalization of Z at P. If Z_P is connected, then $i = 0$. Otherwise, i is the minimum of arithmetic genera of two connected components of Z_P. We say the node P of Z is *of type i*.

Let X be a smooth projective surface over k, Y a smooth projective curve over k, and $f : X \to Y$ a semistable curve over Y of genus $g \geq 2$. By abuse of notation, we denote by $\delta_i(X/Y)$ the number of nodes of type i in all singular fibers of f. Actually, $\delta_i(X/Y) = \deg(\pi^*(\delta_i))$. where $\pi : Y \to \overline{\mathcal{M}}_g$ is a morphism induced by $f : X \to Y$. Then, we have the following (cf. Corollary 4.3).

Theorem D ($\mathrm{char}(k) = 0$)**.** *With notation being as above, we have an inequality*

$$(8g+4)\deg(f_*(\omega_{X/Y})) \geq g\delta_0(X/Y) + \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i(X/Y).$$

As an arithmetic application of Theorem D, we can show the following answer for the effective Bogomolov's conjecture over function fields (cf. Theorem 6.2). (Recently, the Bogomolov's conjecture over number fields was solved by Ullmo [17], but the effective Bogomolov's conjecture is still open.)

Theorem E ($\mathrm{char}(k) = 0$)**.** *We assume that f is not smooth and every singular fiber of f is a tree of stable components, i.e., every node of type 0 on the stable model of each singular fiber is a singularity of an irreducible component, then the effective Bogomolov's conjecture holds for the generic fiber of f, namely, we have the following. Let K be the function field of Y, C the generic fiber of f, $\mathrm{Jac}(C)$ the Jacobian of C, and let $j : C(\overline{K}) \to \mathrm{Jac}(C)(\overline{K})$ be a morphism given by $j(P) = \omega_C - P$. Then, $j(C(\overline{K}))$ is discrete in terms of the seminorm arising from the Neron-Tate height paring on $\mathrm{Jac}(C)(\overline{K})$. More precisely, the maximal radius of ball in which we have only finitely many points coming from $C(\overline{K})$ via j is greater*

than or equal to

$$\sqrt{\frac{(g-1)^2}{g(2g+1)}\left(\frac{g-1}{3}\delta_0(X/Y)+\sum_{i=1}^{\left[\frac{g}{2}\right]}4i(g-i)\delta_i(X/Y)\right)}.$$

Finally, we would like to express our hearty thanks to Institut des Hautes Études Scientifiques where all works of this paper had been done, and to Prof. Bost who pointed out a fatal error of the previous version of it.

1. Pseudo-effectivity, semi-ampleness and weak positivity of divisors

In this section, we will introduce several concepts of positivity and investigate their elementary properties.

Let X be an n-dimensional algebraic variety over k. Let $Z_{n-1}(X)$ be a free abelian group generated by subvarieties of dimension $n-1$, and $\mathrm{Div}(X)$ a subgroup of $Z_{n-1}(X)$ consisting of Cartier divisors on X. We denote $Z_{n-1}(X)$ (resp. $\mathrm{Div}(X)$) modulo the linear equivalence by $A_{n-1}(X)$ (resp. $\mathrm{Pic}(X)$). An element of $Z_{n-1}(X)\otimes\mathbb{Q}$ (resp. $\mathrm{Div}(X)\otimes\mathbb{Q}$) is called a $\mathbb{Q}$-*divisor* (resp. $\mathbb{Q}$-*Cartier divisor*) on X.

We say a $\mathbb{Q}$-divisor D is the limit of a sequence $\{D_m\}_{m=1}^{\infty}$ of $\mathbb{Q}$-divisors, denoted by $D=\lim_{m\to\infty}D_m$, if there are $\mathbb{Q}$-divisors $Z_1,\ldots,Z_l$ and infinite sequences $\{a_{1,m}\}_{m=1}^{\infty},\ldots,\{a_{l,m}\}_{m=1}^{\infty}$ of rational numbers such that $D\sim D_m+\sum_{i=1}^{l}a_{i,m}Z_i$ for all $m\geq 1$ and $\lim_{m\to\infty}a_{i,m}=0$ for all $i=1,\ldots,l$. Similarly, replacing $\mathbb{Q}$-divisor by $\mathbb{Q}$-Cartier divisor in the above sentence, we can define the limit of $\mathbb{Q}$-Cartier divisors. A $\mathbb{Q}$-divisor D is said to be *pseudo-effective* if it is the limit of effective $\mathbb{Q}$-divisors.

Let x be a point of X. A $\mathbb{Q}$-Cartier divisor D on X is said to be *semi-ample at* x if there is a positive integer n such that $nD\in\mathrm{Div}(X)$ and $H^0(X,\mathcal{O}_X(nD))\otimes\mathcal{O}_X\to\mathcal{O}_X(nD)$ is surjective at x. Further, D is said to be *weakly positive at* x if there is an infinite sequence $\{D_m\}_{m=1}^{\infty}$ of $\mathbb{Q}$-Cartier divisors on X such that D_m is semi-ample at x for all $m\geq 1$ and $D=\lim_{m\to\infty}D_m$. Moreover, D is said to be *numerically effective at* x if $(D\cdot C)\geq 0$ for any complete irreducible curves C passing through x. First, let us consider relationship of each positivity.

Proposition 1.1 ($\mathrm{char}(k)\geq 0$)**.** *Let D be a $\mathbb{Q}$-Cartier divisor on X and x a point of X. Then, we have the following.*

(1) *If D is weakly positive at x, then D is pseudo-effective.*
(2) *If D is weakly positive at x, then D is numerically effective at x.*

$$\text{“semi-ample at } x\text{”} \implies \text{“weakly positive at } x\text{”} \implies \text{“numerically effective at } x\text{”}$$
$$\Downarrow$$
$$\text{“pseudo-effective”}$$

Proof. (1) This is obvious because a semi-ample $\mathbb{Q}$-Cartier divisor at x is linearly equivalent to an effective $\mathbb{Q}$-Cartier divisor.

(2) Since D is the limit of semi-ample $\mathbb{Q}$-divisors at x, it is sufficient to show that if D is semi-ample at x, then D is numerically effective at x. Let C be a complete irreducible curve

passing through x. Since D is semi-ample at x, there are a positive integer n and an effective Cartier divisor E such that $nD \sim E$ and $x \notin E$. Thus, $C \not\subseteq E$. Hence $(E \cdot C) = n(D \cdot C) \geq 0$. □

In order to consider further properties, we need to prepare two lemmas.

Lemma 1.2 ($\mathrm{char}(k) \geq 0$)**.** *Let $\pi : X \to Y$ be a proper morphism of quasi-projective varieties over k and y a point of Y such that π is finite over y. Let F be a coherent $\mathcal{O}_X$-module and H an ample line bundle on Y. Then there is a positive integer n_0 such that, for all $n \geq n_0$,*

$$H^0(X, F \otimes \pi^*(H^{\otimes n})) \otimes \mathcal{O}_X \to F \otimes \pi^*(H^{\otimes n})$$

is surjective at each point of $\pi^{-1}(y)$.

Proof. Let n_0 be a positive integer such that, for all $n \geq n_0$, $\pi_*(F) \otimes H^{\otimes n}$ is generated by global sections, i.e.,

$$H^0(Y, \pi_*(F) \otimes H^{\otimes n}) \otimes \mathcal{O}_Y \to \pi_*(F) \otimes H^{\otimes n}$$

is surjective. Thus,

$$H^0(X, F \otimes \pi^*(H^{\otimes n})) \otimes \mathcal{O}_X \to \pi^*\pi_*(F \otimes \pi^*(H^{\otimes n}))$$

is surjective because $\pi_*(F \otimes \pi^*(H^{\otimes n})) = \pi_*(F) \otimes H^{\otimes n}$. On the other hand, since π is finite over y,

$$\pi^*\pi_*(F \otimes \pi^*(H^{\otimes n})) \to F \otimes \pi^*(H^{\otimes n})$$

is surjective at each point of $\pi^{-1}(y)$. Thus, we get our assertion. □

Lemma 1.3 ($\mathrm{char}(k) \geq 0$)**.** *Let $\pi : X \to Y$ be a proper morphism of quasi-projective varieties over k and x a point of X such that π is finite over $\pi(x)$. Let D a $\mathbb{Q}$-Cartier divisor on X and A a $\mathbb{Q}$-Cartier divisor on Y. If D is weakly positive at x and A is ample, then $D + \pi^*(A)$ is semi-ample at x.*

Proof. By our assumption, there are $\mathbb{Q}$-Cartier divisors $Z_1, \ldots, Z_l$, an infinite sequence $\{D_m\}_{m=1}^{\infty}$ of $\mathbb{Q}$-Cartier divisors, and infinite sequences $\{a_{1,m}\}_{m=1}^{\infty}, \ldots, \{a_{l,m}\}_{m=1}^{\infty}$ of rational numbers with the following properties.

(i) $D \sim D_m + \sum_{i=1}^{l} a_{i,m} Z_i$ for sufficiently large m.
(ii) $\lim_{m \to \infty} a_{i,m} = 0$ for all $i = 1, \ldots, l$.
(iii) D_m is semi-ample at x for $m \gg 0$.

By virtue of Lemma 1.2, we can find a positive integer n such that $n\pi^*(A) + Z_i$ and $n\pi^*(A) - Z_i$ are semi-ample at x. We choose a sufficiently large m with $|nla_{i,m}| < 1$ $(i = 1, \ldots, l)$. Then,

$$\frac{1 + nla_{i,m}}{2ln} > 0 \quad \text{and} \quad \frac{1 - nla_{i,m}}{2ln} > 0.$$

On the other hand,

$$D + \pi^*(A) \sim D_m + \sum_{i=1}^{l} \left(\frac{1 + nla_{i,m}}{2ln}(n\pi^*(A) + Z_i) + \frac{1 - nla_{i,m}}{2ln}(n\pi^*(A) - Z_i) \right).$$

Thus, $D + \pi^*(A)$ is semi-ample at x. □

Here we have the following characterization of weak positivity in terms of ample divisors.

Proposition 1.4 ($\mathrm{char}(k) \geq 0$)**.** *Let X be a quasi-projective variety over k, x a point of X, and D a $\mathbb{Q}$-Cartier divisor on X. Then, the following are equivalent.*

(1) *D is weakly positive at x.*
(2) *For any ample $\mathbb{Q}$-Cartier divisors A on X, $D + A$ is semi-ample at x.*

Proof. First we assume (1). Then, by Lemma 1.3, $D + A$ is semi-ample at x.

Next we assume (2). Fix an ample Cartier divisor A on X. We set $E_m = D + (1/m)A$. Then, $D = E_m - (1/m)A$ and E_m is semi-ample at x. Thus, D is weakly positive at x. □

Using the above characterization, we can see that the weak positivity is closed under taking a limit.

Proposition 1.5 ($\mathrm{char}(k) \geq 0$)**.** *Let X be a quasi-projective variety over k, x a point of X, and D a $\mathbb{Q}$-Cartier divisor on X. If D is the limit of weakly positive $\mathbb{Q}$-Cartier divisors at x, then D is weakly positive at x.*

Proof. By our assumption, $D = \lim_{m\to\infty} D_m$, where D_m is a weakly positive $\mathbb{Q}$-Cartier divisors at x. Let A be an ample $\mathbb{Q}$-Cartier divisor. Then, $D = \lim_{m\to\infty} (D_m + (1/m)A)$. By Lemma 1.3, $D_m + (1/m)A$ is semi-ample at x. Thus, D is weakly positive at x. □

Next, let us consider functorial properties of semi-ampleness and weak positivity under pull-back and push-forward.

Proposition 1.6 ($\mathrm{char}(k) \geq 0$)**.** *Let $\pi : X \to Y$ be a morphism of quasi-projective varieties. Let D be a $\mathbb{Q}$-Cartier divisor on Y and x a point of X. If $\pi^*(D)$ is defined, then we have the following. (Note that even if $\pi^*(D)$ is not defined, there is a $\mathbb{Q}$-Cartier divisor D' such that $D' \sim D$ and $\pi^*(D')$ is defined.)*

(1) *If D is semi-ample at $\pi(x)$, then $\pi^*(D)$ is semi-ample at x.*
(2) *If D is weakly positive at $\pi(x)$, then $\pi^*(D)$ is weakly positive at x.*

Proof. (1) By our assumption, $H^0(Y, \mathcal{O}_Y(nD)) \otimes \mathcal{O}_Y \to \mathcal{O}_Y(nD)$ is surjective at $\pi(x)$ for a sufficiently large n. Thus, $H^0(Y, \mathcal{O}_Y(nD)) \otimes \mathcal{O}_X \to \mathcal{O}_X(n\pi^*(D))$ is surjective at x. Here let us consider following commutative diagram:

$$\begin{array}{ccc} H^0(Y, \mathcal{O}_Y(nD)) \otimes \mathcal{O}_X & \xrightarrow{\alpha} & \mathcal{O}_X(n\pi^*(D)) \\ \downarrow & & \| \\ H^0(X, \mathcal{O}_X(n\pi^*(D))) \otimes \mathcal{O}_X & \xrightarrow{\alpha'} & \mathcal{O}_X(n\pi^*(D)) \end{array}$$

Since α is surjective at x, so is α'. Therefore, $\pi^*(D)$ is semi-ample at x.

(2) Let A be a ample divisor on Y such that $\pi^*(A)$ is defined. Then, Lemma 1.3 $D+(1/n)A$ is semi-ample at $\pi(x)$ for all $n > 0$. Thus, by (1), $\pi^*(D) + (1/n)\pi^*(A)$ is semi-ample at x. Therefore, $\pi^*(D)$ is weakly positive at x. □

Proposition 1.7 ($\mathrm{char}(k) \geq 0$)**.** *Let $\pi : X \to Y$ be a proper morphism of quasi-projective varieties over k. Let D be a $\mathbb{Q}$-Cartier divisor on X and y a point of Y such that $\pi_*(D)$ is a $\mathbb{Q}$-Cartier divisor on Y and π is finite over y. We set $\pi^{-1}(y) = \{x_1, \ldots, x_n\}$. Then, we have the following.*

(1) *If D is semi-ample at $x_1, \ldots, x_n$, then $\pi_*(D)$ is semi-ample at y.*
(2) *If D is weakly positive at $x_1, \ldots, x_n$, then $\pi_*(D)$ is weakly positive at y.*

Proof. (1) Clearly, we may assume that D is a Cartier divisor. If we take a sufficient large integer n, then $H^0(X, \mathcal{O}_X(nD)) \otimes \mathcal{O}_X \to \mathcal{O}_X(nD)$ is surjective at $x_1, \ldots, x_n$. Thus, there are sections $s_1, \ldots, s_n$ of $H^0(X, \mathcal{O}_X(nD))$ with $s_i(x_i) \neq 0$ for all $i = 1, \ldots, n$. For $\alpha = (\alpha_1, \ldots \alpha_n) \in k^n$, we set $s_\alpha = \alpha_1 s_1 + \cdots + \alpha_n s_n$. Further, we set $V_i = \{\alpha \in k^n \mid s_\alpha(x_i) = 0\}$. Then, $\dim V_i = n-1$ for all i. Thus, since $\#(k) = \infty$, there is $\alpha \in k^n$ with $\alpha \notin V_1 \cup \cdots \cup V_r$, i.e., $s_\alpha(x_i) \neq 0$ for all i. Let us consider a divisor $E = \mathrm{div}(s_\alpha)$. Then, $E \sim nD$. Thus, $\pi_*(E) \sim n\pi_*(D)$. Here, $x_i \notin E$ for all i. Hence, $y \notin \pi_*(E)$. Therefore, we get our assertion.

(2) Let A be an ample divisor on Y. We set $D_n = D + (1/n)\pi^*(A)$. Then, by Lemma 1.3, D_n is semi-ample at $x_1, \ldots, x_n$. Thus, by (1), $\pi_*(D_n) = \pi_*(D) + (1/n)\deg(\pi)A$ is semi-ample at y. Therefore, $\pi_*(D)$ is weakly positive at y. □

One advantage to use semi-ampleness or weak positivity is that we can avoid bad subvarieties of codimension two.

Proposition 1.8 ($\mathrm{char}(k) \geq 0$)**.** *Let X be a normal quasi-projective variety over k, X_0 a Zariski open set of X, and x a point of X_0. Let D be a $\mathbb{Q}$-Cartier divisor on X and $D_0 = D|_{X_0}$. If $\mathrm{codim}(X \setminus X_0) \geq 2$, then we have the following.*
(1) *D is semi-ample at x if and only if D_0 is semi-ample at x.*
(2) *D is weakly positive at x if and only if D_0 is weakly positive at x.*

Proof. (1) This is obvious because $H^0(X_0, \mathcal{O}_{X_0}(nD_0)) \xrightarrow{\sim} H^0(X, \mathcal{O}_X(nD))$ for all integer n with $nD \in \mathrm{Div}(X)$.

(2) Let A be an ample divisor on X and $A_0 = A|_{X_0}$. Then, by virtue of Proposition 1.4, D is weakly positive at x if and only if $D + \epsilon A$ is semi-ample at x for all positive rational number ϵ. In the same way, D_0 is weakly positive at x if and only if $D_0 + \epsilon A_0$ is semi-ample at x for all positive rational number ϵ. Thus, (1) implies (2). □

Finally, let us consider semi-ampleness and weak positivity over an open set. Let X be a quasi-projective variety over k, U a Zariski open set of X, and D a $\mathbb{Q}$-Cartier divisor on X. We say D is *semi-ample over U* (resp. *weakly positive over U*) if D is semi-ample (resp. weakly positive) at all points of U.

Proposition 1.9 ($\mathrm{char}(k) \geq 0$)**.** *With the notation being as above, we have the following.*
(1) *If D is semi-ample over U, then there is a positive integer n such that nD is a Cartier divisor and $H^0(X, \mathcal{O}_X(nD)) \otimes \mathcal{O}_X \to \mathcal{O}_X(nD)$ is surjective on U.*
(2) *If D is weakly positive over U, then, for any ample $\mathbb{Q}$-Cartier divisors A on X, there is a positive integer n such that $n(D+A)$ is a Cartier divisor and*

$$H^0(X, \mathcal{O}_X(n(D+A))) \otimes \mathcal{O}_X \to \mathcal{O}_X(n(D+A))$$

is surjective on U.

Proof. (1) Clearly we may assume that D is a Cartier divisor. For each $x \in U$, there is a positive integer n_x such that $H^0(X, \mathcal{O}_X(n_x D)) \otimes \mathcal{O}_X \to \mathcal{O}_X(n_x D)$ is surjective at x. Thus, we can find a Zariski open set U_x such that $x \in U_x$, $U_x \subseteq U$ and $H^0(X, \mathcal{O}_X(n_x D)) \otimes \mathcal{O}_X \to$

$\mathcal{O}_X(n_xD)$ is surjective on U_x. Here $U = \bigcup_{x\in U} U_x$ and U is a noetherian scheme. Thus, there is $x_1, \ldots, x_l \in U$ with $U = U_{x_1} \cup \cdots \cup U_{x_l}$. Hence, if we set $n = n_{x_1} \cdots n_{x_l}$, then we have our assertion.

(2) This is a consequence of (1). □

2. Strong semistability of vector bundles

In this section, we will consider semistability of tensor products of semistable vector bundles, which was studied by a lot of authors [6], [7], [11], [10] and etc. This however does not hold in positive characteristic [5], so that we will introduce strong semistability of vector bundles.

Let C be a smooth projective curve over k. For a vector bundle F on C, we set $\mu(F) = \deg(F)/\operatorname{rk}(F)$, which is called the *slope of* F. A vector bundle E on C is said to be *semistable* (resp. *stable*) if, for any proper subbundles F of E, $\mu(F) \leq \mu(E)$ (resp. $\mu(F) < \mu(E)$). Moreover, E is said to be *strongly semistable* if, for any finite morphisms $\pi : C' \to C$ of smooth projective curves over k, $\pi^*(E)$ is semistable. Then, we have the following elementary properties of semistable or strongly semistable vector bundles.

Proposition 2.1 ($\operatorname{char}(k) \geq 0$)**.** *Let E be a vector bundle of rank r on C.*

(1) *Let $\pi : C' \to C$ be a finite morphism of smooth projective curves over k such that $k(C')$ is a separable extension of $k(C)$. If E is semistable, then so is $\pi^*(E)$.*
(2) *Under the assumption of $\operatorname{char}(k) = 0$, E is semistable if and only if E is strongly semistable.*
(3) *Let $f : P = \operatorname{Proj}(\bigoplus_{m=0}^{\infty} \operatorname{Sym}^m(E)) \to C$ be the projective bundle of E and $\mathcal{O}_P(1)$ the tautological line bundle on P. Then, E is strongly semistable if and only if $\omega_{P/C}^{\otimes -1} = \mathcal{O}_P(r) \otimes f^*(\det E)^{\otimes -1}$ is numerically effective.*

Proof. (1) is nothing more that [6, Lemma 1.1] and (2) is a consequence of (1).

(3) First we assume that E is strongly semistable. Let Z be any irreducible curve on P. If Z is contained in a fiber, then obviously $(\omega_{P/C}^{\otimes -1} \cdot Z) > 0$. So we may assume that Z is not contained in any fibers. Let C' be the normalization of Z and $\pi : C' \to Z \to C$ the induced morphism. Let $E' = \pi^*(E)$, $f' : P' = \operatorname{Proj}(\bigoplus_{m=0}^{\infty} \operatorname{Sym}^m(E')) \to C'$ the projective bundle of E', and $\mathcal{O}_{P'}(1)$ the tautological line bundle on P'. Then we have the following commutative diagram.

$$\begin{array}{ccc} P & \xleftarrow{\pi'} & P' \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle f'} \\ C & \xleftarrow[\pi]{} & C' \end{array}$$

By our construction, there is a section Z' of f' such that $\pi'(Z') = Z$. We set $Q' = \mathcal{O}_{P'}(1)|_{Z'}$. Then, there is a surjective homomorphism $E' \to Q'$. Since E' is semistable, we have $\mu(E') \leq \deg(Q')$, which means that $(\omega_{P'/C'}^{\otimes -1} \cdot Z') \geq 0$. Here, $\omega_{P'/C'} = \pi'^*(\omega_{P/C})$. Thus, we get $(\omega_{P/C}^{\otimes -1} \cdot Z) \geq 0$.

Conversely, we assume that $\omega_{P/C}^{\otimes -1}$ is numerically effective on P. Let $\pi : C' \to C$ be a finite morphism of smooth projective curves over k. We set $f' : P' \to C'$ and $\pi' : P' \to P$ as before. Then, $\omega_{P'/C'}^{\otimes -1} = \pi'^*(\omega_{P/C}^{\otimes -1})$ is numerically effective on P'. Let Q be a quotient vector bundle of $E' = \pi^*(E)$ with $s = \operatorname{rk} Q$. The projective bundle $\operatorname{Proj}(\bigoplus_{m=0}^{\infty} \operatorname{Sym}^m(Q)) \to C'$ gives a subvariety V' of P' with $\deg(Q) = (\mathcal{O}_{P'}(1)^s \cdot V')$ and $(\mathcal{O}_{P'}(1)^{s-1} \cdot F' \cdot V') = 1$, where F' is a fiber of f'. Since $\omega_{P'/C'}^{\otimes -1}$ is numerically effective,

$$0 \leq \left((\mathcal{O}_{P'}(r) \otimes f'^*(\det E')^{-1})^s \cdot V'\right) = r^{s-1}(r \deg(Q) - s \deg(E')).$$

Thus, $\mu(E') \leq \mu(Q)$. □

First, let us consider symmetric products of strongly semistable vector bundles.

Theorem 2.2 ($\operatorname{char}(k) \geq 0$)**.** *If E is a strongly semistable vector bundle on C, then so is $\operatorname{Sym}^n(E)$ for all $n \geq 0$.*

Proof. Taking a finite covering of C, we may assume that $\deg(E)$ is divisible by $\operatorname{rk} E$. Let θ be a line bundle on C with $\deg(\theta) = 1$. If we set $E_0 = E \otimes \theta^{\otimes -\frac{\deg(E)}{\operatorname{rk} E}}$, then $\deg(E_0) = 0$ and $\operatorname{Sym}^n(E_0) = \operatorname{Sym}^n(E) \otimes \theta^{\otimes -\frac{n \deg(E)}{\operatorname{rk} E}}$. Thus, to prove our theorem, we may assume $\deg(E) = 0$.

We assume that $\operatorname{Sym}^n(E)$ is not strongly semistable for some $n \geq 2$. By replacing C by a finite covering of C, we may assume that $\operatorname{Sym}^n(E)$ is not semistable. Let $f : P = \operatorname{Proj}(\bigoplus_{n=0}^{\infty} \operatorname{Sym}^n(E)) \to C$ be a projective bundle of E and $\mathcal{O}_P(1)$ the tautological line bundle on P. Let F be the maximal destabilizing sheaf of $\operatorname{Sym}^n(E)$. In particular, F is semistable and $\mu(F) > 0$. We consider a composition of homomorphisms

$$\alpha : f^*(F) \to f^*(\operatorname{Sym}^n(E)) \to \mathcal{O}_P(n).$$

Since $f_*(\alpha)$ induces the inclusion $F \to \operatorname{Sym}^n(E)$, α is a non-trivial homomorphism.

Fix an ample line bundle A on C. Let l be a positive integer with $l\mu(F) > n(r-1)\deg(A)$ and $(l, p) = 1$, where $p = \operatorname{char}(k)$. Here we claim that $\mathcal{O}_P(l) \otimes f^*(A)$ is ample. Let V be a s-dimensional subvariety of P. By virtue of Nakai's criterion, it is sufficient to show $(c_1(\mathcal{O}_P(l) \otimes f^*(A))^s \cdot V) > 0$. If V is contained in a fiber, our assertion is trivial. So we may assume that V is not contained in any fibers. Then,

$$(c_1(\mathcal{O}_P(l) \otimes f^*(A))^s \cdot V) = l^s(c_1(\mathcal{O}_P(1))^s \cdot V) + sl^{s-1}(c_1(\mathcal{O}_P(1))^{s-1} \cdot c_1(f^*(A)) \cdot V).$$

Since $\mathcal{O}_P(1)$ is numerically effective on P by (3) of Proposition 2.1, $(c_1(\mathcal{O}_P(1))^s \cdot V) \geq 0$. Moreover, if x is a general point of C,

$$(c_1(\mathcal{O}_P(1))^{s-1} \cdot c_1(f^*(A)) \cdot V) = \deg(A) \deg(V|_{f^{-1}(x)}) > 0.$$

Therefore, we get our claim.

Thus, there is a positive integer m such that $(\mathcal{O}_P(l) \otimes f^*(A))^{\otimes m}$ is very ample and $(m, p) = 1$. Take general elements $D_1, \ldots, D_{r-1}$ of $|(\mathcal{O}_P(l) \otimes f^*(A))^{\otimes m}|$ such that $\Gamma = D_1 \cap \cdots \cap D_{r-1}$ is a non-singular curve and $f^*(F)|_\Gamma \to \mathcal{O}_P(n)|_\Gamma$ is generically surjective. If x is a general point of C,

$$\deg(\Gamma \to C) = (D_1 \cdots D_{r-1} \cdot f^{-1}(x)) = m^{r-1}(c_1(\mathcal{O}_P(l) \otimes f^*(A))^{r-1} \cdot f^{-1}(x)) = (ml)^{r-1}.$$

Thus, $k(\Gamma)$ is separable over $k(C)$ because $(p, (ml)^{r-1}) = 1$. Hence, $f^*(F)|_\Gamma$ is semistable by (1) of Proposition 2.1. Therefore,

$$\frac{(c_1(f^*(F)) \cdot \Gamma)}{\operatorname{rk} F} \leq (c_1(\mathcal{O}_P(n)) \cdot \Gamma),$$

which implies

$$\frac{(c_1(f^*(F)) \cdot c_1(\mathcal{O}_P(l) \otimes f^*(A))^{r-1})}{\operatorname{rk} F} \leq (c_1(\mathcal{O}_P(n)) \cdot c_1(\mathcal{O}_P(l) \otimes f^*(A))^{r-1}).$$

This gives rise to

$$l^{r-1}\mu(F) \leq n(r-1)l^{r-2}\deg(A),$$

which contradicts to the choice of l with $l\mu(F) > n(r-1)\deg(A)$. □

As Corollary 2.2, we have the following main result of this section.

Corollary 2.3 ($\operatorname{char}(k) \geq 0$)**.** *If E and F are strongly semistable vector bundles on C, then so is $E \otimes F$.*

Proof. Considering a finite covering of C and tensoring line bundles, we may assume that $\deg(E) = \deg(F) = 0$ as in the same way of the beginning part of the proof of Theorem 2.2. Then, $E \oplus F$ is strongly semistable. Thus, by Theorem 2.2, $\operatorname{Sym}^2(E \oplus F)$ is strongly semistable. Here,

$$\operatorname{Sym}^2(E \oplus F) = (E \otimes F) \oplus \operatorname{Sym}^2(E) \oplus \operatorname{Sym}^2(F).$$

Therefore, we can see that $E \otimes F$ is strongly semistable. □

Corollary 2.4 ($\operatorname{char}(k) \geq 0$)**.** *Let X be a smooth projective curve over k and E a vector bundle on X. If E is strongly semistable, then $H^0(X, \operatorname{Sym}^m(\mathcal{E}nd(E)) \otimes L^{\otimes -1}) = 0$ for all $m \geq 0$ and any ample line bundles L on X.*

3. Proof of relative Bogomolov's inequality

Let X be an algebraic variety over k, x a point of X and E a coherent $\mathcal{O}_X$-module on X. We say E is *generated by global sections at x* if $H^0(X, E) \otimes \mathcal{O}_X \to E$ is surjective at x. Let us begin with the following proposition.

Proposition 3.1 ($\operatorname{char}(k) \geq 0$)**.** *Let X be a smooth algebraic variety over k, E a coherent $\mathcal{O}_X$-module, and x a point of X. If E is generated by global sections at x and E is free at x, then* $\det(E)$ *is generated by global sections at x, where* $\det(E)$ *is the determinant line bundle of E in the sense of* [8].

Proof. Let T be the torsion part of E. Then, $\det(E) = \det(E/T) \otimes \det(T)$. If we set

$$D = \sum_{\substack{P \in X, \\ \operatorname{depth}(P)=1}} \operatorname{lenght}(T_P)\overline{\{P\}},$$

then $\det(T) \simeq \mathcal{O}_X(D)$. Here since E is free at x, $x \notin \operatorname{Supp}(D)$. Thus, $\det(T)$ is generated by global sections at x. Moreover, it is easy to see that E/T is generated by global sections at x. Therefore, to prove our proposition, we may assume that E is a torsion free sheaf.

Let r be the rank of E and $\kappa(x)$ the residue field of x. Then, by our assumption, there are sections $s_1, \ldots, s_r$ of E such that $\{s_i(x)\}$ forms a basis of $E \otimes \kappa(x)$. Thus, $s = s_1 \wedge \cdots \wedge s_r$ gives rise to a section of $\det(E) = (\bigwedge^r E)^{**}$ with $s(y) \neq 0$. Hence, we get our proposition. □

Next let us consider the following proposition.

Proposition 3.2 ($\mathrm{char}(k) \geq 0$)**.** *Let $\pi : X \to Y$ be a proper morphism of algebraic varieties over k such that $k(X)$ is a finite extension of $k(X)$. Let y be a point of Y such that π is finite over y.*

(1) *Let $\phi : E \to Q$ be a homomorphism of coherent $\mathcal{O}_X$-modules. If ϕ is surjective at each point of $\pi^{-1}(y)$ and $\pi_*(E)$ is generated by global sections at y, then $\pi_*(Q)$ is generated by global sections at y.*

(2) *Let E_1 and E_2 be coherent $\mathcal{O}_X$-modules. If $\pi_*(E_1)$ and $\pi_*(E_2)$ are generated by global sections at y, then so is $\pi_*(E_1 \otimes E_2)$ at y.*

(3) *Let E be a coherent $\mathcal{O}_X$-module. If $\pi_*(E)$ is generated by global sections at y, then so is $\pi_*(\mathrm{Sym}^n(E))$ at y for every $n > 0$.*

Proof. (1) We can take an open neighborhood U of y such that π is finite over U and $\phi : E \to Q$ is surjective over $\pi^{-1}(U)$. Thus, $\pi_*(E) \to \pi_*(Q)$ is surjective at y. Hence, considering the following diagram:

$$\begin{array}{ccc} H^0(Y, \pi_*(E)) \otimes \mathcal{O}_Y & \longrightarrow & \pi_*(E) \\ \downarrow & & \downarrow \\ H^0(Y, \pi_*(Q)) \otimes \mathcal{O}_Y & \longrightarrow & \pi_*(Q), \end{array}$$

we have our assertion.

(2) Let U be an affine open neighborhood of y such that π is finite over U. We set $U = \mathrm{Spec}(A)$ for some integral domain A. Since π is finite, there is an integral domain B with $\pi^{-1}(U) = \mathrm{Spec}(B)$. Here we take B-modules M_1 and M_2 such that M_1 and M_2 give rise to $E_1|_{\pi^{-1}(U)}$ and $E_2|_{\pi^{-1}(U)}$ respectively. Then, we have a natural surjective homomorphism $M_1 \otimes_A M_2 \to M_1 \otimes_B M_2$. This shows us that $\pi_*(E) \otimes \pi_*(E_2) \to \pi_*(E_1 \otimes E_2)$ is surjective at y. Here, let us consider the following diagram:

$$\begin{array}{ccc} H^0(Y, \pi_*(E_1)) \otimes H^0(Y, \pi_*(E_2)) \otimes \mathcal{O}_Y & \longrightarrow & \pi_*(E_1) \otimes \pi_*(E_2) \\ \downarrow & & \downarrow \\ H^0(Y, \pi_*(E_1 \otimes E_2)) \otimes \mathcal{O}_Y & \longrightarrow & \pi_*(E_1 \otimes E_2), \end{array}$$

where $H^0(Y, \pi_*(E_1)) \otimes H^0(Y, \pi_*(E_2)) \otimes \mathcal{O}_Y \to \pi_*(E_1) \otimes \pi_*(E_2)$ is surjective at y by our assumption. Thus, we get (2).

(3) This is a consequence of (1) and (2) because we have a natural surjective homomorphism $E^{\otimes n} \to \mathrm{Sym}^n(E)$. □

Before starting the main theorem, we need to prepare the following formula derived from Grothendieck-Riemann-Roch theorem.

Lemma 3.3 ($\mathrm{char}(k) \geq 0$)**.** *Let X and Y be algebraic varieties over k, and $f : X \to Y$ a projective morphism over k of $\dim f = d$. Let L and A be line bundles on X. If Y is smooth, then there are elements $Z_1, \ldots, Z_d$ of $A_{\dim Y-1}(Y) \otimes \mathbb{Q}$ such that*

$$c_1\left(Rf_*(L^{\otimes n} \otimes A)\right) \cap [Y] = \frac{f_*(c_1(L)^{d+1} \cap [X])}{(d+1)!} n^{d+1} + \sum_{i=0}^{d} Z_i n^i$$

for all $n > 0$.

Proof. We use the same symbol as in [4]. First of all, $Rf_*(L^{\otimes n} \otimes A) \in K^\circ(Y)$ because Y is smooth. Thus, by [4, Theorem 18.3, (1) and (2)], i.e., Riemann-Roch theorem for singular varieties,

$$\mathrm{ch}(Rf_*(L^{\otimes n} \otimes A)) \cap \tau_Y(\mathcal{O}_Y) = f_*(\mathrm{ch}(L^{\otimes n} \otimes A) \cap \tau_X(\mathcal{O}_X)). \tag{3.3.1}$$

Since $\tau_X(\mathcal{O}_X) = [X] + \text{terms of dimension} < \dim X$ by [4, Theorem 18.3, (5)], it is easy to see that there are $T_0, \ldots, T_d \in A_{\dim Y-1}(X) \otimes \mathbb{Q}$ such that

$$\left(\mathrm{ch}(L^{\otimes n} \otimes A) \cap \tau_X(\mathcal{O}_X)\right)_{\dim Y-1} = \frac{c_1(L)^{d+1} \cap [X]}{(d+1)!} n^{d+1} + \sum_{i=0}^{d} T_i n^i.$$

Thus,

$$f_*(\mathrm{ch}(L^{\otimes n} \otimes A) \cap \tau_X(\mathcal{O}_X))_{\dim Y-1} = \frac{f_*(c_1(L)^{d+1} \cap [X])}{(d+1)!} n^{d+1} + \sum_{i=0}^{d} f_*(T_i) n^i. \tag{3.3.2}$$

On the other hand, since $\tau_Y(\mathcal{O}_Y) = [Y] + \text{terms of dimension} < \dim Y$, if we denote by S $(\dim Y - 1)$-dimensional part of $\tau_Y(\mathcal{O}_Y)$, then

(3.3.3)
$$\left(\mathrm{ch}(Rf_*(L^{\otimes n} \otimes A)) \cap \tau_Y(\mathcal{O}_Y)\right)_{\dim Y-1} = c_1(Rf_*(L^{\otimes n} \otimes A)) \cap [Y] + \mathrm{rk}(Rf_*(L^{\otimes n} \otimes A))S.$$

Here, $\mathrm{rk}(Rf_*(L^{\otimes n} \otimes A)) = \chi(X_\eta, (L^{\otimes n} \otimes A)_\eta)$ is a polynomial of n at most of degree d. Thus, combining (3.3.1), (3.3.2) and (3.3.3), we have our lemma. □

Let us start the main theorem of this paper.

Theorem 3.4 ($\mathrm{char}(k) \geq 0$)**.** *Let X be a quasi-projective variety over k, Y a smooth quasi-projective variety over k, and $f : X \to Y$ a projective morphism over k of $\dim f = 1$. Let E be a locally free sheaf of rank r on X and y a point of Y. We assume that f is flat over y, $X_y = f^{-1}(y)$ is Gorenstein, and, for any ample line bundles L on X_y, there is a positive integer $m_0(L)$ such that $H^0(X_y, \mathrm{Sym}^m(\mathcal{E}nd(E_y)) \otimes L^{\otimes -1}) = 0$ for all $m \geq m_0(L)$. Then, $\mathrm{dis}_{X/Y}(E)$ is weakly positive at y.*

Proof. We set $F = \mathcal{E}nd(E)$. Let A be a very ample line bundle on X such that $A_y \otimes \omega_{X_y}^{-1}$ is ample on X_y. Let B be a general member of $|A^{\otimes 2}|$ such that B is integral, and B is finite over y. Let $\pi : B \to Y$ be a morphism induced by f. Let H be an ample line bundle on Y such that $\pi_*(F_B) \otimes H$ and $\pi_*(A_B) \otimes H$ are generated by global sections at y, where $F_B = F|_B$ and $A_B = A|_B$.

Let $\mu : P = \mathrm{Proj}(\bigoplus_{m=0}^{\infty} \mathrm{Sym}^m(F)) \to X$ be the projective bundle and $\mathcal{O}_P(1)$ the tautological line bundle on P. We set $h = f \cdot \mu : P \to Y$. Let us consider

$$c_1\left(Rh_*((\mathcal{O}_P(1) \otimes h^*(H))^{\otimes m} \otimes \mu^*(A^{\otimes -1}) \otimes h^*(H))\right) \cap [Y]$$

for $m \gg 0$. By Lemma 3.3, there are elements $Z_0, \ldots, Z_{r^2}$ of $A_{\dim Y-1}(Y) \otimes \mathbb{Q}$ such that

$$c_1\left(Rh_*((\mathcal{O}_P(1) \otimes h^*(H))^{\otimes m} \otimes \mu^*(A^{\otimes -1}) \otimes h^*(H))\right) \cap [Y]$$
$$= \frac{h_*(c_1(\mathcal{O}_P(1) \otimes h^*(H))^{r^2+1} \cap [P])}{(r^2+1)!} m^{r^2+1} + \sum_{i=0}^{r^2} Z_i m^i.$$

Since

$$\begin{cases} \mu_*(c_1(\mathcal{O}_P(1))^{r^2+1} \cap [P]) = (c_1(F)^2 - c_2(F)) \cap [X] = -c_2(F) \cap [X], \\ \mu_*(c_1(\mathcal{O}_P(1))^{r^2} \cap [P]) = c_1(F) \cap [X] = 0, \\ \mu_*(c_1(\mathcal{O}_P(1))^{r^2-1} \cap [P]) = [X], \end{cases}$$

and $c_2(F) = 2rc_2(E) - (r-1)c_1(E)^2$, we have

$$\begin{aligned} h_*(c_1(\mathcal{O}_P(1) \otimes h^*(H))^{r^2+1} \cap [P]) &= f_*\left(-c_2(F) \cap [X] + \binom{r^2+1}{2} f^*(c_1(H)^2) \cap [X]\right) \\ &= -f_*(c_2(F) \cap [X]) \\ &= -\operatorname{dis}_{X/Y}(E). \end{aligned}$$

Moreover,

$$R\mu_*((\mathcal{O}_P(1) \otimes h^*(H))^{\otimes m} \otimes \mu^*(A^{\otimes -1}) \otimes h^*(H)) = \mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H).$$

Therefore, we get

$$\sum_{i \geq 0} (-1)^i c_1\left(R^i f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right) \cap [Y]$$
$$= -\frac{1}{(r^2+1)!} \operatorname{dis}_{X/Y}(E) m^{r^2+1} + \sum_{i=0}^{r^2} Z_i m^i.$$

Here we claim the following.

Claim 3.4.1. *If $m \gg 0$, then we have the following.*

(a) $c_1\left(R^i f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right) \cap [Y] = 0$ *for all* $i \geq 2$.
(b) $f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right) = 0$.
(c) $R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)$ *is free at* y.
(d) $R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A \otimes f^*(H)\right) = 0$ *around* y.

(a) : Let Y' be the maximal open set of Y such that f is flat over Y'. If $i \geq 2$, then the support of $R^i f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)$ is contained in $Y \setminus Y'$. Here $\operatorname{codim}(Y \setminus Y') \geq 2$. Thus, we get (a).

(b) and (c) : By our assumption, $H^0(X_y, \mathrm{Sym}^m(F_y) \otimes A_y^{\otimes -1}) = 0$ for $m \gg 0$. Thus, using upper-semicontinuity of dimension of cohomology groups, there is an open neighborhood U_m of y such that f is flat over U_m and $H^0(X_{y'}, \mathrm{Sym}^m(F_{y'}) \otimes A_{y'}^{\otimes -1}) = 0$ for all $y' \in U_m$,

which implies (b). Here, since f is flat over U_m, $\chi(X_{y'}, \mathrm{Sym}^m(F_{y'}) \otimes A_{y'}^{\otimes -1})$ is a constant with respect to $y' \in U_m$. Therefore, so is $\dim_k H^1(X_{y'}, \mathrm{Sym}^m(F_{y'}) \otimes A_{y'}^{\otimes -1})$ with respect to $y' \in U_m$. Thus, we have (c).

(d) : First note that $F^* = F$. By using Serre's duality theorem,

$$\begin{aligned}\dim_k H^1(X_y, \mathrm{Sym}^m(F_y) \otimes A_y) &= \dim_k H^0(X_y, \mathrm{Sym}^m(F_y^*) \otimes \omega_{X_y} \otimes A_y^{\otimes -1}) \\ &= \dim_k H^0(X_y, \mathrm{Sym}^m(F_y) \otimes (A_y \otimes \omega_{X_y}^{\otimes -1})^{\otimes -1}) = 0\end{aligned}$$

for $m \gg 0$. Thus, there is an open neighborhood U'_m of y such that f is flat over U'_m and

$$H^1(X_{y'}, \mathrm{Sym}^m(F_{y'}) \otimes A_{y'}) = 0$$

for all $y' \in U'_m$. Hence, we can see (d).

By (a) and (b) of Claim 3.4.1,

$$\frac{1}{(r^2+1)!}\,\mathrm{dis}_{X/Y}(E) = \frac{c_1\left(R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right) \cap [Y]}{m^{r^2+1}} + \sum_{i=0}^{r^2} \frac{Z_i}{m^{r^2+1-i}}.$$

Hence, it is sufficient to show that

$$c_1\left(R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right) \cap [Y]$$

is semi-ample at y.

Since $\pi_*(F_B \otimes \pi^*(H))$ and $\pi_*(A_B \otimes \pi^*(H))$ are generated by global sections at y, by (2) and (3) of Proposition 3.2, $\pi_*(\mathrm{Sym}^m(F_B \otimes \pi^*(H)) \otimes A_B \otimes \pi^*(H))$ is generated by global sections at y. On the other hand, a short exact sequence

$$\begin{aligned}0 \to \mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H) \to \mathrm{Sym}^m(F \otimes f^*(H)) \otimes A \otimes f^*(H) \\ \to \mathrm{Sym}^m(F_B \otimes \pi^*(H)) \otimes A_B \otimes \pi^*(H) \to 0\end{aligned}$$

gives rise to an exact sequence

$$\begin{aligned}&0 \to f_*(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A \otimes f^*(H)) \to \pi_*(\mathrm{Sym}^m(F_B \otimes \pi^*(H)) \otimes A_B \otimes \pi^*(H)) \\ &\to R^1 f_*(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)) \to R^1 f_*(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A \otimes f^*(H)).\end{aligned}$$

Thus, by (d) of Claim 3.4.1,

$$\phi_m : \pi_*(\mathrm{Sym}^m(F_B \otimes \pi^*(H)) \otimes A_B \otimes \pi^*(H)) \to R^1 f_*(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H))$$

is surjective at y. Therefore, applying (1) of Proposition 3.2 to a case where $\mathrm{id}_Y : Y \to Y$ and $\phi = \phi_m$, $R^1 f_*(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H))$ is generated by global sections at y. Moreover, by virtue of (c) of Claim 3.4.1, $R^1 f_*(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H))$ is free at y. Hence, by Proposition 3.1, $c_1(R^1 f_*(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H))) \cap [Y]$ is semi-ample at y. □

Combining Theorem 3.4 and Corollary 2.4, we have the following corollary.

Corollary 3.5 ($\mathrm{char}(k) \geq 0$)**.** *Let X be a quasi-projective variety over k, Y a smooth quasi-projective variety over k, and $f : X \to Y$ a projective morphism over k with $\dim f = 1$ and $f_*(\mathcal{O}_X) = \mathcal{O}_Y$. Let E be a locally free sheaf of rank r on X and y a point of Y. If f is*

smooth over y and E is strongly semistable on the geometric fiber $X_{\bar{y}}$ over y, then $\mathrm{dis}_{X/Y}(E)$ *is weakly positive at y.*

4. A weakly positive divisor on the moduli space of stable curves

Throughout this section, we assume that $\mathrm{char}(k) = 0$.

Fix an integer $g \geq 2$ and a polynomial $P_g(n) = (6n-1)(g-1)$. Let $H_g \subset \mathrm{Hilb}^{P_g}_{\mathbb{P}^{5g-6}}$ be a subscheme of all tri-canonically embedded stable curves over k, $Z_g \subset H_g \times \mathbb{P}^{5g-6}$ the universal tri-canonically embedded stable curves over k, and $\pi : Z_g \to H_g$ the natural projection. Let Δ be a closed subset of H_g such that π is not smooth over a point of Δ. Then, by [3, Theorem (1.6) and Corollary (1.9)], Z_g and H_g are quasi-projective and smooth over k, and Δ is a divisor with only normal crossings. Let $\Delta = \Delta_0 \cup \cdots \cup \Delta_{[g/2]}$ be the irreducible decomposition of Δ such that, if $x \in \Delta_i \setminus \mathrm{Sing}(\Delta)$, then $\pi^{-1}(x)$ is a stable curve with one node of type i. We set $U = H_g \setminus \Delta$, $H^0_g = H_g \setminus \mathrm{Sing}(\Delta_1 + \cdots + \Delta_{[g/2]})$ and $Z^0_g = \pi^{-1}(H^0_g)$. In [15, §3], we constructed a reflexive sheaf F on Z_g with the following properties.

(1) F is locally free on Z^0_g.

(2) For each $y \in U$, $F|_{\pi^{-1}(y)} = \mathrm{Ker}\left(H^0(\omega_{\pi^{-1}(y)}) \otimes \mathcal{O}_{\pi^{-1}(y)} \to \omega_{\pi^{-1}(y)}\right)$.

(3) $\mathrm{dis}_{Z_g/H_g}(F) = (8g+4)\det(\pi_*(\omega_{Z_g/H_g})) - g\Delta_0 - \sum_{i=1}^{\left[\frac{g}{2}\right]} 4i(g-i)\Delta_i$.

Actually, F can be constructed as follows. First of all, we set

$$E = \mathrm{Ker}\left(\pi^*(\pi_*(\omega_{Z_g/H_g})) \to \omega_{Z_g/H_g}\right).$$

We would like to modify E along singular fibers so that we can get our desired F. For this purpose, we consider $E^0 = E|_{Z^0_g}$. It is easy to see that E^0 is a locally free sheaf on Z^0_g. For each $i \geq 0$, we denote $\Delta_i \cap H^0_g$ by Δ^0_i. If $i \geq 1$, then there is an irreducible decomposition $\pi^{-1}(\Delta^0_i) = C^1_i \cup C^2_i$ such that the generic fiber of $\pi|_{C^1_i} : C^1_i \to \Delta^0_i$ (resp. $\pi|_{C^2_i} : C^2_i \to \Delta^0_i$) is of genus i (resp. $g-i$). Moreover, if we set

$$Q^j_i = \mathrm{Ker}\left(\left(\pi|_{C^j_i}\right)^* \left(\pi|_{C^j_i}\right)_* (\omega_{C^j_i/\Delta^0_i}) \longrightarrow \omega_{C^j_i/\Delta^0_i}\right)$$

for each $i \geq 1$ and $j = 1, 2$, then there is a natural surjective homomorphism

$$\alpha^j_i : E^0\big|_{C^j_i} \to Q^j_i.$$

Here let us consider

$$F^0 = \mathrm{Ker}\left(\bigoplus_{i=1}^{\left[\frac{g}{2}\right]} \left(\alpha^1_i \oplus \alpha^2_i\right) \ : \ E^0 \longrightarrow \bigoplus_{i=1}^{\left[\frac{g}{2}\right]} \left(Q^1_i \oplus Q^2_i\right)\right).$$

As we showed in [15, §3], F^0 is a locally free sheaf on Z^0_g with

$$\mathrm{dis}_{Z^0_g/H^0_g}(F^0) = (8g+4)\det(\pi_*(\omega_{Z^0_g/H^0_g})) - g\Delta^0_0 - \sum_{i=1}^{\left[\frac{g}{2}\right]} 4i(g-i)\Delta^0_i.$$

Let $\nu : Z^0_g \to Z_g$ be the natural inclusion map. Then F can be defined by $\nu_*(F^0)$.

Let $\overline{\mathcal{M}}_g$ (resp. $\mathcal{M}_g$) be the moduli space of stable (resp. smooth) curves of genus g over k. Let $\phi : H_g \to \overline{\mathcal{M}}_g$ be the canonical morphism. Let $\lambda, \delta_0, \ldots, \delta_{[g/2]} \in \mathrm{Pic}(\overline{\mathcal{M}}_g) \otimes \mathbb{Q}$ such that $\phi^*(\lambda) = \det(\pi_*(\omega_{Z_g/H_g}))$ and $\phi^*(\delta_i) = \Delta_i$ for all $0 \leq i \leq [g/2]$. Let us begin with the following lemma.

Lemma 4.1. *Let D be a $\mathbb{Q}$-Cartier divisor on $\overline{\mathcal{M}}_g$ and x a point of $\overline{\mathcal{M}}_g$. If $\phi^*(D)$ is weakly positive at any points of $\phi^{-1}(x)$, then D is weakly positive at x.*

Proof. It is well known that there are a finite morphism $\pi : Y \to \overline{\mathcal{M}}_g$ of normal projective varieties and a stable curve $f : X \to Y$ of genus g such that the induced morphism $Y \to \overline{\mathcal{M}}_g$ by $f : X \to Y$ is π. Then, we have a morphism $\mu : Y \to H_g$ with $\pi = \phi \cdot \mu$. Let us consider $\pi^*(D) = \mu^*(\phi^*(D))$. Here $\phi^*(D)$ is weakly positive at any points of $\phi^{-1}(x)$. Thus, by Proposition 1.6, $\pi^*(D)$ is weakly positive at any points of $\pi^{-1}(x) = \mu^{-1}(\phi^{-1}(x))$. Therefore, $\pi_*\pi^*(D) = \deg(\pi)D$ is weakly positive at x by Proposition 1.7. □

Theorem 4.2. *$(8g+4)\lambda - g\delta_0 - \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i$ is weakly positive over $\mathcal{M}_g$. In particular, it is pseudo-effective, and numerically effective over $\mathcal{M}_g$.*

Proof. Let y be a point of U. By virtue of [16], $F|_{\pi^{-1}(y)}$ is semistable. Thus, by Corollary 3.5, $\mathrm{dis}_{Z_g^0/H_g^0}(F^0)$ is weakly positive at y. Hence, by Proposition 1.8, so is $\mathrm{dis}_{Z_g/H_g}(F)$ at y because $\mathrm{codim}(H_g \setminus H_g^0) = 2$. Thus, $\mathrm{dis}_{Z_g/H_g}(F)$ is weakly positive over $U = \phi^{-1}(\mathcal{M}_g)$. Therefore, by virtue of Lemma 4.1, we get our theorem. □

As corollary, we have the following.

Corollary 4.3. *Let X be a smooth projective surface over k, Y a smooth projective curve over k, and $f : X \to Y$ a semistable curve over Y of genus $g \geq 2$. Then, we have an inequality*

$$(8g+4)\deg(f_*(\omega_{X/Y})) \geq g\delta_0(X/Y) + \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i(X/Y),$$

where $\delta_i(X/Y)$ is the number of nodes of type i in all singular fibers of f.

5. Cones of positive divisors on the moduli space of stable curves

Throughout this section, we assume that $\mathrm{char}(k) = 0$.

Let X be a projective variety over k and $\mathcal{C}$ a certain family of complete irreducible curves on X. A $\mathbb{Q}$-Cartier divisor D on X is said to be *numerically effective for $\mathcal{C}$* if $(D \cdot C) \geq 0$ for all $C \in \mathcal{C}$. We set

$$\mathrm{Nef}(X, \mathcal{C}) = \{D \in NS(X) \otimes \mathbb{Q} \mid D \text{ is numerically effective for } \mathcal{C}\}.$$

Moreover, for subsets A and B in X, we denote by Cur_B^A the set of all irreducible complete curves C on X with $C \subseteq A$ and $C \cap B \neq \emptyset$.

Proposition 5.1. *Let $\mathcal{I}_g$ be the locus of hyperelliptic curves in $\mathcal{M}_g$ and $\overline{\mathcal{M}}_g^0$ a set of all stable curves with at most one node of positive type. Then, we have*

$$\mathrm{Nef}\left(\overline{\mathcal{M}}_g, \mathrm{Cur}_{\mathcal{I}_g}^{\overline{\mathcal{M}}_g^0}\right) \subseteq \left\{ x\lambda + \sum_{i=0}^{[g/2]} y_i\delta_i \;\middle|\; \begin{array}{l} x \geq 0, \\ gx + (8g+4)y_0 \geq 0, \\ i(g-i)x + (2g+1)y_i \geq 0 \quad (1 \leq i \leq [g/2]). \end{array} \right\}.$$

Proof. Let $\overline{\mathcal{M}}_g^s$ be the Satake's compactification of $\mathcal{M}_g$. Then, $\overline{\mathcal{M}}_g^s$ is projective and $\mathrm{codim}(\overline{\mathcal{M}}_g^s \setminus \mathcal{M}_g) \geq 2$. Pick up one point $P \in \mathcal{I}_g$. If we take general hyperplane sections $H_1, \ldots, H_{3g-4}$ passing through P, then $C = H_1 \cap \ldots \cap H_{3g-4}$ is a complete irreducible curve with $C \subseteq \mathcal{M}_g$ and $P \in C$. Thus, $0 \leq (D \cdot C) = x(\lambda \cdot C)$. Hence $x \geq 0$.

To get other inequalities, we need some facts of hyperelliptic fibrations. Details can be found in [2, §4, b].

Let $\mathcal{I}_g$ be the hyperelliptic locus of $\mathcal{M}_g$ and $\overline{\mathcal{I}}_g$ the closure of $\mathcal{I}_g$ in $\overline{\mathcal{M}}_g$. For $i > 0$, $\Delta_i \cap \overline{\mathcal{I}}_g$ is irreducible. $\Delta_0 \cap \overline{\mathcal{I}}_g$ is however reducible and has $1 + [(g-1)/2]$ irreducible components, say, $\Sigma_0, \Sigma_1, \ldots, \Sigma_{[(g-1)/2]}$. Here a general point of Σ_0 represents an irreducible curve of one node, and a general point of Σ_i $(i > 0)$ represents a stable curve consisting of a curve of genus i and a curve of genus $g - i - 1$ joined at two points. The class of Σ_i in $\mathrm{Pic}(\overline{\mathcal{I}}_g) \otimes \mathbb{Q}$ is denoted by σ_i, and by abuse of notation, $\delta_i|_{\overline{\mathcal{I}}_g}$ is denoted by δ_i. Further, $\lambda|_{\overline{\mathcal{I}}_g}$ is denoted by λ. Then, by virtue of [2, Proposition (4.7)],

$$\delta_0 = \sigma_0 + 2\left(\sigma_1 + \cdots + \sigma_{[(g-1)/2]}\right)$$

and

$$(8g+4)\lambda = g\sigma_0 + \sum_{j=1}^{[(g-1)/2]} 2(j+1)(g-j)\sigma_j + \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i.$$

Let us consider D on $\overline{\mathcal{I}}_g$. Using the above relations between λ, δ_i's and σ_j's, we have

$$D = \left(\frac{g}{8g+4}x + y_0\right)\sigma_0 + 2\sum_{j=1}^{[(g-1)/2]}\left(\frac{(j+1)(g-j)}{8g+4}x + y_0\right)\sigma_j + \sum_{i=1}^{[g/2]}\left(\frac{i(g-i)}{2g+1}x + y_i\right)\delta_i.$$

Thus, it is sufficient to show the following lemma. □

Lemma 5.2. *Let $D = \sum_{j=0}^{[(g-1)/2]} a_j\sigma_j + \sum_{i=1}^{[g/2]} b_i\delta_i$ be a $\mathbb{Q}$-Cartier divisor on $\overline{\mathcal{I}}_g$. If $(D \cdot C) \geq 0$ for any irreducible curves C on $\overline{\mathcal{I}}_g$ with $C \cap \mathcal{I}_g \neq \emptyset$ and $C \subseteq \overline{\mathcal{I}}_g^0$, then $a_j \geq 0$ $(j = 0, \ldots, [(g-1)/2])$ and $b_i \geq 0$ $(i = 1, \ldots, [g/2])$, where $\overline{\mathcal{I}}_g^0 = \overline{\mathcal{I}}_g \cap \overline{\mathcal{M}}_g^0$.*

Proof. Applying Proposition A.3 to a case where $a = 0$, and contracting all (-2)-curves in all singular fibers, we have a stable fibered surface $f_0 : X_0 \to Y_0$ such that Y_0 is projective, the generic fiber of f_0 is a smooth hyperelliptic curve of genus g, f_0 is not smooth, and that every singular fiber of f_0 is an irreducible nodal curve with one node. Let $\mu_0 : Y_0 \to \overline{\mathcal{I}}_g^0$ be the induced morphism by $f_0 : X_0 \to Y_0$. Then, $\deg(\mu_0^*(\sigma_0)) > 0$, $\deg(\mu_0^*(\sigma_j)) = 0$ $(1 \leq j \leq [(g-1)/2])$ and $\deg(\mu_0^*(\delta_i)) = 0$ $(1 \leq i \leq [g/2])$. Thus, $a_0 \geq 0$ because $\deg(\mu_0^*(D)) \geq 0$.

For fixed j with $1 \leq j \leq [(g-1)/2]$, by virtue of Proposition A.3, there is a stable fibered surface $f_j : X_j \to Y_j$ such that Y_j is projective, the generic fiber of f_j is a smooth hyperelliptic curve of genus g, f_j is not smooth, and that every singular fiber of f_j is a reducible curve consisting a smooth curve of genus j and a smooth curve of genus $g - j - 1$ joined at two point. Let $\mu_j : Y_j \to \overline{\mathcal{I}}_g$ be the induced morphism by $f_j : X_j \to Y_j$. Then, $\deg(\mu_j^*(\sigma_j)) > 0$, $\deg(\mu_i^*(\sigma_l)) = 0$ $(l \neq j, 0 \leq l \leq [(g-1)/2])$ and $\deg(\mu_j^*(\delta_i)) = 0$ $(1 \leq i \leq [g/2])$. Therefore, $a_j \geq 0$.

Finally, we fix i with $1 \leq i \leq [g/2]$. Using Proposition A.2, there is a stable fibered surface $f_i' : X_i' \to Y_i'$ such that Y_i' is projective, the generic fiber of f_i' is a smooth hyperelliptic curve of genus g, f_i' is not smooth, and that every singular fiber of f_i' is reducible curve with one node of type i. Let $\mu_i' : Y_i' \to \overline{\mathcal{I}}_g$ be the induced morphism by $f_i' : X_i' \to Y_i'$. Then, $\deg({\mu_i'}^*(\delta_i)) > 0$, $\deg({\mu_i'}^*(\sigma_j)) = 0$ $(0 \leq j \leq [(g-1)/2])$ and $\deg({\mu_i'}^*(\delta_l)) = 0$ $(l \neq i, 1 \leq l \leq [g/2])$. Therefore, $b_i \geq 0$. □

Corollary 5.3. *If $\mathcal{C}$ is a set of complete irreducible curves on $\overline{\mathcal{M}}_g$ with $\mathrm{Cur}_{\mathcal{I}_g}^{\overline{\mathcal{M}}_g^0} \subseteq \mathcal{C} \subseteq \mathrm{Cur}_{\mathcal{M}_g}^{\overline{\mathcal{M}}_g}$, then*

$$\mathrm{Nef}\left(\overline{\mathcal{M}}_g, \mathcal{C}\right) = \left\{ x\lambda + \sum_{i=0}^{[g/2]} y_i\delta_i \;\middle|\; \begin{array}{l} x \geq 0, \\ gx + (8g+4)y_0 \geq 0, \\ i(g-i)x + (2g+1)y_i \geq 0 \quad (1 \leq i \leq [g/2]). \end{array} \right\}.$$

Proof. Since $\mathrm{Nef}\left(\overline{\mathcal{M}}_g, \mathcal{C}\right) \subseteq \mathrm{Nef}\left(\overline{\mathcal{M}}_g, \mathrm{Cur}_{\mathcal{I}_g}^{\overline{\mathcal{M}}_g^0}\right)$, the direction "$\subseteq$" is a consequence of Proposition 5.1. Conversely, we assume that $D = x\lambda + \sum_{i=0}^{[g/2]} y_i\delta_i$ satisfies

$$\begin{cases} x \geq 0, \\ gx + (8g+4)y_0 \geq 0, \\ i(g-i)x + (2g+1)y_i \geq 0 \quad (1 \leq i \leq [g/2]). \end{cases}$$

Then, since

$$D = \frac{x}{8g+4}\left((8g+4)\lambda - g\delta_0 - \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i\right) + \left(y_0 + \frac{g}{8g+4}x\right)\delta_0 + \sum_{i=1}^{[g/2]}\left(y_i + \frac{i(g-i)}{2g+1}x\right)\delta_i$$

and $\mathcal{C} \subseteq \mathrm{Cur}_{\mathcal{M}_g}^{\overline{\mathcal{M}}_g}$, we can see that D is numerically effective for $\mathcal{C}$ by using Theorem 4.2. □

In the same way, we can see the following.

Corollary 5.4. *If we set*

$$\mathrm{WP}(\overline{\mathcal{M}}_g; \mathcal{M}_g) = \{D \in \mathrm{Pic}(\overline{\mathcal{M}}_g) \otimes \mathbb{Q} \mid D \text{ is weakly positive over } \mathcal{M}_g\},$$

then

$$\mathrm{WP}(\overline{\mathcal{M}}_g; \mathcal{M}_g) = \left\{ x\lambda + \sum_{i=0}^{[g/2]} y_i\delta_i \;\middle|\; \begin{array}{l} x \geq 0, \\ gx + (8g+4)y_0 \geq 0, \\ i(g-i)x + (2g+1)y_i \geq 0 \quad (1 \leq i \leq [g/2]). \end{array} \right\}.$$

Proof. Note that

$$\mathrm{WP}(\overline{\mathcal{M}}_g; \mathcal{M}_g) \subseteq \mathrm{Nef}\left(\overline{\mathcal{M}}_g, \mathrm{Cur}_{\mathcal{M}_g}^{\overline{\mathcal{M}}_g}\right)$$

and that $(8g+4)\lambda - g\delta_0 - \sum_{i=1}^{[g/2]} 4i(g-i)$ and δ_i's are weakly positive over $\mathcal{M}_g$. □

Remark 5.5. In general, the weak positivity over an open set is a stronger positivity than the numerical effectivity over it. Corollary 5.3 and Corollary 5.4 however say us that, on the moduli space of stable curves $\overline{\mathcal{M}}_g$, the weak positivity over $\mathcal{M}_g$ coincides with the numerical effectivity over $\mathcal{M}_g$.

6. Effective Bogomolov's conjecture over function fields

Let X be a smooth projective surface over k, Y a smooth projective curve over k, and $f : X \to Y$ a generically smooth semistable curve of genus $g \geq 2$ over Y. Let K be the function field of Y, $\overline{K}$ the algebraic closure of K, and C the generic fiber of f. Let $j : C(\overline{K}) \to \mathrm{Jac}(C)(\overline{K})$ be a map given by $j(x) = (2g-2)x - \omega_C$ and $\| \ \|_{NT}$ the semi-norm arising from the Neron-Tate height pairing on $\mathrm{Jac}(C)(\overline{K})$. We set

$$B_C(P; r) = \left\{ x \in C(\overline{K}) \mid \|j(x) - P\|_{NT} \leq r \right\}$$

for $P \in \mathrm{Jac}(C)(\overline{K})$ and $r \geq 0$, and

$$r_C(P) = \begin{cases} -\infty & \text{if } \#\left(B_C(P;0)\right) = \infty, \\ \\ \sup\left\{ r \geq 0 \mid \#\left(B_C(P;r)\right) < \infty \right\} & \text{otherwise.} \end{cases}$$

An effective version of Bogomolov's conjecture claims the following.

Conjecture 6.1 (Effective Bogomolov's conjecture)**.** If f is non-isotrivial, then there is an effectively calculated positive number r_0 with

$$\inf_{P \in \mathrm{Jac}(C)(\overline{K})} r_C(P) \geq r_0.$$

Recently, Ullmo [17] proved that $r_C(P) > 0$ for all $P \in \mathrm{Jac}(C)(\overline{K})$ for a case where K is a number field. As far as we know, the problem to find an effectively calculated r_0 is still open.

Here we need a rather technical condition coming from calculations of green functions along singular fibers. Let $\bar{f} : \overline{X} \to Y$ be the stable model of $f : X \to Y$. Let X_y (resp. $\overline{X}_y$) be a singular fiber of f (resp. $\bar{f}$) over y, and S_y a set of nodes P on $\overline{X}_y$ such that P is not an intersection of two irreducible components of $\overline{X}_y$, i.e., a singularity of an irreducible component. Let $\pi : Z_y \to \overline{X}_y$ be the partial normalization of $\overline{X}_y$ at each node in S_y. We say X_y is a *tree of stable components* if the dual graph of Z_y is a tree graph. In other words, every node of type 0 on $\overline{X}_y$ is a singularity of an irreducible component of $\overline{X}_y$.

As application of Corollary 4.3, we get the following answer of the above conjecture, which is a generalization of [15, Theorem 5.2].

Theorem 6.2 ($\mathrm{char}(k) = 0$)**.** *If f is not smooth and every singular fiber of f is a tree of stable components, then*

$$\inf_{P \in \mathrm{Jac}(C)(\overline{K})} r_C(P) \geq \sqrt{\frac{(g-1)^2}{g(2g+1)} \left(\frac{g-1}{3} \delta_0(X/Y) + \sum_{i=1}^{\left[\frac{g}{2}\right]} 4i(g-i)\delta_i(X/Y) \right)}.$$

Before starting the proof of Theorem 6.2, let us recall several facts of green functions on a metrized graph. For details of metrized graphs, see [19].

Let G be a connected metrized graph and D an $\mathbb{R}$-divisor on G. If $\deg(D) \neq -2$, then there are a unique measure $\mu_{(G,D)}$ on G and a unique function $g_{(G,D)}$ on $G \times G$ with the following properties.

(a) $\displaystyle\int_G \mu_{(G,D)} = 1$.

(b) $g_{(G,D)}(x,y)$ is symmetric and continuous on $G \times G$.

(c) For a fixed $x \in G$, $\Delta_y(g_{(G,D)}(x,y)) = \delta_x - \mu_{(G,D)}$.

(d) For a fixed $x \in G$, $\displaystyle\int_G g_{(G,D)}(x,y)\mu_{(G,D)}(y) = 0$.

(e) $g_{(G,D)}(D,y) + g_{(G,D)}(y,y)$ is a constant for all $y \in G$.

The constant $g_{(G,D)}(D,y) + g_{(G,D)}(y,y)$ is denoted by $c(G,D)$. Further we set

$$\epsilon(G,D) = 2\deg(D)c(G,D) - g_{(G,D)}(D,D).$$

We would like to calculate the invariant $\epsilon(G,D)$ for the metrized graph G with the polarization D. First of all, let us see two examples, which will be elementary pieces for our calculations.

Example 6.3 (cf. [13, Lemma 3.2])**.** Let C be a circle of length l and O a point on C. Then,

$$g_{(C,0)}(O,O) = \frac{l}{12} \quad \text{and} \quad \epsilon(C,0) = 0.$$

Example 6.4 (cf. [15, Lemma 4.4])**.** Let G be a segment of length l, and P and Q terminal points of G. Let a and b be real numbers with $a+b \neq 0$, and D an $\mathbb{R}$-divisor on G given by $D = (2a-1)P + (2b-1)Q$. Then,

$$\epsilon(G,D) = \left(\frac{4ab}{a+b} - 1\right) l, \quad g_{(G,D)}(P,P) = \frac{b^2}{(a+b)^2} l \quad \text{and} \quad g_{(G,D)}(Q,Q) = \frac{a^2}{(a+b)^2} l.$$

Let G_1 and G_2 be metrized graphs. Fix points $x_1 \in G_1$ and $x_2 \in G_2$. The one point sum $G_1 \vee G_2$ with respect to x_1 and x_2, defined by $G_1 \times \{x_2\} \cup \{x_1\} \times G_2$ in $G_1 \times G_2$, is a metrized graph obtained by joining $x_1 \in G_1$ and $x_2 \in G_2$. The joining point, which is $\{x_1\} \times \{x_2\}$ in $G_1 \times G_2$, is denoted by $j(G_1 \vee G_2)$. Any $\mathbb{R}$-divisor on G_i $(i = 1,2)$ can be viewed as an $\mathbb{R}$-divisor on $G_1 \vee G_2$. Then, our basic tools for our calculations is the following.

Proposition 6.5 (cf. [15, Proposition 4.2])**.** *Let G_1 and G_2 be connected metrized graphs, and D_1 and D_2 $\mathbb{R}$-divisors on G_1 and G_2 respectively with $\deg(D_i) \neq -2$ $(i = 1,2)$. Let $G = G_1 \vee G_2$, $O = j(G_1 \vee G_2)$, and $D = D_1 + D_2$ on $G_1 \vee G_2$. If $\deg(D_1 + D_2) \neq -2$, then we have the following formulae, where $d_i = \deg(D_i)$ $(i = 1,2)$ and $r_{G_2}(O,P)$ is the resistance between O and P on G_2.*

(1) *If $P \in G_2$, then*

$$\begin{aligned} g_{(G,D)}(P,P) = {} & \frac{d_1}{d_1+d_2+2} r_{G_2}(O,P) + \frac{d_2+2}{d_1+d_2+2} g_{(G_2,D_2)}(P,P) \\ & - \frac{d_1(d_2+2)}{(d_1+d_2+2)^2} g_{(G_2,D_2)}(O,O) + \frac{(d_1+2)^2}{(d_1+d_2+2)^2} g_{(G_1,D_1)}(O,O). \end{aligned}$$

(2)

$$\epsilon(G,D)=\epsilon(G_1,D_1)+\epsilon(G_2,D_2)+\frac{2d_2(d_1+2)g_{(G_1,D_1)}(O,O)+2d_1(d_2+2)g_{(G_2,D_2)}(O,O)}{d_1+d_2+2}.$$

Combining the above proposition and Example 6.3, we have the following.

Corollary 6.6. *Let G be a connected metrized graph and D an $\mathbb{R}$-divisor on G with $\deg(D)\neq -2$. Let C be a circle of length l. Then,*

$$\epsilon(G\vee C,D)=\epsilon(G,D)+\frac{\deg(D)}{3(\deg D+2)}l.$$

Let G be a connected metrized graph. We assume that G is a tree, i.e., there is no loop in G. Let $\mathrm{Vert}(G)$ (resp. $\mathrm{Ed}(G)$) be the set of vetexes (resp. edges) of G. For a function $\alpha:\mathrm{Vert}(G)\to\mathbb{R}$, we define a divisor $D(\alpha)$ on G as

$$D(\alpha)=\sum_{x\in\mathrm{Vert}(G)}(2\alpha(x)-2+v(x))x,$$

where $v(x)$ is the number of branches starting from x. It is easy to see that

$$\deg(D(\alpha))+2=2\sum_{x\in\mathrm{Vert}(G)}\alpha(x).$$

To give an exact formula for $\epsilon(G,D(\alpha))$, we need to introduce the following notation. Let e be an edge of G, P and Q terminal points of e, and $e^\circ=e\setminus\{P,Q\}$. Since G is a connected tree, there are two connected sub-graphs G'_e and G''_e such that $G\setminus e^\circ=G'_e\coprod G''_e$. Then, we have the following.

Proposition 6.7. *With the same notation as above, if $\alpha(x)\geq 0$ for all $x\in\mathrm{Vert}(G)$ and $\sum_{x\in\mathrm{Vert}(G)}\alpha(x)\neq 0$, then*

$$\epsilon(G,D(\alpha))=\sum_{e\in\mathrm{Ed}(G)}\left(\frac{4\left(\sum_{x\in\mathrm{Vert}(G'_e)}\alpha(x)\right)\left(\sum_{x\in\mathrm{Vert}(G''_e)}\alpha(x)\right)}{\sum_{x\in\mathrm{Vert}(G)}\alpha(x)}-1\right)l(e),$$

where $l(e)$ is the length of e.

Proof. For a positive number t, we set $\alpha_t(x)=\alpha(x)+t$. Then, it is easy to see that

$$\lim_{t\downarrow 0}\epsilon(G,D(\alpha_t))=\epsilon(G,D(\alpha)).$$

Thus, in order to prove our proposition, we may assume that $\alpha(x)>0$ for all $x\in\mathrm{Vert}(G)$.

We fix $P\in\mathrm{Vert}(G)$. For $e\in\mathrm{Ed}(G)$, we denote by $G_{P,e}$ the connected component of $G\setminus e^\circ$ not containing P, i.e., if $P\notin G'_e$, then $G_{P,e}=G'_e$; otherwise, $G_{P,e}=G''_e$. With this notation, let us consider the following claim.

Claim 6.7.1.

$$g_{(G,D(\alpha))}(P,P)=\sum_{e\in\mathrm{Ed}(G)}\frac{\left(\sum_{x\in\mathrm{Vert}(G_{P,e})}\alpha(x)\right)^2}{\left(\sum_{x\in\mathrm{Vert}(G)}\alpha(x)\right)^2}l(e).$$

We prove this claim by induction on $\#(\mathrm{Ed}(G))$. If $\#(\mathrm{Ed}(G)) = 0, 1$, then our assertion is obvious by Example 6.4. Thus, we may assume that $\#(\mathrm{Ed}(G)) \geq 2$.

First, we suppose that P is not a terminal point. Let G' be one branch starting from P, and G'' a connected sub-graph such that $G' \cup G'' = G$ and $G' \cap G'' = \{P\}$. We define $\alpha' : \mathrm{Vert}(G') \to \mathbb{R}$ and $\alpha'' : \mathrm{Vert}(G'') \to \mathbb{R}$ by

$$\alpha'(x) = \begin{cases} 1 & \text{if } x = P \\ \alpha(x) & \text{otherwise} \end{cases}$$

and $\alpha'' = \alpha|_{\mathrm{Vert}(G'')}$. Then, we have $G = G' \vee G''$ and $D(\alpha) = D(\alpha') + D(\alpha'')$. Thus, using (1) of Proposition 6.5 and hypothesis of induction, we can easily see our claim.

Next we suppose that P is a terminal point. Pick up $e \in \mathrm{Ed}(G)$ such that P is a terminal of e. Let O be another terminal of e. We set $G' = e$ and $G'' = (G \setminus e) \cup \{O\}$. Moreover, we define $\alpha' : \mathrm{Vert}(G') = \{P, O\} \to \mathbb{R}$ and $\alpha'' : \mathrm{Vert}(G'') \to \mathbb{R}$ by $\alpha'(P) = \alpha(P)$, $\alpha'(O) = 1$ and $\alpha'' = \alpha|_{\mathrm{Vert}(G'')}$. Then, $G = G' \vee G''$ and $D(\alpha) = D(\alpha') + D(\alpha'')$. Thus, using (1) of Proposition 6.5, Example 6.4 and hypothesis of induction, we can see our claim after easy calculations.

Let us go back to the proof of Proposition 6.7. We prove it by induction on $\#(\mathrm{Ed}(G))$. If $\#(\mathrm{Ed}(G)) = 0, 1$, then our assertion comes from Example 6.4. Thus, we may assume that $\#(\mathrm{Ed}(G)) \geq 2$. Let us pick up a terminal edge e of G. Let $\{O, P\}$ be terminals of e such that P gives a terminal of G. We set $G_1 = e$ and $G_2 = (G \setminus e) \cup \{O\}$. Moreover, we define $\alpha_1 : \{O, P\} = \mathrm{Vert}(G_1) \to \mathbb{R}$ and $\alpha_2 : \mathrm{Vert}(G_2) \to \mathbb{R}$ by $\alpha_1(O) = 1$, $\alpha_1(P) = \alpha(P)$, and $\alpha_2 = \alpha|_{\mathrm{Vert}(G_2)}$. Then, $G = G_1 \vee G_2$ and $D(\alpha) = D(\alpha_1) + D(\alpha_2)$. Thus, if we set

$$\begin{cases} A = \sum_{x \in \mathrm{Vert}(G)} \alpha(x), \\ a = \alpha(P), \\ A_{e'} = \sum_{x \in \mathrm{Vert}(G_{O,e'})} \alpha(x) \quad \text{for } e' \in \mathrm{Vert}(G) \setminus \{e\}, \end{cases}$$

then, by (2) of Proposition 6.5, Example 6.4, Claim 6.7.1 and hypothesis of induction, we have

$$\begin{aligned} \epsilon(G, D(\alpha)) &= \left(\frac{4a}{a+1} - 1\right) l(e) + \sum_{e' \in \mathrm{Vert}(G) \setminus \{e\}} \left(\frac{4A_{e'}(A - a - A_{e'})}{A - a} - 1\right) l(e') \\ &\quad + \frac{4(A-a-1)(a+1)}{A} \frac{a^2}{(a+1)^2} l(e) + \frac{4a(A-a)}{A} \sum_{e' \in \mathrm{Vert}(G) \setminus \{e\}} \frac{A_{e'}^2}{(A-a)^2} l(e') \\ &= \left(\frac{4a(A-a)}{A} - 1\right) l(e) + \sum_{e' \in \mathrm{Vert}(G) \setminus \{e\}} \left(\frac{4A_{e'}(A - A_{e'})}{A} - 1\right) l(e'). \end{aligned}$$

Therefore, we get our proposition. □

Corollary 6.8 ($\mathrm{char}(k) \geq 0$)**.** *Let X be a smooth projective surface over k, Y a smooth projective curve over k, and $f : X \to Y$ a generically smooth semistable curve over Y of genus $g \geq 2$. Let X_y be a singular fiber of f over $y \in Y$, and $X_y = C_1 + \cdots + C_n$ the irreducible decomposition of X_y. Let G_y be the metrized graph given by the configuration of X_y, v_i the*

vertex of G_y corresponding to C_i, and ω_y a divisor on G_y defined by $\omega_y = \sum_i (\omega_{X/Y} \cdot C_i) v_i$. If X_y is a tree of stable components, then

$$\epsilon(G_y, \omega_y) = \frac{g-1}{3g}\delta_0(X_y) + \sum_{i=1}^{\left[\frac{g}{2}\right]} \left(\frac{4i(g-i)}{g} - 1 \right) \delta_i(X_y).$$

Proof. Let $\bar{f} : \overline{X} \to Y$ be the stable model of $f : X \to Y$, and S_y a set of nodes P on $\overline{X}_y$ such that P is a singularity of an irreducible component. Let $\pi : Z_y \to \overline{X}_y$ be the partial normalization of $\overline{X}_y$ at each node in S_y. Let $\overline{G}_y$ be the dual graph of Z_y. Let $l_1, \cdots, l_r$ be circles in G_y corresponding to nodes in S_y. Then, $G_y = \overline{G}_y \vee l_1 \vee \cdots \vee l_r$. Moreover, if g_i is the arithmetic genus of C_i and $\alpha : \mathrm{Vert}(G_y) \to \mathbb{R}$ is given by $\alpha(v_i) = g_i$, then $\omega_y = D(\alpha)$. Here, by virtue of Proposition 6.7,

$$\epsilon(\overline{G}_y, \omega_y) = \sum_{i=1}^{\left[\frac{g}{2}\right]} \left(\frac{4i(g-i)}{g} - 1 \right) \delta_i(X_y).$$

Therefore, it follows from Corollary 6.6 that

$$\epsilon(G_y, \omega_y) = \frac{g-1}{3g}\delta_0(X_y) + \sum_{i=1}^{\left[\frac{g}{2}\right]} \left(\frac{4i(g-i)}{g} - 1 \right) \delta_i(X_y).$$

□

Let us start the proof of Theorem 6.2. First of all, note the following fact (cf. [19, Theorem 5.6], [13, Corollary 2.3] or [14, Theorem 2.1]). If $(\omega^a_{X/Y} \cdot \omega^a_{X/Y})_a > 0$, then

$$\inf_{P \in \mathrm{Jac}(C)(\overline{K})} r_C(P) \geq \sqrt{(g-1)(\omega^a_{X/Y} \cdot \omega^a_{X/Y})_a},$$

where $(\ \cdot\ )_a$ is the admissible pairing.

By the definition of admissible pairing, we can see

$$\left(\omega^a_{X/Y} \cdot \omega^a_{X/Y}\right)_a = \left(\omega_{X/Y} \cdot \omega_{X/Y}\right) - \sum_{y \in Y} \epsilon(G_y, \omega_y).$$

On the other hand, by Corollary 4.3, we have

$$(8g+4)\deg(f_*(\omega_{X/Y})) \geq g\delta_0(X/Y) + \sum_{i=1}^{\left[\frac{g}{2}\right]} 4i(g-i)\delta_i(X/Y).$$

Thus, using Noether formula, the above inequality implies

$$(\omega_{X/Y} \cdot \omega_{X/Y}) \geq \frac{g-1}{2g+1}\delta_0(X/Y) + \sum_{i=1}^{\left[\frac{g}{2}\right]} \left(\frac{12i(g-i)}{2g+1} - 1 \right) \delta_i(X/Y).$$

Therefore, we have our theorem by Corollary 6.8. □

Moreover, using Ullmo's result [17] and Proposition 6.7, we have the following.

Corollary 6.9. *Let K be a number field, O_K the ring of integers, and $f : X \to \mathrm{Spec}(O_K)$ a regular semistable arithmetic surface of genus $g \geq 2$ over O_K. Let S be a subset of $\mathrm{Spec}(O_K) \setminus \{0\}$ such that the stable model of the geometric fiber $X_{\bar{P}}$ at P is a tree of stable components for each $P \in S$. Then, we have*

$$\left(\omega^{Ar}_{X/O_K} \cdot \omega^{Ar}_{X/O_K}\right) > \sum_{P \in S} \left\{ \frac{g-1}{3g}\delta_0(X_{\bar{P}}) + \sum_{i=1}^{[g/2]} \left(\frac{4i(g-i)}{g} - 1 \right) \delta_i(X_{\bar{P}}) \right\} \log \#(O_K/P).$$

7. Generalization to higher dimensional fibrations

In this section, we consider a generalization of Corollary 3.5 to higher dimensional fibrations.

First of all, let us recall the definition of semistability of vector bundles. Let V be a smooth projective variety of dimension d over k, and $H_1, \ldots, H_{d-1}$ ample line bundles on V. A vector bundle E on V is said to be semistable with respect to $H_1, \ldots, H_{d-1}$ if for any non-zero subsheaves G of E,

$$\frac{(c_1(G) \cdot c_1(H_1) \cdots c_1(H_{d-1}))}{\mathrm{rk}\, G} \leq \frac{(c_1(E) \cdot c_1(H_1) \cdots c_1(H_{d-1}))}{\mathrm{rk}\, E}.$$

Let us start to consider a generalization of Corollary 3.5 to higher dimensional fibrations. Let $f : X \to Y$ be a projective morphism of quasi-projective varieties over k such that the generic fiber of $f : X \to Y$ is a geometrically irreducible smooth variety of dimension $d \geq 1$. Let $H_1, \ldots, H_{d-1}$ be line bundles on X and E a vector bundle on X of rank r. Then, $((2rc_2(E) - (r-1)c_1(E)^2) \cdot c_1(H_1) \cdots c_1(H_{d-1})) \cap [X]$ is a cycle on X of dimension $\dim Y - 1$. Thus, $f_*\left(((2rc_2(E) - (r-1)c_1(E)^2) \cdot c_1(H_1) \cdots c_1(H_{d-1})) \cap [X]\right)$, denoted by $\mathrm{dis}_{X/Y}(E; H_1, \ldots, H_{d-1})$, is a divisor on Y. Then, we have the following theorem.

Theorem 7.1 ($\mathrm{char}(k) = 0$)**.** *We assume that Y is smooth over k and $H_1, \ldots, H_{d-1}$ are ample. Let y be a point of Y such that f is smooth over y and $E|_{X_{\bar{y}}}$ is semistable with respect to $H_1|_{X_{\bar{y}}}, \ldots, H_{d-1}|_{X_{\bar{y}}}$. Then, the discriminant divisor $\mathrm{dis}_{X/Y}(E; H_1, \ldots, H_{d-1})$ is weakly positive at y.*

Proof. We prove this theorem by induction on d. If $d = 1$, then our assertion is nothing more than Corollary 3.5. So we assume $d \geq 2$. We choose a sufficiently large integer n so that $H_{d-1}^{\otimes n}$ is very ample, i.e., there is an embedding $\iota : X \hookrightarrow \mathbb{P}^N$ with $H_{d-1}^{\otimes n} \simeq \iota^*(\mathcal{O}_{\mathbb{P}^N}(1))$. By Bertini's theorem, we can find a general member $\Gamma \in |\mathcal{O}_{\mathbb{P}^N}(1)|$ such that $X \cap \Gamma$ is integral and $f^{-1}(y) \cap \Gamma$ is smooth. We set $Z = X \cap \Gamma$ and $g = f|_Z : Z \to Y$. Let $X_{\bar{\eta}}$ and $Z_{\bar{\eta}}$ be geometric generic fibers of $f : X \to Y$ and $g : Z \to Y$ respectively. Then, we have an exact sequence

$$0 \to H_{d-1}^{\otimes -n}\big|_{X_{\bar{\eta}}} \to \mathcal{O}_{X_{\bar{\eta}}} \to \mathcal{O}_{Z_{\bar{\eta}}} \to 0.$$

Since n is sufficiently large, $H^0(H_{d-1}^{\otimes -n}\big|_{X_{\bar{\eta}}}) = H^1(H_{d-1}^{\otimes -n}\big|_{X_{\bar{\eta}}}) = 0$. Thus, $\dim H^0(\mathcal{O}_{Z_{\bar{\eta}}}) = 1$, which means that $Z_{\bar{\eta}}$ is connected. On the other hand, $Z_{\bar{\eta}}$ is smooth over $\overline{k(Y)}$ because $g : Z \to Y$ is smooth over y. Hence $Z_{\bar{\eta}}$ is irreducible. Moreover, since n is sufficiently large, $g^{-1}(y) \in \left| H_{d-1}^{\otimes n}\big|_{f^{-1}(y)} \right|$ and $g^{-1}(y)$ is smooth, by virtue of [12, Theorem 3.1], $E|_{Z_{\bar{y}}}$

is semistable with respect to $H_1|_{Z_{\bar{y}}}, \ldots, H_{d-2}|_{Z_{\bar{y}}}$. Therefore, by hypothesis of induction, $\mathrm{dis}_{Z/Y}(E|_Z; H_1|_Z, \ldots, H_{d-2}|_Z)$ is weakly positive at y. On the other hand, we have

$$\begin{aligned}\mathrm{dis}_{Z/Y}(E|_Z; H_1|_Z, \ldots, H_{d-2}|_Z) &= \mathrm{dis}_{X/Y}(E; H_1, \ldots, H_{d-2}, H_{d-1}^{\otimes n}) \\ &= n\,\mathrm{dis}_{X/Y}(E; H_1, \ldots, H_{d-1}).\end{aligned}$$

Hence, $\mathrm{dis}_{X/Y}(E; H_1, \ldots, H_{d-1})$ is weakly positive at y. □

Appendix A. A construction of a certain fibration of hyperelliptic curves

In this section, we would like to construct a certain fibration of hyperelliptic curves. Throughout this section, we assume that $\mathrm{char}(k) = 0$.

Let us begin with the following lemma.

Lemma A.1. *For non-negative integers a_1 and a_2, there are a morphism $f_1 : X_1 \to Y_1$ of smooth projective varieties over k, an effective divisor D_1 on X_1, a line bundle L_1 on X_1, and a line bundle M_1 on Y_1 with the following properties.*

(1) $\dim X_1 = 2$ *and* $\dim Y_1 = 1$.
(2) *Let Σ_1 be the set of all critical values of f_1, i.e., $P \in \Sigma_1$ if and only if $f_1^{-1}(P)$ is a singular variety. Then, for any $P \in Y_1 \setminus \Sigma_1$, $f_1^{-1}(P)$ is a smooth rational curve.*
(3) *$\Sigma_1 \neq \emptyset$, and for any $P \in \Sigma_1$, $f_1^{-1}(P)$ is a reduced curve consisting of two smooth rational curves E_P^1 and E_P^2 joined at one point transversally.*
(4) *D_1 is smooth over k and $f_1|_{D_1} : D_1 \to Y_1$ is etale.*
(5) *$(E_P^1 \cdot D_1) = a_1 + 1$ and $(E_P^2 \cdot D_1) = a_2 + 1$ for any $P \in \Sigma_1$. Moreover, D_1 does not pass through $E_P^1 \cap E_P^2$.*
(6) *There is a map $\sigma : \Sigma_1 \to \{1, 2\}$ such that*

$$D_1 \in \left| L_1^{\otimes a_1 + a_2 + 2} \otimes f_1^*(M_1) \otimes \mathcal{O}_{X_1}\left(- \sum_{P \in \Sigma_1} (a_{\sigma(P)} + 1) E_P^{\sigma(P)} \right) \right|.$$

(7) $\deg(M_1)$ *is divisible by $(a_1 + 1)(a_2 + 1)$.*



Proof. First of all, let us consider a function $\theta(x)$ defined by

$$\theta(x) = (a_1 + a_2 + 1) \int_0^x t^{a_1}(t-1)^{a_2} dt.$$

Then, $\theta(x)$ is a monic polynomial of degree $a_1 + a_2 + 1$ over $\mathbb{Q}$. Moreover, it is easy to see that

$$\theta'(x) = (a_1 + a_2 + 1)x^{a_1}(x-1)^{a_2}, \qquad \theta(0) = 0 \quad \text{and} \quad \theta(1) = (-1)^{a_2}(a_1 + a_2 + 1)\frac{(a_1)!(a_2)!}{(a_1 + a_2)!}.$$

Thus, there are distinct non-zero algebraic numbers $\alpha_1, \ldots, \alpha_{a_2}$ and $\beta_1, \ldots, \beta_{a_1}$ such that

$$\theta(x) = x^{a_1+1}(x - \alpha_1) \cdots (x - \alpha_{a_2})$$

and

$$\theta(x) - \theta(1) = (x-1)^{a_2+1}(x - 1 - \beta_1) \cdots (x - 1 - \beta_{a_1}).$$

Here we set

$$F(X, Y) = Y^{a_1+a_2+1}\theta(X/Y) = X^{a_1+1}(X - \alpha_1 Y) \cdots (X - \alpha_{a_2} Y)$$

and

$$G(X, Y, S, T) = TF(X, Y) - Y^{a_1+a_2+1}S.$$

Then, F is a homogeneous polynomial of degree a_1+a_2+1 over $\mathbb{Q}$, and G is a bi-homogeneous polynomial of bi-degree $(a_1+a_2+1, 1)$ in $\mathbb{Q}[X, Y] \otimes_{\mathbb{Q}} \mathbb{Q}[S, T]$. Let D' (resp. D'') be a curve on $\mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)}$ given by an equation $\{G = 0\}$ (resp. $\{Y = 0\}$), where $\mathbb{P}^1_{(X,Y)} = \mathrm{Proj}(k[X, Y])$ and $\mathbb{P}^1_{(S,T)} = \mathrm{Proj}(k[S, T])$. Moreover, we set $D = D' + D''$. Let $p : \mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)} \to \mathbb{P}^1_{(X,Y)}$ and $q : \mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)} \to \mathbb{P}^1_{(S,T)}$ be the natural projections. Then, D' (resp. D'') is an element of a linear system $|p^*(\mathcal{O}_{\mathbb{P}^1}(a_1 + a_2 + 1)) \otimes q^*(\mathcal{O}_{\mathbb{P}^1}(1))|$ (resp $|p^*(\mathcal{O}_{\mathbb{P}^1}(1))|$). Thus, $D \in |p^*(\mathcal{O}_{\mathbb{P}^1}(a_1 + a_2 + 2)) \otimes q^*(\mathcal{O}_{\mathbb{P}^1}(1))|$, $(D' \cdot D'') = 1$ and $D' \cap D'' = \{((1 : 0), (1 : 0))\}$. Here we claim the following.

Claim A.1.1. (a) *D' is a smooth rational curve.*

(b) *Let $\pi' : D' \to \mathbb{P}^1_{(S,T)}$ be a morphism induced by the projection $q : \mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)} \to \mathbb{P}^1_{(S,T)}$. If we set $Q_1 = ((0 : 1), (0 : 1))$, $Q_2 = ((1 : 1), (\theta(1), 1))$ and $Q_3 = ((1 : 0), (1 : 0))$, then the set of ramification points of π' is $\{Q_1, Q_2, Q_3\}$. Further, the ramification indexes at Q_1, Q_2 and Q_3 are $a_1 + 1$, $a_2 + 1$ and $a_1 + a_2 + 1$ respectively.*

Proof. (a) Since $F(X, Y)$ has no factor of Y, a morphism $e : \mathbb{P}^1_{(X,Y)} \to D'$ given by

$$e(x : y) = \big((x : y), (F(x, y) : y^{a_1+a_2+1})\big)$$

is well defined. Moreover, if we set $e' = p|_{D'}$, then it is easy to see that $e \cdot e' = \mathrm{id}_{D'}$ and $e' \cdot e = \mathrm{id}_{\mathbb{P}^1}$. Thus, D' is a smooth rational curve.

(b) Pick up a point $(\lambda : \mu) \in \mathbb{P}^1_{(S,T)}$. Then, $G_{(\lambda,\mu)} = \mu F(X, Y) - Y^{a_1+a_2+1}\lambda$ is a homogeneous polynomial of degree $a_1 + a_2 + 1$.

First, we assume that $\mu \neq 0$, hence we may assume that $\mu = 1$. Then, Y is not a factor of $G_{(\lambda,1)}(X, Y)$, which means that ${\pi'}^{-1}((\lambda : 1))$ sits in an affine open set $\mathrm{Spec}(k[X/Y, S/T])$. Thus,

$${\pi'}^{-1}((\lambda : 1)) = \{((\gamma : 1), (\lambda : 1)) \mid \theta(\gamma) - \lambda = 0\}.$$

Hence, in order to get ramification points of π', we need to see multiple roots of $\phi(x) = \theta(x) - \lambda$. Here we will check that $\phi(x)$ has a multiple root if and only if λ is either 0 or $\theta(1)$. Moreover, if λ is 0 (resp. $\theta(1)$), then 0 (resp. 1) is the only multiple root of $\phi(x)$ with multiplicity $a_1 + 1$ (resp. $a_2 + 1$).

Let γ be a multiple root of $\phi(x) = 0$. Then, $\phi(\gamma) = \phi'(\gamma) = 0$. Here,

$$\phi'(x) = (a_1 + a_2 + 1)x^{a_1}(x-1)^{a_2}.$$

Thus, γ is either 0 or 1. If $\gamma = 0$, then $\lambda = \theta(0) = 0$. If $\gamma = 1$, then $\lambda = \theta(1)$. In the same way, we can easily check the remaining part of our assertion.

Therefore, we get two ramification points Q_1 and Q_2 whose ramification indexes are $a_1 + 1$ and $a_2 + 1$ respectively.

Next, we assume that $\mu = 0$, hence we may assume $\lambda = 1$. Then, $G_{(\lambda,\mu)} = -Y^{a_1+a_2+1}$. Thus, P_3 is a ramification point whose ramification index is $a_1 + a_2 + 1$. □

Here we set $P_i = q(Q_i)$ $(i = 1, 2, 3)$, $b_1 = a_1 + 1$, $b_2 = a_2 + 1$, and $b_3 = a_1 + a_2 + 1$.

Claim A.1.2. *There is a cyclic covering $h_1 : Y_1 \to \mathbb{P}^1_{(S,T)}$ of smooth projective curves such that $\deg(h_1) = 3b_1b_2b_3$ and that, for any $i = 1, 2, 3$ and any $P \in h_1^{-1}(P_i)$, the ramification index of h_1 at P is b_i.*

Proof. Since $b_1b_2 + b_2b_3 + b_3b_1 \leq 3b_1b_2b_3$, there is an effective and reduced divisor d on $\mathbb{P}^1_{(S,T)}$ such that $P_i \notin \mathrm{Supp}(d)$ for each $i = 1, 2, 3$ and

$$b_2b_3P_1 + b_3b_1P_2 + b_1b_2P_3 + d \in |\mathcal{O}_{\mathbb{P}^1}(3b_1b_2b_3)|.$$

Let w be a section of $H^0(\mathcal{O}_{\mathbb{P}^1}(3b_1b_2b_3))$ with $\mathrm{div}(w) = b_2b_3P_1 + b_3b_1P_2 + b_1b_2P_3 + d$. Then, w gives a ring structure on $\bigoplus_{i=0}^{3b_1b_2b_3-1} \mathcal{O}_{\mathbb{P}^1}(-i)$. Let Y_1 be the normalization of

$$\mathrm{Spec}\left(\bigoplus_{i=0}^{3b_1b_2b_3-1} \mathcal{O}_{\mathbb{P}^1}(-i)\right)$$

and $h_1 : Y_1 \to \mathbb{P}^1$ the induced morphism. Then, by our choice of w, it is easy to see that $h_1 : Y_1 \to \mathbb{P}^1$ satisfies the desired properties. □

Let $p_1 : \mathbb{P}^1_{(X,Y)} \times Y_1 \to \mathbb{P}^1_{(X,Y)}$ and $q_1 : \mathbb{P}^1_{(X,Y)} \times Y_1 \to Y_1$ be the natural projections, and $u_1 = \mathrm{id} \times h_1 : \mathbb{P}^1_{(X,Y)} \times Y_1 \to \mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)}$. Then, we have a commutative diagram:

$$\begin{array}{ccc} \mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)} & \xleftarrow{u_1} & \mathbb{P}^1_{(X,Y)} \times Y_1 \\ \Big\downarrow{\scriptstyle q} & & \Big\downarrow{\scriptstyle q_1} \\ \mathbb{P}^1_{(S,T)} & \xleftarrow[h_1]{} & Y_1 \end{array}$$

We set $h_1^{-1}(P_1)$, $h_1^{-1}(P_2)$ and $h_1^{-1}(P_3)$ as follows.

$$\begin{cases} h_1^{-1}(P_1) = \{P_{1,1}, \ldots, P_{1,3b_2b_3}\}, \\ h_1^{-1}(P_2) = \{P_{2,1}, \ldots, P_{1,3b_3b_1}\}, \\ h_1^{-1}(P_3) = \{P_{3,1}, \ldots, P_{3,3b_1b_2}\}. \end{cases}$$

Then, there is a unique $Q_{i,j}$ on $\mathbb{P}^1_{(X,Y)} \times Y_1$ with $q_1(Q_{i,j}) = P_{i,j}$ and $u_1(Q_{i,j}) = Q_i$.

Claim A.1.3. (a) *$u_1^*(D)$ is etale over Y_1 outside $\{Q_{i,j}\}_{i,j}$. In particular, $u_1^*(D)$ is smooth over k outside $\{Q_{i,j}\}_{i,j}$.*

(b) *If we set $c_1 = a_1 + 1$, $c_2 = a_2 + 1$ and $c_3 = a_1 + a_2 + 2$, then $u_1^*(D)$ has an ordinary c_i-fold point at $Q_{i,j}$ for all i, j. Moreover, each tangent of $u_1^*(D)$ at $Q_{i,j}$ is different from the fiber $q_1^{-1}(P_{i,j})$.*

Proof. (a) is trivial because $q|_D : D \to \mathbb{P}^1_{(S,T)}$ is etale outside $\{Q_1, Q_2, Q_3\}$. Since $u_1^*(D'') = p_1^{-1}((1:0))$, in order to see (b), it is sufficient to check the following. $u_1^*(D')$ has an ordinary b_i-fold point at $Q_{i,j}$ for all i, j. Moreover, for i=1,2, each tangent of $u_1^*(D')$ at $Q_{i,j}$ is different from the fiber $q_1^{-1}(P_{i,j})$, and each tangent of $u_1^*(D')$ at $Q_{3,j}$ is different from the fiber $q_1^{-1}(P_{3,j})$ and $p_1^{-1}((1:0))$.

First we assume $i = 1$. Let z be a local parameter of Y_1 at $P_{1,j}$, $x = X/Y$, and $s = S/T$. Then, (x, z) gives a local parameter of $\mathbb{P}^1_{(X,Y)} \times Y_1$ at $Q_{1,j}$. Since $s = v(z)z^{a_1+1}$ for some $v(z)$ with $v(0) \neq 0$, $u_1^*(D')$ is defined by

$$x^{a_1+1}(x - \alpha_1) \cdots (x - \alpha_{a_2}) - v(z)z^{a_1+1} = 0$$

around $Q_{1,j}$. Thus, since $\alpha_1 \cdots \alpha_{a_2} \neq 0$, $Q_{1,j}$ is an ordinary $(a_1 + 1)$-fold point and each tangent is different from $\{z = 0\}$.

Next we assume $i = 2$. Let z be a local parameter of Y_1 at $P_{2,j}$, $x' = X/Y - 1$, and $s' = S/T - \theta(1)$. Then, (x', z) gives a local parameter of $\mathbb{P}^1_{(X,Y)} \times Y_1$ at $Q_{2,j}$. Since $s' = v(z)z^{a_2+1}$ for some $v(z)$ with $v(0) \neq 0$, $u_1^*(D')$ is defined by

$$(x')^{a_2+1}(x' - \beta_1) \cdots (x' - \beta_{a_1}) - v(z)z^{a_2+1} = 0$$

around $Q_{2,j}$. Thus, we can see our assertion in this case because $\beta_1 \cdots \beta_{a_1} \neq 0$.

Finally we assume that $i = 3$. Let z be a local parameter of Y_1 at $P_{3,j}$, $y = Y/X$, and $t = T/S$. Since $t = v(z)z^{a_1+a_2+1}$ for some $v(z)$ with $v(0) \neq 0$, $u_1^*(D')$ is defined by

$$v(z)z^{a_1+a_2+1}(1 - \alpha_1 y) \cdots (1 - \alpha_{a_2} y) - y^{a_1+a_2+1} = 0$$

around $Q_{3,j}$. Thus, $Q_{3,j}$ is an ordinary $(a_1 + a_2 + 1)$-fold point and each tangent is different from $\{z = 0\}$ and $\{y = 0\}$. □

Let $\mu_1 : Z_1 \to \mathbb{P}^1_{(X,Y)} \times Y_1$ be blowing-ups at all points $Q_{i,j}$, and $E_{i,j}$ (-1)-curve over $Q_{i,j}$. Let $\overline{D}_1$ be the strict transform of $u_1^*(D)$ by μ_1. Then, by the previous claim, $\overline{D}_1$ is etale over Y_1 and

$$\overline{D}_1 \in \left| \mu_1^*(p_1^*(\mathcal{O}_{\mathbb{P}^1}(1)))^{\otimes a_1+a_2+2} \otimes \mu_1^*(q_1^*(h_1^*(\mathcal{O}_{\mathbb{P}^1}(1)))) \otimes \mathcal{O}_{Z_1}\left(-\sum_{i,j} c_i E_{i,j}\right) \right|.$$

Let F_j be the strict transform of the fiber $q_1^{-1}(P_{3,j})$. Note that $F_j \cap \overline{D}_1 = \emptyset$ for all j. Since F_j's are (-1)-curve, we can contract them. Let $\nu_1 : Z_1 \to X_1$ be the contraction of F_j's, and $f_1 : X_1 \to Y_1$ the induced morphism.

$$\begin{array}{ccccccc}
\mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)} & \xleftarrow{u_1} & \mathbb{P}^1_{(X,Y)} \times Y_1 & \xleftarrow{\mu_1} & Z_1 & \xrightarrow{\nu_1} & X_1 \\
\downarrow{\scriptstyle q} & & \downarrow{\scriptstyle q_1} & & \downarrow & & \downarrow{\scriptstyle f_1} \\
\mathbb{P}^1_{(S,T)} & \xleftarrow[h_1]{} & Y_1 & = & Y_1 & = & Y_1
\end{array}$$

Here we set $D_1 = (\nu_1)_*(\overline{D}_1)$, $L_1 = (\nu_1)_*(\mu_1^*(p_1^*(\mathcal{O}_{\mathbb{P}^1}(1))))^{**}$, and

$$M_1 = h_1^*(\mathcal{O}_{\mathbb{P}^1}(1)) \otimes \mathcal{O}_{Y_1}\left(-\sum_j c_3 P_{3,j}\right) \simeq \mathcal{O}_{Y_1}\left(-\sum_j P_{3,j}\right).$$

Then,

$$D_1 \in \left| L_1^{\otimes a_1+a_2+2} \otimes f_1^*(M_1) \otimes \mathcal{O}_{X_1}\left(-\sum_{\substack{i=1,2,\\ j\geq 1}} b_i \nu_1(E_{i,j})\right)\right|.$$

Therefore, by our construction of $f_1 : X_1 \to Y_1$, D_1, L_1 and M_1, it is easy to see all properties (1) — (6) in Lemma A.1. □

Proposition A.2. *Let g and a be integers with $g \geq 1$ and $0 \leq a \leq [g/2]$. Then, there is a smooth projective surface X over k, a smooth projective curve C over k, and a surjective morphism $f : X \to Y$ over k with the following properties.*

(1) *The generic fiber of f is a smooth hyperelliptic curve of genus g.*
(2) *f is not smooth and every fiber is reduced. Moreover, every singular fiber of f is a nodal curve consisting of a smooth curve of genus a and a smooth curve of genus $g-a$ joined at one point.*

Proof. Applying Lemma A.1 to a case where $a_1 = 2a$ and $a_2 = 2g-2a$, we fix a conic fibration as in Lemma A.1. Adding one point to Σ_1, if necessarily, we can take an effective and reduced divisor d on Y_1 such that $\Sigma_1 \subseteq \mathrm{Supp}(d)$ and $\deg(d)$ is even. Thus, there is a line bundle ϑ on Y_1 with $\mathcal{O}_{Y_1}(d) \simeq \vartheta^{\otimes 2}$, which produces a double covering $h_2 : Y \to Y_1$ of smooth projective curves such that h_2 is ramified over Σ_1. Let $\mu_2 : X_2 \to X_1 \times_{Y_1} Y$ be the minimal resolution of singularities of $X_1 \times_{Y_1} Y$. We set the induced morphisms as follows.

$$\begin{array}{ccc} X_1 & \xleftarrow{u_2} & X_2 \\ {\scriptstyle f_1}\downarrow & & \downarrow{\scriptstyle f_2} \\ Y_1 & \xleftarrow{h_2} & Y \end{array}$$

Let Σ_2 be the set of all critical values of f_2. Here, for all $Q \in \Sigma_2$, $f_2^{-1}(Q)$ is reduced, and there is an irreducible decomposition $f_2^{-1}(Q) = \overline{E}_Q^1 + \overline{E}_Q^2 + B_Q$ such that $u_2(\overline{E}_Q^i) = E_{h_2(Q)}^i$ for $i = 1, 2$ and B_Q is a (-2)-curves. We set $D_2 = u_2^*(D_1)$ and $B = \sum_{Q\in\Sigma_2} B_Q$. Then, D_2 is etale over Y and $D_2 + B$ is smooth over k because $D_2 \cap B = \emptyset$. Moreover,

$$D_2 \in \left| u_2^*(L_1)^{\otimes 2g+2} \otimes f_2^*(h_2^*(M_1)) \otimes \mathcal{O}_{X_2}\left(-u_2^*\left(\sum_{P\in\Sigma_1}(a_{\sigma(P)}+1)E_P^{\sigma(P)}\right)\right)\right|.$$

Let $\sigma_2 : \Sigma_2 \to \{1, 2\}$ be a map given by $\sigma_2 = \sigma \cdot h_2$. Then

$$u_2^*\left(\sum_{P\in\Sigma_1}(a_{\sigma(P)}+1)E_P^{\sigma(P)}\right) = \sum_{Q\in\Sigma_2}(a_{\sigma_2(Q)}+1)(2\overline{E}_Q^{\sigma_2(Q)} + B_Q).$$

Therefore,

$$D_2 + B \in \left| u_2^*(L_1)^{\otimes 2g+2} \otimes f_2^*(h_2^*(M_1)) \otimes \mathcal{O}_{X_2}\left(- \sum_{Q \in \Sigma_2} (2(a_{\sigma_2(Q)}+1)\overline{E}_Q^{\sigma_2(Q)} + a_{\sigma_2(Q)} B_Q) \right) \right|.$$

Here, since $\deg(h_2^*(M_1)) = 2\deg(M_1)$, $h_2^*(M_1)$ is divisible by 2 in $\mathrm{Pic}(Y)$. Further, a_i is even for each $i = 1, 2$. Thus,

$$u_2^*(L_1)^{\otimes 2g+2} \otimes f_2^*(h_2^*(M_1)) \otimes \mathcal{O}_{X_2}\left(- \sum_{Q \in \Sigma_2} (2(a_{\sigma_2(Q)}+1)\overline{E}_Q^{\sigma_2(Q)} + a_{\sigma_2(Q)} B_Q) \right)$$

is divisible by 2 in $\mathrm{Pic}(X_2)$, i.e., there is a line bundle H on X_2 with

$$H^{\otimes 2} \simeq u_2^*(L_1)^{\otimes 2g+2} \otimes f_2^*(h_2^*(M_1)) \otimes \mathcal{O}_{X_2}\left(- \sum_{Q \in \Sigma_2} (2(a_{\sigma_2(Q)}+1)\overline{E}_Q^{\sigma_2(Q)} + a_{\sigma_2(Q)} B_Q) \right).$$

Hence, we can construct a double covering $\mu_3 : X_3 \to X_2$ of smooth projective surfaces such that μ_3 is ramified over $D_2 + B$. Let $f_3 : X_3 \to Y$ be the induced morphism. Then, there is an irreducible decomposition

$$f^{-1}(Q) = \overline{C}_Q^1 + \overline{C}_Q^2 + 2\overline{B}_Q$$

as cycles such that $\mu_3(\overline{C}_Q^i) = \overline{E}_Q^i$ $(i = 1, 2)$ and $\mu_3(\overline{B}_Q) = B_Q$. Here it is easy to check that $\overline{B}_Q$ is a (-1)-curve. Thus, we have a contraction $\nu_3 : X_3 \to X$ of $\overline{B}_Q$'s, and an induced morphism $f : X \to Y$.

$$\begin{array}{ccccccc} X_1 & \xleftarrow{u_2} & X_2 & \xleftarrow{\mu_3} & X_3 & \xrightarrow{\nu_3} & X \\ {\scriptstyle f_1}\downarrow & & {\scriptstyle f_2}\downarrow & & {\scriptstyle f_3}\downarrow & & {\scriptstyle f}\downarrow \\ Y_1 & \xleftarrow{h_2} & Y & = & Y & = & Y \end{array}$$

We denote $\nu_3(\overline{C}_Q^i)$ by C_Q^i. Then, C_Q^1 (resp. C_Q^2) is a smooth projective curve of genus a (resp. $g - a$), $(C_Q^1 \cdot C_Q^2) = 1$, and $f^{-1}(Q) = C_Q^1 + C_Q^2$. Thus, $f : X \to Y$ is our desired fibration. □

In the same way, we can also show the following proposition.

Proposition A.3. *Let g and a be integers with $g \geq 1$ and $0 \leq a \leq [(g-1)/2]$. Then, there is a smooth projective surface X over k, a smooth projective curve C over k, and a surjective morphism $f : X \to Y$ over k with the following properties.*

(1) *The generic fiber of f is a smooth hyperelliptic curve of genus g.*
(2) *f is not smooth and every fiber is reduced. Moreover, every singular fiber of f is a nodal curve consisting of a smooth curve of genus a and a smooth curve of genus $g - a - 1$ joined at two points.*

Proof. Applying Lemma A.1 to a case where $a_1 = 2a + 1$ and $a_2 = 2g - 2a - 1$, we fix a conic fibration as in Lemma A.1. In this case, $\deg(M_1)$ is even. Thus,

$$L_1^{\otimes 2g+2} \otimes f_1^*(M_1) \otimes \mathcal{O}_{X_1}\left(- \sum_{P \in \Sigma_1} (a_{\sigma(P)}+1) E_P^{\sigma(P)} \right)$$

is divisible by 2 in $\mathrm{Pic}(X_1)$. Therefore, there is a double covering $\mu : X \to X_1$ of smooth projective surfaces such that μ_3 is ramified over D_1. Then, the induced morphism $f : X \to Y_1$ is the desired fibration. □

References

[1] J.-B. Bost, Semi-stability and height of cycles, Invent. Math., 118 (1994), 223–253.
[2] M. Cornalba and J. Harris, Divisor classes associated to families of stable varieties, with application to the moduli space of curves, Ann. Scient. Ec. Norm. Sup., 21 (1988), 455–475.
[3] P. Deligne and D. Mumford, The irreducibility of the space of curves of given genus, Publ. Math. IHES, 36 (1969), 75–110.
[4] W. Fulton, Intersection Theory, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Band 2, (1984), Springer-Verlag.
[5] D. Gieseker, Stable vector bundles and the Frobenius morphism, Ann. Scient. Ec. Norm. Sup., 6(1973) 95-101.
[6] D. Gieseker, On a theorem of Bogomolov on Chern classes of stable bundles, Amer. J. Math., 101 (1979), 79-85.
[7] R. Hartshorne, Ample vector bundles on curves, Nagoya Math. J. 43 (1971) 73-90.
[8] F. Knudsen and D. Mumford, The projectivity of the moduli space of stable curves, I: Preliminaries on "det" and "Div", Math. Scand. 39 (1976), 19-55.
[9] M. Maruyama, Openness of a family of torsion free sheaves, J. Math. Kyoto Univ., 16 (1976), 627-637.
[10] M. Maruyama, The theorem of Grauer-Mulich-Spindler, Math. Ann. 255 (1981) 317-333.
[11] Y. Miyaoka, The Chern Classes and Kodaira Dimension of a Minimal Variety, Advanced Studies in Pure Mathematics 10, 1987, Algebraic Geometry, Sendai, 1985, 449-476, Kinokuniya.
[12] A. Moriwaki, Arithmetic Bogomolov-Gieseker's inequality, Amer. J. Math. 117, 1325-1347 (1995).
[13] A. Moriwaki, Bogomolov conjecture over function fields for stable curves with only irreducible fibers, to appear in Comp. Math..
[14] A. Moriwaki, Bogomolov conjecture for curves of genus 2 over function fields, to appear in J. Math. Kyoto Univ..
[15] A. Moriwaki, A sharp slope inequality for general stable fibrations of curves, J. reine angew. Math. 480 (1996), 177–195.
[16] A. Paranjape and S. Ramanan, On the canonical ring of an algebraic curve, Algebraic Geometry and Commutative Algebra in Honor of Masayoshi NAGATA, vol. II (1987), 501–516.
[17] E, Ullmo, Positivité et discretion des points algébriques des courbes, preprint.
[18] G. Xiao, Fibered algebraic surfaces with low slope, Math. Ann., 276 (1987), 449–466.
[19] S. Zhang, Admissible pairing on a curve, Invent. Math., 112 (1993), 171–193.

Department of Mathematics, Faculty of Science, Kyoto University, Kyoto, 606-01, Japan
E-mail address: moriwaki@kusm.kyoto-u.ac.jp