

# Moduli of Lagrangian immersions in pair-of-pants decompositions and mirror symmetry

Siu-Cheong Lau  
Boston University

Decemeber 2017

Joint work with Cheol-Hyun Cho and Hansol Hong

# Outline

- ▶ Overview.
- ▶ Construction for pair-of-pants decompositions.
- ▶ An application to wall-crossing.

# Section 1

## Overview and background

# Moduli theory in the B-side

- ▶ Moduli theory for vector bundles was well established.

## Theorem (**Donaldson, Uhlenbeck-Yau**)

*A slope-semistable holomorphic vector bundle admits a Hermitian Yang-Mills metric.*

- ▶ GIT and stability conditions were essential to the construction.
- ▶ **Bridgeland** developed a general mathematical theory of stability conditions for triangulated categories.
- ▶ **Toda** developed foundational techniques to construct Bridgeland stability conditions for derived categories of coherent sheaves.
- ▶ Moduli spaces undergo birational changes (such as flops) in a variation of stability conditions.
- ▶ **How about moduli of Lagrangians in the mirror A-side?**

# Moduli theory in the B-side

- ▶ Moduli theory for vector bundles was well established.

## Theorem (**Donaldson, Uhlenbeck-Yau**)

*A slope-semistable holomorphic vector bundle admits a Hermitian Yang-Mills metric.*

- ▶ GIT and stability conditions were essential to the construction.
- ▶ **Bridgeland** developed a general mathematical theory of stability conditions for triangulated categories.
- ▶ **Toda** developed foundational techniques to construct Bridgeland stability conditions for derived categories of coherent sheaves.
- ▶ Moduli spaces undergo birational changes (such as flops) in a variation of stability conditions.
- ▶ How about moduli of Lagrangians in the mirror A-side?

# Moduli theory in the B-side

- ▶ Moduli theory for vector bundles was well established.

## Theorem (**Donaldson, Uhlenbeck-Yau**)

*A slope-semistable holomorphic vector bundle admits a Hermitian Yang-Mills metric.*

- ▶ GIT and stability conditions were essential to the construction.
- ▶ **Bridgeland** developed a general mathematical theory of stability conditions for triangulated categories.
- ▶ **Toda** developed foundational techniques to construct Bridgeland stability conditions for derived categories of coherent sheaves.
- ▶ Moduli spaces undergo birational changes (such as flops) in a variation of stability conditions.
- ▶ **How about moduli of Lagrangians in the mirror A-side?**

# Ingredients for moduli theory of Lagrangians

- ▶ *Complexification.* The naive moduli spaces are affine manifolds with singularities. Complexification is needed in order to compactify. Technically we need to work over the Novikov ring.
- ▶ *Quantum correction.* The canonical complex structures need to be corrected using Lagrangian Floer theory [Fukaya-Oh-Ohta-Ono]. The combinatorial structure of quantum corrections for SYZ fibrations was deeply studied by **Kontsevich-Soibelman** and **Gross-Siebert**.
- ▶ *Landau-Ginzburg model.* The moduli in general are singular varieties. They are described as critical loci of holomorphic functions.
- ▶ *Singular Lagrangians.* **Kontsevich** proposed to study them using cosheaves of categories. **Nadler** is developing a theory of arboreal singularities.

# Ingredients for moduli theory of Lagrangians

- ▶ *Complexification.* The naive moduli spaces are affine manifolds with singularities. Complexification is needed in order to compactify. Technically we need to work over the Novikov ring.
- ▶ *Quantum correction.* The canonical complex structures need to be corrected using Lagrangian Floer theory [**Fukaya-Oh-Ohta-Ono**]. The combinatorial structure of quantum corrections for SYZ fibrations was deeply studied by **Kontsevich-Soibelman** and **Gross-Siebert**.
- ▶ *Landau-Ginzburg model.* The moduli in general are singular varieties. They are described as critical loci of holomorphic functions.
- ▶ *Singular Lagrangians.* **Kontsevich** proposed to study them using cosheaves of categories. **Nadler** is developing a theory of arboreal singularities.




# Ingredients for moduli theory of Lagrangians

- ▶ *Complexification.* The naive moduli spaces are affine manifolds with singularities. Complexification is needed in order to compactify. Technically we need to work over the Novikov ring.
- ▶ *Quantum correction.* The canonical complex structures need to be corrected using Lagrangian Floer theory [**Fukaya-Oh-Ohta-Ono**]. The combinatorial structure of quantum corrections for SYZ fibrations was deeply studied by **Kontsevich-Soibelman** and **Gross-Siebert**.
- ▶ *Landau-Ginzburg model.* The moduli in general are singular varieties. They are described as critical loci of holomorphic functions.
- ▶ *Singular Lagrangians.* **Kontsevich** proposed to study them using cosheaves of categories. **Nadler** is developing a theory of arboreal singularities.

# Ingredients for moduli theory of Lagrangians


- ▶ *Complexification.* The naive moduli spaces are affine manifolds with singularities. Complexification is needed in order to compactify. Technically we need to work over the Novikov ring.
- ▶ *Quantum correction.* The canonical complex structures need to be corrected using Lagrangian Floer theory [**Fukaya-Oh-Ohta-Ono**]. The combinatorial structure of quantum corrections for SYZ fibrations was deeply studied by **Kontsevich-Soibelman** and **Gross-Siebert**.
- ▶ *Landau-Ginzburg model.* The moduli in general are singular varieties. They are described as critical loci of holomorphic functions.
- ▶ *Singular Lagrangians.* **Kontsevich** proposed to study them using cosheaves of categories. **Nadler** is developing a theory of arboreal singularities.

# Why care about Lagrangian immersions




- ▶ Lagrangian immersions have a well-defined Floer theory by **Akaho-Joyce**.
- ▶ They are the main sources of wall-crossing phenomenon in the SYZ setting.
- ▶ The deformation space of a Lagrangian immersion is 'bigger' than its smoothing and covers a local family of Lagrangians (including singular Lagrangians).
- ▶ A Lagrangian immersion in the pair-of-pants was used by **Seidel** and **Sheridan** to prove homological mirror symmetry for Fermat-type hypersurfaces.

# Why care about Lagrangian immersions




- ▶ Lagrangian immersions have a well-defined Floer theory by **Akaho-Joyce**.
- ▶ They are the main sources of wall-crossing phenomenon in the SYZ setting.
- ▶ The deformation space of a Lagrangian immersion is 'bigger' than its smoothing and covers a local family of Lagrangians (including singular Lagrangians).
- ▶ A Lagrangian immersion in the pair-of-pants was used by **Seidel** and **Sheridan** to prove homological mirror symmetry for Fermat-type hypersurfaces.

# Why care about Lagrangian immersions



- ▶ Lagrangian immersions have a well-defined Floer theory by **Akaho-Joyce**.
- ▶ They are the main sources of wall-crossing phenomenon in the SYZ setting.
- ▶ The deformation space of a Lagrangian immersion is 'bigger' than its smoothing and covers a local family of Lagrangians (including singular Lagrangians).
- ▶ A Lagrangian immersion in the pair-of-pants was used by **Seidel** and **Sheridan** to prove homological mirror symmetry for Fermat-type hypersurfaces.

# Why care about Lagrangian immersions



- ▶ Lagrangian immersions have a well-defined Floer theory by **Akaho-Joyce**.
- ▶ They are the main sources of wall-crossing phenomenon in the SYZ setting.
- ▶ The deformation space of a Lagrangian immersion is 'bigger' than its smoothing and covers a local family of Lagrangians (including singular Lagrangians).
- ▶ A Lagrangian immersion in the pair-of-pants was used by **Seidel** and **Sheridan** to prove homological mirror symmetry for Fermat-type hypersurfaces.

## Family Floer theory

- ▶ **Fukaya** proposed to study mirror symmetry by using  $CF(L_b, \cdot)$  for fibers  $L_b$  of a Lagrangian torus fibration.
- ▶ **Tu** took this approach to construct mirror spaces away from singular fibers.
- ▶ **Abouzaid** constructed family Floer functors for torus bundles and showed that the functor is fully faithful.
- ▶ We consider moduli of Lagrangians which are not necessarily tori.
- ▶ Instead of Fukaya trick, **we consider pseudo-isomorphisms between Lagrangian immersions which intersect with each other in a certain manner, and obtain the gluing information from cocycle conditions.** (In particular we do not need diffeomorphisms.)
- ▶ Lagrangian immersions help in gluing. The immersed sectors provide  $\Lambda_+$  (or even  $\Lambda_0$ ) space of deformations, which is much 'bigger' than the deformation space of flat  $\mathbf{C}^\times$ -connections.

## Family Floer theory


- ▶ **Fukaya** proposed to study mirror symmetry by using  $CF(L_b, \cdot)$  for fibers  $L_b$  of a Lagrangian torus fibration.
- ▶ **Tu** took this approach to construct mirror spaces away from singular fibers.
- ▶ **Abouzaid** constructed family Floer functors for torus bundles and showed that the functor is fully faithful.
- ▶ We consider moduli of Lagrangians which are not necessarily tori.
- ▶ Instead of Fukaya trick, **we consider pseudo-isomorphisms between Lagrangian immersions which intersect with each other in a certain manner, and obtain the gluing information from cocycle conditions.** (In particular we do not need diffeomorphisms.)
- ▶ Lagrangian immersions help in gluing. The immersed sectors provide  $\Lambda_+$  (or even  $\Lambda_0$ ) space of deformations, which is much 'bigger' than the deformation space of flat  $\mathbf{C}^\times$ -connections.



## Section 2


### Pair-of-pants decompositions

# Punctured Riemann surfaces




- ▶ Consider a pair-of-pants decomposition of a punctured Riemann surface.
- ▶ Homological mirror symmetry was proved for punctured Riemann surface by **Abouzaid-Auroux-Efimov-Katzarkov-Orlov, Bocklandt, and Heather Lee**.
- ▶ Our focus is on the construction of moduli rather than HMS. The construction comes with a natural functor which derives HMS.
- ▶ We have a family of Lagrangian immersions as shown in the figure. **We will take finitely many representing Lagrangians and glue their deformation spaces together.**
- ▶  $\dim = 1$  contains the essential ingredients. In higher dimensions, Seidel's Lagrangians are replaced by Sheridan's Lagrangians.

# Punctured Riemann surfaces




- ▶ Consider a pair-of-pants decomposition of a punctured Riemann surface.
- ▶ Homological mirror symmetry was proved for punctured Riemann surface by **Abouzaid-Auroux-Efimov-Katzarkov-Orlov, Bocklandt, and Heather Lee**.
- ▶ Our focus is on the construction of moduli rather than HMS. The construction comes with a natural functor which derives HMS.
- ▶ We have a family of Lagrangian immersions as shown in the figure. **We will take finitely many representing Lagrangians and glue their deformation spaces together.**
- ▶  $\dim = 1$  contains the essential ingredients. In higher dimensions, Seidel's Lagrangians are replaced by Sheridan's Lagrangians.

# Review on local moduli construction




- ▶ First consider a pair-of-pants, with the Lagrangian immersion constructed by Seidel.
- ▶  $CF(L, L) = \text{Span}\{\mathbf{1}, X, Y, Z, \bar{X}, \bar{Y}, \bar{Z}, \text{pt}\}$ .
- ▶ We proved that  $xX + yY + zZ$  is weakly unobstructed for  $x, y, z \in \mathbf{C}$ . It is important that the areas of the two triangles are the same.
- ▶ Thus the local moduli is  $(\mathbf{C}^3, W)$ , where  $W = xyz$ .
- ▶ The pair-of-pants can be compactified to  $\mathbb{P}_{a,b,c}^1$ . We used this to construct and compute the mirror, and derived homological mirror symmetry. In an ongoing work with **Amorim** we prove closed-string mirror symmetry. For elliptic orbifolds the coefficients of  $W$  are modular forms [L.-Zhou].

# Review on local moduli construction



- ▶ First consider a pair-of-pants, with the Lagrangian immersion constructed by Seidel.
- ▶  $CF(L, L) = \text{Span}\{\mathbf{1}, X, Y, Z, \bar{X}, \bar{Y}, \bar{Z}, \text{pt}\}$ .
- ▶ We proved that  $xX + yY + zZ$  is weakly unobstructed for  $x, y, z \in \mathbf{C}$ . It is important that the areas of the two triangles are the same.
- ▶ Thus the local moduli is  $(\mathbf{C}^3, W)$ , where  $W = xyz$ .
- ▶ The pair-of-pants can be compactified to  $\mathbb{P}_{a,b,c}^1$ . We used this to construct and compute the mirror, and derived homological mirror symmetry. In an ongoing work with **Amorim** we prove closed-string mirror symmetry. For elliptic orbifolds the coefficients of  $W$  are modular forms [**L.-Zhou**].


# Gluing



- ▶ Consider the four-punctured sphere as shown above.
- ▶ We need to glue the deformation spaces of the two Seidel Lagrangians  $S_1$  and  $S_2$ .
- ▶ There are two main processes: smoothing and gauge change.
- ▶ Let's pretend to work over  $\mathbf{C}$  at this stage. The gluing we need is

$$(\mathbf{C}^3, W) \xleftrightarrow{\text{smoothing}} (\mathbf{C}^\times \times \mathbf{C}^2) \xleftrightarrow{\text{gauge change}} (\mathbf{C}^\times \times \mathbf{C}^2) \xleftrightarrow{\text{smoothing}} \mathbf{C}^3.$$


# Gluing



- ▶ Consider the four-punctured sphere as shown above.
- ▶ We need to glue the deformation spaces of the two Seidel Lagrangians  $S_1$  and  $S_2$ .
- ▶ There are two main processes: smoothing and gauge change.
- ▶ Let's pretend to work over  $\mathbf{C}$  at this stage. The gluing we need is

$$(\mathbf{C}^3, W) \xleftrightarrow{\text{smoothing}} (\mathbf{C}^\times \times \mathbf{C}^2) \xleftrightarrow{\text{gauge change}} (\mathbf{C}^\times \times \mathbf{C}^2) \xleftrightarrow{\text{smoothing}} \mathbf{C}^3.$$


# Choice of gauge change



- ▶ There is a vanishing sphere in a smoothing at an immersed point of  $S$ . In this case it is simply the union of two points.
- ▶ Put a flat  $\mathbf{C}^\times$  connection on the smoothing  $C$ , which is acting by  $t \in \mathbf{C}^\times$  when passing through the two points.
- ▶ The position of the two points are different for smoothings on the left and on the right.
- ▶ When a gauge point  $T$  is moved across the immersed point  $Y$ , the  $A_\infty$  algebras are related by  $\tilde{y} = ty$ .
- ▶ There are different ways of moving the gauge points to match them. This results in  $\tilde{y} = t^a y$ ,  $\tilde{z} = t^b z$  with  $a + b = 2$ .




# Choice of gauge change




- ▶ There is a vanishing sphere in a smoothing at an immersed point of  $S$ . In this case it is simply the union of two points.
- ▶ Put a flat  $\mathbf{C}^\times$  connection on the smoothing  $C$ , which is acting by  $t \in \mathbf{C}^\times$  when passing through the two points.
- ▶ The position of the two points are different for smoothings on the left and on the right.
- ▶ When a gauge point  $T$  is moved across the immersed point  $Y$ , the  $A_\infty$  algebras are related by  $\tilde{y} = ty$ .
- ▶ There are different ways of moving the gauge points to match them. This results in  $\tilde{y} = t^a y$ ,  $\tilde{z} = t^b z$  with  $a + b = 2$ .

## Choice of gauge change



- ▶ There is a vanishing sphere in a smoothing at an immersed point of  $S$ . In this case it is simply the union of two points.
- ▶ Put a flat  $\mathbf{C}^\times$  connection on the smoothing  $C$ , which is acting by  $t \in \mathbf{C}^\times$  when passing through the two points.
- ▶ The position of the two points are different for smoothings on the left and on the right.
- ▶ When a gauge point  $T$  is moved across the immersed point  $Y$ , the  $A_\infty$  algebras are related by  $\tilde{y} = ty$ .
- ▶ There are different ways of moving the gauge points to match them. This results in  $\tilde{y} = t^a y$ ,  $\tilde{z} = t^b z$  with  $a + b = 2$ .

# Choice of gauge change




- ▶ There is a vanishing sphere in a smoothing at an immersed point of  $S$ . In this case it is simply the union of two points.
- ▶ Put a flat  $\mathbf{C}^\times$  connection on the smoothing  $C$ , which is acting by  $t \in \mathbf{C}^\times$  when passing through the two points.
- ▶ The position of the two points are different for smoothings on the left and on the right.
- ▶ When a gauge point  $T$  is moved across the immersed point  $Y$ , the  $A_\infty$  algebras are related by  $\tilde{y} = ty$ .
- ▶ There are different ways of moving the gauge points to match them. This results in  $\tilde{y} = t^a y$ ,  $\tilde{z} = t^b z$  with  $a + b = 2$ .

# Smoothing

- ▶ We need to glue the deformation spaces of  $S$  and  $C$ .
- ▶ They are given by  $(x, y, z) \in \mathbf{C}^3$  and  $(t, y_0, z_0) \in \mathbf{C}^\times \times \mathbf{C}^2$  respectively.
- ▶ Intuitively to match the superpotentials  $xyz$  and  $ty_0z_0$ , we simply put  $x = t, y = y_0, z = z_0$ .
- ▶ We take  $S_1$  to be the deformed Seidel Lagrangian which intersects  $C$  as in the figure.
- ▶ Then we use cocycle conditions to deduce the gluing between  $S$  and  $C$ .


# Smoothing

- ▶ We need to glue the deformation spaces of  $S$  and  $C$ .
- ▶ They are given by  $(x, y, z) \in \mathbf{C}^3$  and  $(t, y_0, z_0) \in \mathbf{C}^\times \times \mathbf{C}^2$  respectively.
- ▶ Intuitively to match the superpotentials  $xyz$  and  $ty_0z_0$ , we simply put  $x = t, y = y_0, z = z_0$ .
- ▶ We take  $S_1$  to be the deformed Seidel Lagrangian which intersects  $C$  as in the figure.
- ▶ Then we use cocycle conditions to deduce the gluing between  $S$  and  $C$ .




# Isomorphisms in smoothing

- ▶ We consider  $a_1 + b_1 \in CF((C, \nabla_t), (S, xX))$  and  $c_2 + d_2 \in CF((S, xX), (C, \nabla_t))$ .
- ▶ Consider cocycle conditions on  $a_1 + b_1$  and  $c_2 + d_2$ . Once the cocycle conditions are satisfied, they give isomorphisms between the two objects.




## A paradox



- ▶  $S$  is isomorphic to  $S_1$ , and  $(S_1, xX)$  is isomorphic to  $(C, \nabla_t)$  for  $t = x \neq 0$ . Thus  $(S, xX)$  is isomorphic to  $C$ .
- ▶ But  $S$  and  $C$  are disjoint, and hence there is no morphism between them!
- ▶ We need to take a closer look at areas of holomorphic discs.
- ▶ Following **Fukaya-Oh-Ohta-Ono**, we shall use the Novikov ring  $\Lambda_0 = \{\sum_{i=0}^{\infty} a_i T^{A_i} : 0 \leq A_0 \leq A_1 \leq \dots\}$  to filter deformations into different energy levels.


## A paradox



- ▶  $S$  is isomorphic to  $S_1$ , and  $(S_1, xX)$  is isomorphic to  $(C, \nabla_t)$  for  $t = x \neq 0$ . Thus  $(S, xX)$  is isomorphic to  $C$ .
- ▶ But  $S$  and  $C$  are disjoint, and hence there is no morphism between them!
- ▶ We need to take a closer look at areas of holomorphic discs.
- ▶ Following **Fukaya-Oh-Ohta-Ono**, we shall use the Novikov ring  $\Lambda_0 = \{\sum_{i=0}^{\infty} a_i T^{A_i} : 0 \leq A_0 \leq A_1 \leq \dots\}$  to filter deformations into different energy levels.




## Area constraints for isomorphisms




- ▶ For the cocycle conditions on  $(S_1, x_1 X_1) \rightarrow (C, \nabla_t)$ , indeed we have  $t = T^{A_1 + \dots + A_5 - A_7} x_1$ .
- ▶ Note that  $t \in \mathbf{C}^\times$  while  $x_1 \in \Lambda_0$ . When  $A_1 + \dots + A_5 > A_7$ , the condition is never satisfied.
- ▶ Hence  $(S_1, x_1 X_1)$  can never be isomorphic to  $(C, \nabla_t)$  if  $A_1 + \dots + A_5 > A_7$ .
- ▶ When  $A_1 + \dots + A_5 = A_7$ , the change of coordinates  $x_1 = t$  does not involve Novikov parameters.
- ▶ For the cocycle conditions on  $(S, xX) \rightarrow (S_1, x_1 X_1)$ , we have  $x_1 = T^A x$  for certain  $A > 0$ . Thus  $\text{val}(x_1) \geq A > 0$  in order to have them to be isomorphic.
- ▶ The two regions  $\text{val}(x_1) = 0$  and  $\text{val}(x_1) \geq A > 0$  are disjoint. This solves the paradox.

## Area constraints for isomorphisms



- ▶ For the cocycle conditions on  $(S_1, x_1 X_1) \rightarrow (C, \nabla_t)$ , indeed we have  $t = T^{A_1 + \dots + A_5 - A_7} x_1$ .
- ▶ Note that  $t \in \mathbf{C}^\times$  while  $x_1 \in \Lambda_0$ . When  $A_1 + \dots + A_5 > A_7$ , the condition is never satisfied.
- ▶ Hence  $(S_1, x_1 X_1)$  can never be isomorphic to  $(C, \nabla_t)$  if  $A_1 + \dots + A_5 > A_7$ .
- ▶ When  $A_1 + \dots + A_5 = A_7$ , the change of coordinates  $x_1 = t$  does not involve Novikov parameters.
- ▶ For the cocycle conditions on  $(S, xX) \rightarrow (S_1, x_1 X_1)$ , we have  $x_1 = T^A x$  for certain  $A > 0$ . Thus  $\text{val}(x_1) \geq A > 0$  in order to have them to be isomorphic.
- ▶ The two regions  $\text{val}(x_1) = 0$  and  $\text{val}(x_1) \geq A > 0$  are disjoint. This solves the paradox.

## Area constraints for isomorphisms




- ▶ For the cocycle conditions on  $(S_1, x_1 X_1) \rightarrow (C, \nabla_t)$ , indeed we have  $t = T^{A_1 + \dots + A_5 - A_7} x_1$ .
- ▶ Note that  $t \in \mathbf{C}^\times$  while  $x_1 \in \Lambda_0$ . When  $A_1 + \dots + A_5 > A_7$ , the condition is never satisfied.
- ▶ Hence  $(S_1, x_1 X_1)$  can never be isomorphic to  $(C, \nabla_t)$  if  $A_1 + \dots + A_5 > A_7$ .
- ▶ When  $A_1 + \dots + A_5 = A_7$ , the change of coordinates  $x_1 = t$  does not involve Novikov parameters.
- ▶ For the cocycle conditions on  $(S, xX) \rightarrow (S_1, x_1 X_1)$ , we have  $x_1 = T^A x$  for certain  $A > 0$ . Thus  $\text{val}(x_1) \geq A > 0$  in order to have them to be isomorphic.
- ▶ The two regions  $\text{val}(x_1) = 0$  and  $\text{val}(x_1) \geq A > 0$  are disjoint. This solves the paradox.

## More pair-of-pants


- ▶ For the four-punctured sphere with two pants in the decomposition, we can choose  $S_1$  as above in each pants such that the gluing  $x = t$  does not involve the Novikov parameter.
- ▶ Then the  $\mathbf{C}$ -valued mirror  $(\mathcal{O}(-1) \oplus \mathcal{O}(-1), W)$  embeds well in the  $\Lambda$ -valued mirror  $(\Lambda_0^3 \cup \Lambda_0^3, W)$ .
- ▶ However suppose there is one more pants in the decomposition. Then we need to consider another Seidel Lagrangian  $S'_1$ .
- ▶ The gluing between  $S_1$  and  $S'_1$  is  $x_1 = T^A x'_1$  for certain  $A > 0$ . Unavoidably the gluing involves the Novikov parameter.

## More pair-of-pants




- ▶ For the four-punctured sphere with two pants in the decomposition, we can choose  $S_1$  as above in each pants such that the gluing  $x = t$  does not involve the Novikov parameter.
- ▶ Then the  $\mathbf{C}$ -valued mirror  $(\mathcal{O}(-1) \oplus \mathcal{O}(-1), W)$  embeds well in the  $\Lambda$ -valued mirror  $(\Lambda_0^3 \cup \Lambda_0^3, W)$ .
- ▶ However suppose there is one more pants in the decomposition. Then we need to consider another Seidel Lagrangian  $S_1'$ .
- ▶ The gluing between  $S_1$  and  $S_1'$  is  $x_1 = T^A x_1'$  for certain  $A > 0$ . Unavoidably the gluing involves the Novikov parameter.

## More pair-of-pants




- ▶ For the four-punctured sphere with two pants in the decomposition, we can choose  $S_1$  as above in each pants such that the gluing  $x = t$  does not involve the Novikov parameter.
- ▶ Then the  $\mathbf{C}$ -valued mirror  $(\mathcal{O}(-1) \oplus \mathcal{O}(-1), W)$  embeds well in the  $\Lambda$ -valued mirror  $(\Lambda_0^3 \cup \Lambda_0^3, W)$ .
- ▶ However suppose there is one more pants in the decomposition. Then we need to consider another Seidel Lagrangian  $S_1'$ .
- ▶ The gluing between  $S_1$  and  $S_1'$  is  $x_1 = T^A x_1'$  for certain  $A > 0$ . Unavoidably the gluing involves the Novikov parameter.

# Moduli over $\Lambda$



- ▶ The moduli over  $\Lambda$  is depicted above.
- ▶ The  $\mathbf{C}$ -valued toric Calabi-Yau is not embedded, since the gluing  $x_1 = T^A x_1'$  does not preserve  $\mathbf{C}$ .
- ▶ On the other hand, the  $\mathbf{C}$ -valued critical locus of  $W$  is embedded in the  $\Lambda$ -valued moduli. Exact Lagrangians are transformed to matrix factorizations supported on the  $\mathbf{C}$ -valued critical locus.



# A compactification



- ▶ The gluing  $x = t$ ,  $y = y_0$ ,  $z = z_0$  looks pretty trivial. Let's compactify to get more interesting gluing.
- ▶ As an example, compactify the four-punctured sphere to a sphere. we still consider the moduli of double-circles.
- ▶ The gluing is further quantum-corrected by discs emanated from infinite divisors.
- ▶ The new gluing is  $t = x$ ,  $y_0 = y_1 + t_1^{-1}$ ,  $z_0 = z_1 - t_1^{-1}$ .  
 $W = xyz - y_1 + x_1 + z_1$ . (The Novikov parameter is suppressed for simplicity.)
- ▶ Unlike the case for anti-canonical divisors, gluing needs to be corrected upon compactification.





# A compactification




- ▶ The gluing  $x = t, y = y_0, z = z_0$  looks pretty trivial. Let's compactify to get more interesting gluing.
- ▶ As an example, compactify the four-punctured sphere to a sphere. we still consider the moduli of double-circles.
- ▶ The gluing is further quantum-corrected by discs emanated from infinite divisors.
- ▶ The new gluing is  $t = x, y_0 = y_1 + t_1^{-1}, z_0 = z_1 - t_1^{-1}$ .  
 $W = xyz - y_1 + x_1 + z_1$ . (The Novikov parameter is suppressed for simplicity.)
- ▶ Unlike the case for anti-canonical divisors, gluing needs to be corrected upon compactification.

# A compactification




- ▶ The gluing  $x = t, y = y_0, z = z_0$  looks pretty trivial. Let's compactify to get more interesting gluing.
- ▶ As an example, compactify the four-punctured sphere to a sphere. we still consider the moduli of double-circles.
- ▶ The gluing is further quantum-corrected by discs emanated from infinite divisors.
- ▶ The new gluing is  $t = x, y_0 = y_1 + t_1^{-1}, z_0 = z_1 - t_1^{-1}$ .  
 $W = xyz - y_1 + x_1 + z_1$ . (The Novikov parameter is suppressed for simplicity.)
- ▶ Unlike the case for anti-canonical divisors, gluing needs to be corrected upon compactification.

## Changing stability and flop



- ▶ Let's go back to the four-punctured sphere. We can take another pair-of-pants decomposition.
- ▶ It corresponds to another choice of a quadratic differential (stability condition [**Haiden-Katzarkov-Kontsevich**]).
- ▶ This results in a flop of the moduli space  $(\mathcal{O}(-1) \oplus \mathcal{O}(-1), W)$ .
- ▶ We can also consider a quadratic differential with double zeros. Then the moduli is the non-commutative resolution of the conifold corresponding to a quiver ([**Cho-Hong-L. 15**]).
- ▶ In [**Fan-Hong-L.-Yau**] we studied a 3d version of this (deformed conifold).

## Changing stability and flop




- ▶ Let's go back to the four-punctured sphere. We can take another pair-of-pants decomposition.
- ▶ It corresponds to another choice of a quadratic differential (stability condition [**Haiden-Katzarkov-Kontsevich**]).
- ▶ This results in a flop of the moduli space  $(\mathcal{O}(-1) \oplus \mathcal{O}(-1), W)$ .
- ▶ We can also consider a quadratic differential with double zeros. Then the moduli is the non-commutative resolution of the conifold corresponding to a quiver ([**Cho-Hong-L. 15**]).
- ▶ In [**Fan-Hong-L.-Yau**] we studied a 3d version of this (deformed conifold).

## Section 3


### Wall-crossing

# Immersed sphere




- ▶ The two-dimensional immersed sphere leads to wall-crossing phenomena for SYZ Lagrangian fibrations.
- ▶ **Fukaya** studied this immersion and studied the relation with the mirror equation. Here we realize it from cocycle conditions.
- ▶ It has two degree-one immersed generators  $U$  and  $V$ . It gives the deformations  $b = uU + vV$  where  $(u, v) \in (\Lambda_0 \times \Lambda_+) \cup (\Lambda_+ \times \Lambda_0)$ .
- ▶ There are constant holomorphic discs with corners  $U, V, \dots, U, V$ . To ensure there are only finitely many terms under every energy level, we only allow one of  $u, v$  having valuation zero.

# Immersed sphere



- ▶ The two-dimensional immersed sphere leads to wall-crossing phenomena for SYZ Lagrangian fibrations.
- ▶ **Fukaya** studied this immersion and studied the relation with the mirror equation. Here we realize it from cocycle conditions.
- ▶ It has two degree-one immersed generators  $U$  and  $V$ . It gives the deformations  $b = uU + vV$  where  $(u, v) \in (\Lambda_0 \times \Lambda_+) \cup (\Lambda_+ \times \Lambda_0)$ .
- ▶ There are constant holomorphic discs with corners  $U, V, \dots, U, V$ . To ensure there are only finitely many terms under every energy level, we only allow one of  $u, v$  having valuation zero.


# Immersed sphere



- ▶ The two-dimensional immersed sphere leads to wall-crossing phenomena for SYZ Lagrangian fibrations.
- ▶ **Fukaya** studied this immersion and studied the relation with the mirror equation. Here we realize it from cocycle conditions.
- ▶ It has two degree-one immersed generators  $U$  and  $V$ . It gives the deformations  $b = uU + vV$  where  $(u, v) \in (\Lambda_0 \times \Lambda_+) \cup (\Lambda_+ \times \Lambda_0)$ .
- ▶ There are constant holomorphic discs with corners  $U, V, \dots, U, V$ . To ensure there are only finitely many terms under every energy level, we only allow one of  $u, v$  having valuation zero.




# Immersed sphere




- ▶ It has two degree-one immersed generators  $U$  and  $V$ . It gives the deformations  $b = uU + vV$  where  $(u, v) \in (\Lambda_0 \times \Lambda_+) \cup (\Lambda_+ \times \Lambda_0)$ .
- ▶ There are constant holomorphic discs with corners  $U, V, \dots, U, V$ . To ensure there are only finitely many terms under every energy level, we only allow one of  $u, v$  having valuation zero.
- ▶ The constant discs with corners  $U, V, \dots, U, V$  contributing to pt of  $m_0^b$  cancel with that with corners  $V, U, \dots, V, U$ . Thus  $b = uU + vV$  is weakly unobstructed.

# Gluing




- ▶ Consider the immersed sphere  $S$  and a Chekanov torus  $T$ .  $S$  is made by deforming the immersed fiber.
- ▶ We glue the deformations  $(u, v) \in \Lambda_0 \times \Lambda_+$  with the deformations  $(x, y) \in (\mathbf{C}^\times \oplus \Lambda_+)^2$  of  $T$ .

# Gluing




- ▶ We glue the deformations  $(u, v) \in \Lambda_0 \times \Lambda_+$  with the deformations  $(x, y) \in (\mathbf{C}^\times \oplus \Lambda_+)^2$  of  $T$ .
- ▶ The cocycle conditions give  $u = y$  and  $x = uv - 1$ . Note that the second equation is on the region  $x = -1 + \Lambda_+$ .
- ▶ In other words,  $\text{CF}(T, T)$  gives an extension of  $\text{CF}(S, S)$  from  $v \in \Lambda_+$  to  $v \in \Lambda_0$  with  $uv \neq 1$ .
- ▶ Similarly we can glue the immersed sphere  $S$  (with  $(u, v) \in \Lambda_+ \times \Lambda_0$ ) with a Clifford torus  $T'$ . It is  $v = y'$  and  $x' = uv - 1$ .
- ▶ The resulting moduli is  $\{(u, v) \in \Lambda_0 \times \Lambda_0 : uv \neq 1\}$ .

# Gluing




- ▶ We glue the deformations  $(u, v) \in \Lambda_0 \times \Lambda_+$  with the deformations  $(x, y) \in (\mathbf{C}^\times \oplus \Lambda_+)^2$  of  $T$ .
- ▶ The cocycle conditions give  $u = y$  and  $x = uv - 1$ . Note that the second equation is on the region  $x = -1 + \Lambda_+$ .
- ▶ In other words,  $\text{CF}(T, T)$  gives an extension of  $\text{CF}(S, S)$  from  $v \in \Lambda_+$  to  $v \in \Lambda_0$  with  $uv \neq 1$ .
- ▶ Similarly we can glue the immersed sphere  $S$  (with  $(u, v) \in \Lambda_+ \times \Lambda_0$ ) with a Clifford torus  $T'$ . It is  $v = y'$  and  $x' = uv - 1$ .
- ▶ The resulting moduli is  $\{(u, v) \in \Lambda_0 \times \Lambda_0 : uv \neq 1\}$ .

# Grassmannians



- ▶ With **Hansol Hong and Yoosik Kim**, we are applying this to construct the compactified mirror of  $\text{Gr}(2, n)$ .
- ▶ Flag manifolds have Gelfand-Cetlin systems serving as Lagrangian torus fibrations.
- ▶ Immersed spheres are important in that case because they appear as critical fibers of the superpotential.
- ▶ For instance, by **Nohara-Ueda** for  $\text{Gr}(2, 4)$  there is a critical point  $(0, 0)$  which corresponds to a certain fiber  $\mathbf{S}^3 \times \mathbf{S}^1$  of the Gelfand-Cetlin system.
- ▶ The trouble is  $\mathbf{S}^3$  is rigid!

# Grassmannians



- ▶ Consider the symplectic reduction picture on  $\mathbf{S}^2$ . We push in one singular point and consider the corresponding moduli.
- ▶ There is one monotone Lagrangian torus above and below the wall respectively. We glue the deformation spaces of an immersed sphere (times  $T^2$ ) with that of the two monotone tori like in the last slide.
- ▶ This recovers the **Rietsch** Lie theoretical mirror for  $\text{Gr}(2, 4)$ .